



# Article Estimates on Some General Classes of Holomorphic Function Spaces

Amnah E. Shammaky<sup>1</sup> and Ahmed El-Sayed Ahmed<sup>2,\*</sup>

- <sup>1</sup> Department of Mathematics, Faculty of Science, Jazan University, Jazan 45142, Saudi Arabia; aeshamakhi@jazanu.edu.sa
- <sup>2</sup> Mathematics Department, Faculty of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia
- \* Correspondence: a.elsayed@tu.edu.sa

**Abstract:** In this current manuscript, some general classes of weighted analytic function spaces in a unit disc are defined and studied. Special functions significant in both analytic  $T(p,q,m,s;\Psi)$  norms and analytic  $\Psi$ -Bloch norms serve as a framework for introducing new families of analytic classes. An application in operator theory is provided by establishing important properties of the composition-type operator  $C_{\phi}$  such as the boundedness and compactness with the help of the defined new classes.

**Keywords:** composition operators; Bloch functions;  $T(p, p - 2, m, s; \Psi)$ -classes; Carleson measure

AMS 2020 Classification: 47B33; 46E15; 30H30

## 1. Introduction

The subjects of complex classes of Banach function spaces and composition operators have gained extremely considerable popularity and high importance during the last few decades. These important subjects have mainly demonstrated significant applications and interesting, widespread, different fields of mathematical analysis. In the last few years, regarding the above-mentioned evolved motivations, some extended complex classes of function spaces associated with certain weights have been introduced and studied actively; see the related references herein. The main goal of this article is to introduce some new relevant classes of related holomorphic function spaces with the help of some general relevant weights and then establish some numerous, potentially useful interesting properties and general formulas. Some important properties of the composition operator  $C_{\phi}$  such as the boundedness and compactness are investigated using the defined general classes in the considered complex disc.

Mainly definitions, notations and basic concepts will be recalled in the next section. The considered class that contains all the analytic functions in the unit disk  $\mathbb{D} = \{w \in \mathbb{C} : |w| < 1\}$  is symbolized by  $H(\mathbb{D})$ .

**Definition 1** ([1,2]). *Let*  $g \in H(\mathbb{D})$  *and*  $\alpha \in (0, \infty)$ *. When* 

$$\|g\|_{\mathcal{B}^lpha} = \sup_{w\in\mathbb{D}}(1-|w|^2)^lpha|g'(w)| < \infty, \ w\in\mathbb{D}$$

Thus, g is said to belong to the known  $\alpha$ -Bloch space  $\mathcal{B}^{\alpha}$ . Additionally, the specific space  $\mathcal{B}^{1}$  is called the analytic Bloch space  $\mathcal{B}$ .

**Definition 2** ([3]). Let  $g \in H(\mathbb{D})$  and suppose that  $p \in (1, \infty)$ . If

$$\|g\|_{B_p}^p = \sup_{w\in\mathbb{D}} \int_{\mathbb{D}} |g'(w)|^p (1-|w|^2)^{p-2} d\xi(w) < \infty,$$



Citation: Shammaky, A.E.; El-Sayed Ahmed, A. Estimates on Some General Classes of Holomorphic Function Spaces. *Symmetry* **2021**, *13*, 528. https://doi.org/10.3390/ sym13040528

Academic Editor: Juan Luis García Guirao

Received: 6 March 2021 Accepted: 22 March 2021 Published: 24 March 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). then g belongs to the Besov space  $B_p$ .

Suppose that  $\varphi : \mathbb{D} \longrightarrow \mathbb{D}$ . Assume  $d\xi(w) = dx dy$  defines the relevant Euclidean area element on  $\mathbb{D}$ . Using  $\varphi$ , the composition operator  $C_{\varphi}$  is given by

$$C_{\varphi}g = g \circ \varphi$$

where  $g \in H(\mathbb{D})$ . This operator maps analytic functions *g* to analytic functions.

Composition operators are among the most interesting and widely studied of the different types of operators. The study of their properties (boundedness and compactness) is very important. For several studies of composition operators on some weighted classes of analytic function spaces, refer to [4–8] and others.

Let  $F : X_1 \to X_2$  be a linear operator; then, F is called compact if it maps relevant bounded sets in  $X_1$  to sets in  $X_2$ , where such sets have compact closure. For the spaces  $X_1$  and  $X_2$  of the Banach type, it is said that F is compact from  $X_1$  to  $X_2$  if each bounded sequence  $\{a_n\} \in X_1$  and the sequence  $\{Fa_n\} \in X_2$  contains a subsequence converging to some limits in  $X_2$ .

Let  $M_h$  and  $M_h^*$  be two quantities, both of them depending on the function h, for which  $h \in H(\mathbb{D})$ ; thus, we can write  $M_h \approx M_h^*$ , when have a constant  $\eta > 0$ , for which

$$\frac{1}{\eta}M_h^* \le L_g \le \eta M_h^*.$$

Let  $b \in \mathbb{D}$  and  $0 < |w| = \rho < 1$ . The symbol  $\sigma(b, \rho)$  stands for the relevant pseudo-hyperbolic complex disc such that (see [9])

$$|\sigma(b,\rho)| = \frac{(1-|b|^2)^2}{(1-\rho^2|b|^2)^2}\rho^2.$$

#### 2. Analytic Bloch Characterizations

In the present article, some properties of analytic functions belonging to  $T(p, q, m, s; \Psi)$  are studied. Additionally, certain Carleson measure characterizations of the relevant compact composition operator  $C_{\varphi}$  on  $T(p, q, m, s; \Psi)$  spaces are given.

Interesting characterizations of  $C_{\varphi}$  on  $T(p, q, m, s; \Psi)$  spaces are also presented. In this relevant study, the relevant function  $\Psi$  is supposed to be a nondecreasing bounded and continuous function, where  $\Psi : (0, 1) \to (0, \infty)$ .

**Definition 3.** For a given nondecreasing continuous function  $\Psi : (0,1) \to (0,\infty)$ , the function  $g \in H(\mathbb{D})$  is said to belong to the space  $\mathcal{B}_{\Psi}$  if

$$\|g\|_{\mathcal{B}_{\Psi}} = \sup_{w \in \mathbb{D}} \Psi(|w|) |g'(w)| < \infty.$$

**Remark 1.** In the above definition, suppose that  $\Psi(t) = (1-t)^{\alpha}$ ,  $0 < t < \infty$ ; then, we obtain the definition of an  $\alpha$ -Bloch space where  $0 < \alpha < \infty$ .

Let the Green's function  $g(w, a) = \ln \left| \frac{1 - \bar{a}w}{a - w} \right| = \ln \frac{1}{|\varphi_a(w)|}$ , where  $\varphi_a(w) = \frac{1 - \bar{a}w}{a - w}$  stands for the Möbius transformation.

By the help of the function g(w, a), the following definition can be given.

**Definition 4.** Let  $0 , <math>-2 < q < \infty$ ,  $0 < m < \infty$  and  $0 < s < \infty$ . Additionally, let q + s + m > -1. For a given nondecreasing continuous function  $\Psi : (0,1) \rightarrow (0,\infty)$ , the function  $g \in H(\mathbb{D})$  is said to belong to the space  $T(p,q,m,s;\Psi)$  if

$$\|g\|_{T(p,q,m,s;\Psi)}^{p} = \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |g'(w)|^{p} (1-|w|^{2})^{q} \Psi^{p}(|w|) g^{s}(w,a) (1-|\varphi_{a}(w)|^{2})^{m} d\xi(w) < \infty.$$
(1)

**Remark 2.** The analytic  $T(p,q,m,s;\Psi)$  function classes involve some various special analytic classes of complex-type function spaces. For example, when m = 0 and  $\Psi \equiv 1$ , the analytic F(p,q,s) classes can be obtained (see [10]). Additionally, when q = 0, m = 0, p = 2 and  $\Psi \equiv 1$ , the analytic  $Q_s$  classes can be followed ([2]). Furthermore, upon setting s = 0 and  $\Psi \equiv 1$ , the known analytic Besov-type classes are also obtained (see [3,11]).

In this current manuscript, the conditions of q + s + m > -1 and the boundedness of the weighted holomorphic function  $\Psi$  are necessary to guarantee nontrivial holomorphic spaces. Moreover, these classes of complex-type function spaces have interesting, clear importance and different applications in various research areas, for instance, in operator theory and in measure theory as well as in differential equations. In the present article, useful interesting discussions with the help of a class of composition operators on the defined analytic  $T(p, q, m, s; \Psi)$ -type classes are introduced.

**Theorem 1.** Let  $0 < |w| = \rho < 1$ ,  $0 , and <math>0 \le m < \infty$  with q + s + m > -1. Assume that  $0 \le s < \infty$ . Let  $\Psi : (0,1) \to (0,\infty)$  be a given nondecreasing bounded and continuous function. Then, for  $g \in H(\mathbb{D})$ , the next fundamental quantities are equivalent: (A)  $\|g\|_{\mathcal{B}_{\Psi}}^{p}$ 

$$\sup_{a\in\mathbb{D}}\frac{1}{|\sigma_{(a,\rho)}|^{1-\frac{p}{2}}}\int_{U(a,r)}|g'(w)|^p\Psi^p(|w|)d\xi(w)<\infty,$$

(C)

$$\sup_{a\in\mathbb{D}}\int_{\sigma_{(a,\rho)}}|g'(w)|^p(1-|w|^2)^{p-2}\Psi^p(|w|)d\xi(w)<\infty,$$

(D)

$$\sup_{a\in\mathbb{D}}\int_{\sigma_{(a,\rho)}}|g'(w)|^{p}(1-|w|^{2})^{p-2}\Psi^{p}(|w|)(1-|\varphi_{a}(w)|^{2})^{m+s}d\xi(w)<\infty,$$

(E)

$$\sup_{a\in\mathbb{D}}\int_{\sigma_{(a,\rho)}}|g'(w)|^p(1-|w|^2)^{p-2}\Psi^p(|w|)g^{m+s}(w,a)d\xi(w)<\infty,$$

(F)

$$\sup_{a\in\mathbb{D}}\int_{\sigma_{(a,\rho)}}|g'(w)|^p\left(\log\frac{1}{|w|}\right)^p\Psi^p(|w|)|\varphi_a'(w)|^2d\xi(w)<\infty$$

**Proof.** The proof can be obtained as the corresponding results in [1,12,13] with very simple modifications.  $\Box$ 

### 3. Nevanlinna-Type Functions

Composition operators on weighted classes of function spaces are intensively researched by several authors (see [4,5,14–19] and others).

Nevanlinna-type counting functions are playing a significant role in such studies. Some types of Nevanlinna counting functions for different function spaces are defined. For instance, in [4], the known Nevanlinna-type counting holomorphic function is defined to help with studying some interesting useful properties of composition-type operators on holomorphic-type F(p,q,s) classes.

Next, an alternative definition for the Nevanlinna-type counting holomorphic function will be introduced to characterize certain properties of  $C_{\varphi}$  on holomorphic  $T(p, q, m, s; \Psi)$ -type classes.

**Definition 5.** *The alternative definition of the Nevanlinna-type counting holomorphic function for the holomorphic*  $T(p,q,m,s;\Psi)$ *-type classes is given by:* 

$$L_{p,q,m,s;\omega}(w) = \sum_{\varphi(z)=w} \{ |\varphi'(z)|^{p-2} (1-|z|^2)^q \Psi^p(|z|) g^s(z,a) (1-|\varphi_a(z)|^2)^m \},$$
(2)

for  $w \in \varphi(\mathbb{D})$ ,  $p \in (2, \infty)$ ,  $q \in (-2, \infty)$ ,  $s \in (0, \infty)$  and  $0 < m < \infty$ .

**Remark 3.** In Definition 5, let  $\Psi(|z|) \equiv 1$  and m = 0; then, we obtain the Nevanlinna-type function as introduced in [4]. Additionally, if we let  $\Psi(|z|) \equiv 1$  and s = m = 0, then we obtain the counting function that is given in [20].

The alternative known Nevanlinna function is introduced by using the change of variables formula in  $T(p, q, m, s; \Psi)$  classes as follows:

For  $g \in T(p,q,m,s;\Psi)$ ,  $2 \le p < \infty$ ,  $-2 < q < \infty$ ,  $0 < s < \infty$  and  $0 < m < \infty$ ,

$$\begin{split} \|C_{\varphi}g\|_{T(p,q,m,s;\Psi)}^{p} &= \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |(g\circ\varphi)'(z)|^{p} (1-|z|^{2})^{q} \Psi^{p}(|z|) g^{s}(z,a) (1-|\varphi_{a}(z)|^{2})^{m} d\xi(z) \\ &= \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |g'(\varphi(z))|^{p} |\varphi'(z)|^{p} (1-|z|^{2})^{q} \Psi^{p}(|z|) g^{s}(z,a) (1-|\varphi_{a}(z)|^{2})^{m} d\xi(z). \end{split}$$

By using the known change in variables as obtained in [4,20], we yield

$$\|C_{\varphi}g\|_{T(p,q,m,s;\Psi)}^{p} = \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |g'(w)|^{p} L_{p,q,m,s;\Psi}(w) d\xi(w).$$
(3)

Next, there are some restrictions of  $C_{\varphi}$  to  $T(p,q,m,s;\Psi)$  that shall be clarified. Then,  $C_{\varphi}$  is a bounded operator if and only if there is a positive constant *K* such that

$$\|C_{\varphi}g\|_{T(p,q,m,s;\Psi)}^{p} \le K\|g\|_{T(p,q,m,s;\Psi)}^{p}$$

$$\tag{4}$$

for all  $g \in T(p,q,m,s;\Psi)$ .

$$\sup_{a\in\mathbb{D}}\int_{\mathbb{D}}|g'(w)|^{p}L_{p,q,m,s;\Psi}(w)d\xi(w)\leq C\|g\|_{T(p,q,m,s;\Psi)}^{p}$$

for all  $g \in T(p,q,m,s;\Psi)$ .

Suppose that  $h \in (0, 1)$ ,  $\theta \in [0, 2\pi]$ ; also let

$$\Omega(h,\theta) = \{re^{it} : r \in (1-h,1) \text{ with } h > |\theta-t|\},$$
$$S(h,\theta) = \{re^{it} : h > |re^{it} - re^{i\theta}|\}.$$

It is known that any positive-type measure  $\mu$  on  $\mathbb{D}$  is a Carleson-type measure when we find a positive  $M_1$  such that

$$\mu(S(h,\theta)) \le M_1 h$$
, for  $h \in (0,1)$  and  $\theta \in [0 \le 2\pi]$ .

Now, we clarify things by using the considered measures, which induce certain types of Carleson conditions, playing an important role in understanding how the analytic function  $\varphi$  mapping  $\mathbb{D}$  onto itself provides some types of bounded composition operators on the classes  $W = (T(p,q,m,s;\Psi) \text{ or } \mathcal{B}_{\Psi}.$ 

For various intensive and essential results on Carleson measures, see [4,5,20,21] and others.

**Definition 6.** Assume that  $\mu$  is a positive-type measure on  $\mathbb{D}$ , and  $t W = \mathcal{B}_{\Psi}$  or  $T(p,q,m,s;\Psi)$  for  $0 , <math>q \in (-2,\infty)$ ,  $s \in (0,\infty)$  and  $m \in (0,\infty)$ . Thus, the measure  $\mu$  is a (W, p)-Carleson-type measure when we find a constant C > 0; hence

$$\int_{\mathbb{D}} |g'(w)|^p d\mu(w) \le C \|g\|_W^p,$$

for all  $g \in W$ .

**Definition 7.** For  $p \in (1, \infty)$ , the specific measure  $\mu$  is called a vanishing-type p-Carleson measure when

$$\lim_{h\to 0} \sup_{\theta\in[0,2\pi)} F(\mu,h) = 0,$$

where  $F(\mu, h) = \frac{\mu(S(h,\theta))}{h^p}$ .

With the help of (2) and (3), we can deduce that the considered operator  $C_{\varphi}$  is bounded on  $T(p,q,m,s;\Psi)$  if the measure  $L_{p,q,m,s;\Psi}(w) d\xi(w)$  is a  $(T(p,q,m,s;\Psi),p)$ -Carleson measure.

Next, some global and interesting characterizations of compact composition operators on  $T(p,q,m,s;\Psi)$  classes in terms of  $p_1$ -Carleson-type measures are given.

**Theorem 2.** Let  $2 \le p < \infty$ ,  $0 < m < \infty$  and  $0 < s < \infty$ . For a given nondecreasing continuous function  $\Psi : (0,1) \to (0,\infty)$ , the next fundamental quantities can be equivalent:

(a) The measure  $\mu$  is a  $(T(p, p-2, m, s; \Psi), p)$ -Carleson-type measure,

(b) We can find a positive constant  $C_1$ , for which  $\mu(S(h,\theta)) \leq C_1 h^p$ ,  $\forall 0 < h < 1$  and  $0 \leq \theta < 2\pi$ ,

(c) We can find a specific positive constant  $C_2$  for which

$$\sup_{a\in\mathbb{D}}\int_{\mathbb{D}}|\varphi_{a}'(z)|^{p}d\mu(z) \leq C_{2}, \ \forall \ a\in\mathbb{D}.$$

Proof. Now, assume that (a) holds. Thus, by using Theorem 1 and Definition 7, we infer that

$$\int_{\mathbb{D}} |g'(z)|^p d\mu(z) \le C \int_{\mathbb{D}} |g'(z)|^p (1-|z|^2)^{p-2} \Psi^p(|z|) g^s(z,a) (1-|\varphi_a(z)|^2)^m d\xi(z),$$

for all  $g \in T(p, p-2, m, s; \Psi)$ . This can be obtained for

$$g(z) = \varphi_a(z) = \frac{a-z}{1-\bar{a}z}$$

Thus,

$$\begin{split} \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |\varphi_{a}'(z)|^{p} d\mu(z) &\leq C \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |\varphi_{a}'(z)|^{p} (1 - |z|^{2})^{p-2} \Psi^{p}(|z|) g^{s}(z, a) (1 - |\varphi_{a}(z)|^{2})^{m} d\xi(z) \\ &\leq C \|\varphi_{a}\|_{T(p, p-2, m, s; \Psi)}^{p} \leq C \text{ const.}, \end{split}$$

for all  $a \in \mathbb{D}$ . This gives (c). It is not hard to prove the equivalence between (b) and (c). Now, assume that (b) holds; we shall prove that (a) is true. For  $z = re^{i\theta}$ , let

$$\sigma_1(z) = \left\{ t : |t - z| < \frac{1 - |z|}{2} \right\},\$$
$$\sigma_2(z) = \left\{ t : |t - z| < 1 - |z| \right\}.$$

Then,

$$\sigma_1(z) \subseteq \sigma_2(z) \subseteq S(2(1-|z|),\theta).$$

Furthermore, if  $a \in \sigma_1(z)$ , then

$$\frac{1}{2} \le \frac{1-|t|}{1-|z|} \le \frac{3}{2}.$$

Let  $g \in T(p, q, m, s; \Psi)$ ; because the function g is analytic,

$$g'(z) = \frac{4}{\pi(1-|z|)^2} \int_{\sigma_1(z)} g'(t) d\xi(t).$$

By Jensen's inequality (see [22]), we obtain

$$|g'(z)|^p \leq rac{4}{\pi(1-|z|)^2} \int_{\sigma_1(z)} |g'(t)|^p d\xi(t).$$

Thus,

$$\begin{split} \int_{\mathbb{D}} |g'(z)|^{p} d\mu(z) &\leq \int_{\mathbb{D}} \frac{4}{\pi (1-|z|)^{2}} \left( \int_{\sigma_{1}(z)} |g'(t)|^{p} d\xi(t) \right) d\mu(z) \\ &\leq \frac{4}{\pi} \int_{\mathbb{D}} \left( \int_{\sigma_{1}(z)} |g'(t)|^{p} \left( \frac{3}{2(1-|t|)} \right)^{2} d\xi(t) \right) d\mu(z) \\ &\leq \frac{9}{\pi} \int_{\mathbb{D}} \int_{\mathbb{D}} \int_{\mathbb{D}} |g'(t)|^{p} \Lambda_{\sigma_{1}(z)}(t) (1-|t|)^{-2} d\xi(t) d\mu(z) \\ &\leq \frac{9}{\pi} \int_{\mathbb{D}} |g'(t)|^{p} (1-|t|)^{-2} \int_{\mathbb{D}} \Lambda_{\sigma_{1}(z)}(t) d\mu(z) d\xi(t), \end{split}$$

where  $\Lambda_{\sigma_1(z)}(t)$  stands for the characteristic function with  $\Lambda_{\sigma_1(z)}(t) \leq \Lambda_{S(2(1-|z|),\theta)}(t)$ ,  $z = re^{i\theta}$ , because  $a \in \sigma_1(z)$ , which means that

$$|t-e^{i\theta}|<2(1-|t|).$$

Now, by applying (b), we infer that

$$\int_{\mathbb{D}} \Lambda_{\sigma_1(z)} d\mu(z) \leq \mu(S(2(1-|t|),\theta)) \leq A2^p(1-|t|)^p.$$

Therefore,

$$\begin{split} \int_{\mathbb{D}} |g'(z)|^p d\mu(z) &\leq \frac{9}{\pi} A 2^p \int_{\mathbb{D}} |g'(t)|^p (1-|t|)^{p-2} d\xi(t) \\ &\leq C \int_{\mathbb{D}} |g'(t)|^p (1-|t|)^{p-2} d\xi(t), \end{split}$$

where  $C_1$  is a positive constant. From Theorem 1, the statements (C) and (E) are equivalent; hence,

$$\int_{\mathbb{D}} |g'(z)|^p d\mu(z) \leq C \int_{\mathbb{D}} |g'(t)|^p (1-|t|)^{p-2} \Psi^p(|z|) g^s(z,a) (1-|\varphi_a(z)|^2)^m d\xi(t)$$
  
 
$$\leq C \|g\|_{T(p,p-2,m,s;\Psi)},$$

which is (a). The proof is thereby finished.  $\Box$ 

Using Theorem 2, the following result can be proved.

**Theorem 3.** Let  $\varphi$  be an analytic function on  $\mathbb{D}$ ,  $2 \le p < \infty$ ,  $0 < m < \infty$  and  $0 < s < \infty$ . Let  $\Psi : (0,1) \to (0,\infty)$  be a given nondecreasing continuous function. Then, the operator  $C_{\varphi}$  is bounded on  $T(p, p - 2, m, s; \Psi) \iff$ 

$$\sup_{a\in\mathbb{D}}\|C_{\varphi} \ \varphi_a\|_{T(p,p-2,m,s;\Psi)}<\infty.$$

Next, some interesting results on  $T(p, q, m, s; \Psi)$  spaces are proved.

**Lemma 1.** Let  $\Psi$  :  $(0,1) \rightarrow (0,\infty)$  be a given nondecreasing bounded and continuous function. Let  $0 , <math>0 < q < \infty$ ,  $0 < s < \infty$   $0 < m < \infty$ . Suppose that

$$\int_0^1 \frac{r\,dr}{\Psi(r)} < \infty$$

Then:

(*i*) Every bounded sequence  $(h_n) \subset T(p,q,s,m;\Psi)$  is converging, uniformly bounded in compact specific sets.

(*ii*) For any specific sequence  $(h_n)$  on  $T(p,q,s,m;\Psi)$ ,  $||h_n||_{T(p,q,s,m;\Psi)} \to 0$  by the uniform convergence  $h_n - h_n(0) \to 0$  on compact sets.

**Proof.** From Theorem 1, we have

$$\|g\|_{\mathcal{B}_{\Psi}} \leq \lambda_1 \|g\|_{T(p,q,m,s;\Psi)},$$

 $\lambda_1$  is a positive constant. If  $z \in U(0, r)$ , 0 < r < 1, then we get

$$|h_n - h_n(0)| = \left| \int_0^1 h'_n(zt) z dt \right| \le \lambda_1 ||h_n||_{\mathcal{B}_{\Psi}} \le \lambda ||h_n||_{T(p,q,m,s;\Psi)}.$$

where  $\lambda$  and  $\lambda_1$  are positive constants. The lemma is therefore completely established.  $\Box$ 

**Lemma 2.** Let  $\Psi : (0,1) \to (0,\infty)$  be a given nondecreasing continuous function. Suppose that  $2 \le p < \infty, 0 < q < \infty, 0 < m < \infty$ , and  $0 < s < \infty$ . Let  $X_1, X_2 = T(p,q,m,s;\Psi)$  or  $\mathcal{B}_{\Psi}$ . Then,

 $C_{\varphi}: X_1 \to X_2$  is a compact operator  $\iff \|C_{\varphi}g_n\|_{X_2} \to 0$ 

where  $(g_n) \subset X_1$  is a bounded sequence converging uniformly to zero on compact sets as  $n \to \infty$ ,  $g_n \to 0$ .

**Proof.** For the spaces  $T(p, q, m, s; \Psi)$  or  $\mathcal{B}_{\Psi}$ , the quantities (i) and (ii) can be proved in view of Lemma 2.10 [20]. From Lemma 1, it is not hard to clarify that (i) holds. To clarify that the quantity (ii) holds, assume that  $(g_n)$  is a sequence in the closed unit ball of  $X_1$ . Then, by Lemma 1, the sequence  $(g_n)$  is uniformly bounded on compact sets. Applying Montel's theorem (see [23]), we can find a subsequence  $(g_{n_k}), (n_1 < n_2 < ...)$  such that  $g_{n_k} \to f$  uniformly on compact sets, for some  $f \in H(\mathbb{D})$ . Now, we clarify that  $f \in X_1$ .

(i) If  $X_1 = T(p,q,m,s; \Psi)$ ,  $(2 \le p < \infty, -2 < q < \infty, 0 < m < \infty, 0 < s < \infty)$ , then by applying Fatou's theorem, we deduce that

$$\begin{split} \int_{\mathbb{D}} |g'(z)|^p S(\Psi, g, \varphi_a) \, d\xi(z) &= \int_{\mathbb{D}} \lim_{k \to \infty} |g'_{n_k}(z)|^p S(\Psi, g, \varphi_a) \, d\xi(z) \\ &\leq \liminf_{k \to \infty} \int_{\mathbb{D}} |g'_{n_k}(z)|^p S(\Psi, g, \varphi_a) \, d\xi(z), \\ &= \liminf_{k \to \infty} \|g_{n_k}\|_{T(p,q,m,s;\Psi)}^p < \infty, \end{split}$$

where

$$S(\Psi, g, \varphi_a) = (1 - |z|^2)^q \Psi^p(|z|) g^s(z, a) (1 - |\varphi_a(z)|^2)^m$$

(ii) If  $X_1 = \mathcal{B}_{\Psi}$  , we get

$$|g'(z)|(1-|z|^2)\Psi(|z|) = \lim_{k\to\infty} |g'_{n_k}(z)|\Psi(|z|)(1-|z|^2) \leq \lim_{k\to\infty} \|g'_{n_k}\|_{\mathcal{B}_{\Psi}} < \infty,$$

Lemma 1 gives

 $C_{\varphi}: X_1 \to X_2$  is a compact operator  $\iff \|C_{\varphi}g_n\|_{X_2} \to 0$ 

where the sequence  $(g_n) \subset X_1$  is a bounded specific sequence that converges uniformly to zero on compact sets as  $n \to \infty$ . Hence,  $|g_n(\varphi(0))| + ||C_{\varphi}g_n||_{X_2} \to 0$ , as  $n \to \infty$ . The lemma is therefore proved.

For the compactness of the operator  $C_{\varphi}$  on  $T(p,q,m,s;\Psi)$  spaces, the next results can be proved.  $\Box$ 

**Theorem 4.** Let  $\Psi : (0,1) \to (0,\infty)$  be a given nondecreasing continuous function. Let  $2 \le p < p_1 < \infty, 0 < q < \infty$ , and  $0 < s < \infty$ . Then, the next essential quantities are equivalent:

(a) The operator  $C_{\varphi}$ :  $T(p,q,m,s;\Psi) \rightarrow T(p_1,q,m,s;\Psi)$  is compact.

- (b) The measure  $L_{p_1,q,m,s;\Psi}(w)d\xi(w)$  is a vanishing  $p_1$ -Carleson measure.
- (c) The norm  $\|C_{\varphi} \ \varphi_a\|_{T(p_1,q,m,s;\Psi)} \to 0$  as  $|a| \to 1$ .

**Proof.** Using (3), we infer that

$$\|C_{\varphi} \varphi_a\|_{T(p_1,q,m,s;\Psi)}^{p_1} = \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |\varphi_a'(w)|^{p_1} L_{p_1,q,m,s;\Psi}(w) d\xi(w)$$

Using Lemma 2.1 in [24], we have (b) $\iff$ (c). Next, we show that (a) $\implies$ (c). Let us assume that

$$C_{\varphi}: T(p,q,m,s;\Psi) \rightarrow T(p_1,q,m,s;\Psi)$$

is a specific compact operator. Because the set  $\{\varphi_a : a \in \mathbb{D}\}$  is bounded in classes  $T(p,q,m,s;\Psi)$ ,

$$\|\varphi_a\|_{T(p,q,m,s;\Psi)} = \|z \circ \varphi_a\|_{T(p,q,m,s;\Psi)},$$

with the norm of  $\varphi_a$  in  $T(p, q, m, s; \Psi)$  such that

$$\|\varphi_a(0)\| + \|\varphi_a\|_{T(p,q,m,s;\Psi)} < 1 + \|\varphi_a\|_{T(p,q,m,s;\Psi)} < \infty.$$

Additionally, the convergence  $(\varphi_a - a) \longrightarrow 0$  as  $|a| \rightarrow 1$  is uniform on specific compact sets, since

$$|\varphi_a - a| = |z| \frac{1 - |a|^2}{|1 - \bar{a}z|}, \text{ for } r = |z|; z \in \mathbb{D}.$$

Hence, by Lemma 2, we obtain

$$\|C_{\varphi}(\varphi_a - a)\|_{T(p_1, q, m, s; \Psi)} \to 0$$
, as  $|a| \to 1$ .

Now, we clarify that (b) $\Rightarrow$ (a). Let  $(g_n)$  be a certain bounded sequence in  $T(p,q,m,s;\Psi)$  that converges to 0 uniformly on compact sets. Using the mean-value property for the  $g'_n$ , shows that

$$g'_{n}(\zeta) = \frac{4}{\pi(1-|\zeta|)^{2}} \int_{|\zeta-z| < \frac{1-|\zeta|}{2}} |g'_{n}(z)| d\zeta(z).$$
(5)

Jensen's inequality gives

$$\sigma_{1}(\zeta) = \left\{ z : |\zeta - z| < \frac{1 - |\zeta|}{2} \right\},\$$
$$|g_{n}'(\zeta)|^{p_{1}} \le \frac{4}{\pi (1 - |\zeta|)^{2}} \int_{\sigma_{1}(\zeta)} |g_{n}'(z)|^{p_{1}} d\zeta(z).$$
(6)

Thus, by using (6) and applying the known Fubini's theorem, the following inequality can be deduced

,

$$\begin{split} \|C_{\varphi}g_{n}\|_{T(p_{1},q,m,s;\Psi)}^{p_{1}} &= \sup_{a\in\mathbb{D}}\int_{\mathbb{D}}|g_{n}'(\zeta)|^{p_{1}}L_{p_{1},q,m,s;\Psi}(\zeta)d\xi(\zeta) \\ &\leq \sup_{a\in\mathbb{D}}\int_{\mathbb{D}}\frac{4}{\pi(1-|\zeta|)^{2}}\left(\int_{\sigma_{1}(\zeta)}|g_{n}'(z)|^{p_{1}}d\xi(z)\right)L_{p_{1},q,m,s;\Psi}(\zeta)d\xi(\zeta). \end{split}$$

Then,

$$\|C_{\varphi}g_{n}\|_{T(p_{1},q,m,s;\Psi)}^{p_{1}} \leq \frac{4}{\pi} \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |g_{n}'(z)|^{p_{1}} \left( \int_{\mathbb{D}} \frac{\Lambda_{\sigma_{1}(\zeta)}(z)}{(1-|\zeta|)^{2}} L_{p_{1},q,m,s;\Psi}(\zeta) d\xi(\zeta) \right) d\xi(z), \quad (7)$$

When  $|\zeta - z| < \frac{1-|\zeta|}{2}$ ,  $\zeta \in S(2(1-|z|), \theta)$ , where  $z = |z|e^{i\theta}$ , since

$$|\zeta - e^{i\theta}| \le |z - \zeta| + |e^{i\theta} - z| < \frac{1 - |\zeta|}{2} + |\frac{z}{|z|} - z| < 2(1 - |z|).$$

Hence, when  $|\zeta - z| < \frac{1-|w|}{2}$ ,

$$\frac{1}{(1-|\zeta|)^2} < c_1 \frac{1}{(1-|z|)^2},$$

where  $c_1$  is a positive specific constant. Then,

$$\begin{split} \|C_{\varphi}g_{n}\|_{T(p_{1},q,m,s;\Psi)}^{p_{1}} &\leq c_{1} \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} \frac{|g_{n}'(z)|^{p_{1}}}{(1-|z|)^{2}} \left(\int_{S(2(1-|z|),\theta)} L_{p_{1},q,m,s;\Psi}(\zeta) \, d\zeta(\zeta)\right) d\zeta(z) \\ &= c_{2} \sup_{a\in\mathbb{D}} \left(\int_{|z|>1-\frac{\delta}{2}} + \int_{|z|\leq 1-\frac{\delta}{2}} \frac{|g_{n}'(z)|^{p_{1}}}{(1-|z|)^{2}} \left(\int_{S(2(1-|z|),\theta)} L_{p_{1},q,m,s;\Psi}(b) d\zeta(\zeta)\right) d\zeta(z) \right) \\ &= c_{3} \sup_{a\in\mathbb{D}} (J+J^{*}) \end{split}$$

for any  $0 < \delta < 1$ , where  $c_2$  and  $c_3$  are specific positive constants. By fixing  $\varepsilon > 0$  with  $\delta > 0$  for which  $\ heta \in [0,2\pi]$  with  $\ h_1 < \delta$  ,

$$\sup_{a\in\mathbb{D}}\int_{S(h,\theta)}L_{p_{1},q,m,s,\varphi;\Psi}(w)d\xi(w) \leq \varepsilon h_{1}^{p_{1}}.$$
(8)

Therefore, we deduce that

$$J \leq \varepsilon 2^{p_1} \int_{|z|>1-\frac{\delta}{2}} \frac{|g'_n(z)|^{p_1}}{(1-|z|)^2} (1-|z|)^{p_1} d\xi(z)$$
  
$$\leq \varepsilon 2^{p_1} \int_{|z|>1-\frac{\delta}{2}} |g'_n(z)|^{p_1} (1-|z|)^{p_1-2} d\xi(z)$$
  
$$\leq \varepsilon c_4 ||g_n||_{\mathcal{B}^{p_1}} < \varepsilon c_5,$$

where  $c_4$  and  $c_5$  are positive constants.

Additionally, we have

$$J^{*} \leq k \sup_{a \in \mathbb{D}} \int_{|z| \leq 1 - \frac{\delta}{2}} \frac{|g'_{n}(z)|^{p_{1}}}{(1 - |z|)^{2}} \left( \int_{S(2(1 - |z|), \theta)} L_{p_{1}, q, s, m; \Psi}(w) d\xi(w) \right) d\xi(z)$$
  
=  $k_{1} \sup_{a \in \mathbb{D}} \left( \int_{\mathbb{D}} L_{p_{1}, q, s, m, \Psi}(w) d\xi(w) \right) \int_{|z| \leq 1 - \frac{\delta}{2}} |g'_{n}(z)|^{p_{1}} d\xi(z) < const.,$ 

where  $k, k_1$  and  $k_2$  are positive constants. For the case when n is large enough, because the convergence type of  $g'_n \to 0$  is uniform on considered compact sets, the following inequality can be deduced

$$\|C_{\varphi}g'_{n}\|_{T(p_{1},q,m,s;\Psi)}^{p_{1}} < c^{*} \sup_{a \in \mathbb{D}}(J+J^{*}) < \varepsilon \quad const.,$$

for a larger *n*, where  $c^*$  is a specific positive constant. Then,  $\|C_{\varphi}g'_n\|_{T(p_1,q,m,s;\Psi)}^{p_1} \to 0$ , when  $n \to \infty$ , so Lemma 2 implies that

$$C_{\varphi}: T(p,q,m,s;\Psi) \rightarrow T(p_1,q,m,s;\Psi)$$

is an actually compact operator. The proof is thereby clearly established.  $\Box$ 

**Remark 4.** For our current study, when  $\varphi(w) = w$ , the composition operator  $C_{\varphi}$  is not a compact operator. The condition is  $\varphi(e^{i\theta}) < 1$  (almost everywhere). Thus, the operator  $C_{\varphi}$  is compact whenever

$$\int_0^{2\pi} \frac{d\theta}{1-\varphi(e^{i\theta})} < \infty.$$

#### 4. Conclusions

This manuscript covers a global study of a class of composition operators acting between a new general family of analytic function spaces on a disk  $\mathbb{D}$ . The elementary properties of the new class of functions are characterized with a class of specific univalent functions.

Motivated by a nondecreasing and continuous function as well as the Green's function, some certain weighted holomorphic classes of functions are defined and carefully discussed in the present manuscript. Important and global characterizations for the weighted classes  $T(p, q, m, s; \Psi)$  with the general weighted Bloch-type class  $\mathcal{B}_{\Psi}$  are introduced. Concerning the defined classes, the values of all the parameters have significant and large effects in the obtained results. Some interesting specific properties for the class of composition operators are discussed. The essential width for the operator  $C_{\varphi}$  is clarified between the defined weighted classes. The compactness and boundedness properties of the operator  $C_{\varphi}$  are discussed with the help of Carleson-type measure roles as well as the alternative considered Nevanlinna function.

For future research on the classes  $T(p,q,m,s;\Psi)$ , we make the following remarks.

**Remark 5.** How can we establish some properties for other classes of operators, such as superposition and integral operators on the analytic  $T(p, q, m, s; \Psi)$  classes?

**Remark 6.** There are some interesting recent studies on hyperbolic-type spaces (see [25,26]). Thus, it is possible to discuss some problems by using hyperbolic  $T(p,q,m,s;\Psi)$ -type classes.

**Remark 7.** Generalizing the analytic  $T(p, q, m, s; \Psi)$ -type spaces to the known quaternion analysis can be considered a future open problem, too. For some numerous recent studies with the help of quaternion techniques, one may refer to [27–30] and others.

**Author Contributions:** Methodology, A.E.S. and A.E.-S.A.; formal analysis, A.E.S. and A.E.-S.A.; investigation, A.E.S. and A.E.-S.A.; data curation, A.E.S. and A.E.-S.A.; writ-ing—original draft preparation, A.E.S. and A.E.-S.A.; writing—review and editing, A.E.S. and A.E.-S.A.; visualization, A.E.S. and A.E.-S.A.; project administration, A.E.S. and A.E.-S.A. Each of author has contributed to the current study equally. All authors have read and agreed to the published version of the manuscript.

Funding: There was no external funding.

Data Availability Statement: There are no data supporting this manuscript.

Acknowledgments: Ahmed El-Sayed thanks Taif University researchers for supporting the current important study with project number TURSP-2020/159, Taif University, Saudi Arabia. The authors deeply thank the reviewers for their valuable remarks and interesting comments.

Conflicts of Interest: The authors declare that both of them have no competing interests.

#### References

- 1. Rashwan, R.A.; El-Sayed Ahmed, A.; Kamal, A. Some characterizations of weighted Bloch space. *Eur. J. Pure Appl. Math.* **2009**, *2*, 250–267.
- 2. Xiao, J. Holomorphic Q Classes; Springer LNM 1767: Berlin, Gremany, 2001.
- 3. Stroethoff, K. Besov-type characterizations for the Bloch space. Bull. Austral. Math. Soc. 1989, 39, 405–420. [CrossRef]
- 4. El-Sayed Ahmed, A.; Bakhit, M. Composition operators on some holomorphic Banach function spaces. *Math. Scand.* 2009, 104, 275–295. [CrossRef]
- 5. El-Sayed Ahmed, A.; Bakhit, M. Composition operators acting between some weighted Möbius invariant spaces. *Ann. Funct. Anal.* **2011**, *2*, 138–152. [CrossRef]
- Tang, S.; Wu, P. Composition operator, boundedness, compactness, hyperbolic Bloch-type space B<sup>\*</sup>, hyperbolic-type space. J. Funct. Spaces 2020, 23, 5390732. [CrossRef]
- 7. Tien, P.T.; Khoi, L.H. Weighted composition operators between different Fock spaces. Potential Anal. 2019, 50, 171–195. [CrossRef]
- Tien, P.T.; Khoi, L.H. Differences of weighted composition operators between the Fock spaces. *Monatsh. Math.* 2019, 188, 183–193. [CrossRef]
- 9. Garnett, J.B. Bounded Analytic Functions; Academic Press: New York, NY, USA, 1981.
- 10. Zhao, R. On a General Family of Function Spaces; Suomalainen Tiedeakatemia: Helsinki, Finland, 1996; pp. 1–56.
- 11. El-Sayed Ahmed, A.; Youssif, M. Classes of weighted tent function spaces and mixed norms with some applications. *Ital. J. Pure Appl. Math.* **2020**, *43*, 402–415.
- 12. Rashwan, R.A.; El-Sayed Ahmed, A.; Kamal, A. Integral characterizations of weighted Bloch spaces and  $Q_{K,\omega}(p,q)$  spaces. *Math. Cluj.* **2009**, *51*, 63–76.
- 13. Zhao, R. On *α*-Bloch functions and VMOA. *Acta Math. Sci.* **1996**, *61*, 349–360. [CrossRef]
- 14. Colonna, F. Weighted composition operators between and BMOA. Bull. Korean Math. Soc. 2013, 50, 185–200. [CrossRef]
- 15. Cowen, C.C.; Maccluer, B.D. Composition Operators on Spaces of Analytic Functions. In *Studies in Advanced Mathematics*; CRC Press: Boca Raton, FL, USA, 1995.
- 16. Liang, Y. New characterizations for differences of weighted differentiation composition operators from a Bloch-type space to a weighted-type space. *Period. Math. Hung.* **2018**, *77*, 119–138. [CrossRef]
- 17. Li, H.; Ma, T. Generalized composition operators from  $\mathcal{B}_{\mu}$  spaces to  $Q_{K,\omega}(p,q)$  spaces. *Abstr. Appl. Anal.* **2014**, 2014, 6.
- 18. Manhas, J.S.; Zhao, R. New estimates of essential norms of weighted composition operators between Bloch type spaces. *J. Math. Anal. Appl.* **2012**, *389*, 32–47. [CrossRef]
- 19. Thompson, D. Normaloid weighted composition operators on H<sup>2</sup>. J. Math. Anal. Appl. 2018, 467, 1153–1162. [CrossRef]
- 20. Tjani, M. Compact Composition Operators on Some Möobius Invariant Banach Spaces. Ph.D. Thesis, Michigan State University, East Lansing, MI, USA, 1996.
- Essén, M.; Wulan, H.; Xiao, J. Several Function- theoretic aspects characterizations of Möbius invariant Q<sub>K</sub>-spaces. *J. Funct. Anal.* 2006, 230, 78–115. [CrossRef]
- 22. Bakula, M.K.; Matić, M.; Pécarić, J. On some general inequalities related to Jensens inequality. In *International Series of Numerical Mathematics*; Birkhäuser Verlag: Basel, Switzerland, 2008; Volume 157, pp. 233–243.
- 23. Conway, J. Functions of One Complex Variable I; Springer: Berlin, Germany, 1978; ISBN 0-387-90328-3.
- 24. Aulaskari, R.; Stegenga, D.A.; Xiao, J. Some subclasses of BMOA and their characterization in terms of Carleson measures. *Rocky Mt. Math.* **1996**, *26*, 485–506. [CrossRef]
- 25. El-Sayed Ahmed, A. Refinements on some classes of complex function spaces. Symmetry 2021, 13, 339. [CrossRef]
- 26. El-Sayed Ahmed, A.; Shammaky, A.E. Certain classes of operators on some weighted hyperbolic function spaces. *J. Math.* **2021**, 2021, 6664398. [CrossRef]
- 27. El-Sayed Ahmed, A. Hyperholomorphic Q-classes. Math. Comput. Model. 2012, 55, 1428–1435. [CrossRef]
- 28. El-Sayed Ahmed, A.; Ahmadi, A. On weighted Bloch spaces of quaternion-valued functions. AIP Conf. Proc. 2011, 1389, 272–275.

- 29. El-Sayed Ahmed, A.; Gürlebeck, K.; Reséndis, L.F.; Tovar, L.M. Characterizations for the Bloch space by *B<sup>p; q</sup>* spaces in Clifford analysis. *Complex Var. Elliptic Equ.* **2006**, *l51*, 119–136. [CrossRef]
- 30. El-Sayed Ahmed, A.; Omran, S. Extreme points and some quaternion valued functions in the Unit Ball of ℝ<sup>3</sup>. *Adv. Appl. Clifford Algebr.* **2018**, *28*, 31. [CrossRef]