

Article

Frame Transformation Relations and Symmetry Analysis of Fluxional Symmetric Rotor Dimers

Horace T. Crogman ^{1,*}, Bumgyunmiga Choi ¹, Harrison B. Chen ¹ and William G. Harter ²

¹ Physics Department, La Sierra University, Riverside, CA 92515, USA;
E-Mails: migachoi@gmail.com (B.C.); harrison8024@hotmail.com (H.B.C.)

² Physics Department, University of Arkansas, Fayetteville, AR 72701, USA;
E-Mail: wharter@urak.edu

* Author to whom correspondence should be addressed; E-Mail: hcrogman@gmail.com;
Tel: +1-951-785-2272; Fax: +1-951-785-2215.

Received: 8 November 2012; in revised form: 14 January 2013 / Accepted: 17 January 2013 /
Published: 5 February 2013

Abstract: The theory of Frame transformation relations between the states of Born Oppenheimer and the weak coupling approximations is developed for polyatomic molecules. The symmetry relations are a generalization of the frame transformation relations derived by Harter and Crogman for coupled rotor molecules. A key internal symmetry label (named “soul”) is defined so that it remains a constant label for frame transformation relations, and is conserved during vibronic transitions, ionization, and even dissociation provided the nuclear spin-rotation interaction is relatively small. Simplified procedures are given for obtaining selection rules, statistical weights, and matrix elements of multipole operators for common molecules having various point symmetries.

Keywords: frame; transformation; cluster; dimers; spin; rotation; fluxional

1. Introduction

It is well known that permutations or symmetric groups have an important role in the physics of many-body systems. Their importance stems from the fact that a system of identical particles has permutational symmetry and there exists an inter-relation between the permutation and the unitary groups. It is the theory of group representations that physicists use to solve and understand a variety of problems; the work of Frobenius in the 19th century and others such as Young [1], Yamanouchi [2,3],

Littlewood [4], and Gel'fand [5–7] have established the theory of permutational group representations as a powerful tool in spite of the fact that the theory is quite difficult. With the aid of computers, the group representations theory has become a remarkably effective tool for the modern physicist.

The theory of permutation group representations is quite useful to describe many phenomenon of molecules, ions, and radicals which are made of identical points like particles (electrons, nuclei, atoms, and molecular sub-complexes such as CH_3). Algorithms for representations, dimensions and characters that were developed by Shur [8], Littlewood [4], Yamanouchi [2,3], Robinson [9], Coleman [10], Harter [11] and others act as a powerful tool to explore the symmetry dynamics of molecular systems. Symmetry operations involving interchange of identical particles by rotations and reflections are the reasons that finite point groups such as D_m , C_m , O_h and T_d are important symmetries.

Experience has taught us that elementary quantum objects with the same quantum numbers are indistinguishable and mathematical description of multi-quantum states must not lead to a new quantum state due to a permutation of constituents. Furthermore, Fermi-Pauli and Bose-Einstein's permutational symmetry principles strongly restrict the allowed quantum configurations, spectra, and spin states. This results ultimately in having only the particular constituents in the periodic atomic chart, the chart of the Nuclides, and finally in the allowed elementary particles of the standard quark QCD.

Harter *et al.* [12] have developed an approximate theory for high J spin-rotational levels and very high resolution spectra of octahedral XY_6 molecules. Additionally, Harter and Patterson [13] were among the first to apply tableaux to molecular spin states and statistical weight calculations. The spin-statistics theorem says that the systems with integer spins are bosons, that is, symmetric to all permutations while the systems with half-integer spins are fermions, that is, antisymmetric to odd permutations. This theorem has a rather difficult proof based on the PCT theorem [14] and quantum field theory. This symmetry principle will influence all other symmetries defined for multi-particle states, for example, the possible values of the total angular momentum of molecules as discussed below. The effects of Spin-statistics can be rather subtle, one example is the influence of nuclear spins on the quantum numbers of molecules which is seen in the difference between ortho- and para-hydrogen. In the ortho form, the two proton spins are parallel, but in the para form they are paired to a total spin of 0. As a result of the fermion characteristic of the constituent protons, the allowed rotational degeneracies of the two forms are different, thus giving different heat capacities. The changes in the proton spin states are normally very slow, making the two forms appear as two macroscopically different gases.

Level and spectral clusters, and related tunneling dynamics are intimately related to nuclear, atomic, or molecular permutations. However, floppy molecules do operations that transcend the geometry of 3D crystal point group operations. One of the earliest attempts of permutation operations to non-rigid molecular symmetry was put forward by Longuet-Higgins [15]. He classified the spin states and states motions for molecular symmetry systems such as ethane and hydrazine. Longuet-Higgins points out that there are certain symmetry elements of non-rigid molecules; for example, the twisting of ethane which leads to configurations which cannot be brought into near-coincidence with the original one by a simple rotation of the whole molecule in space. The work of Hougen *et al.* [16], Harter *et al.* [17–22], and Bunker and Jensen [23] have allowed for an easier approach to compute spin weight, and rovibration wavefunctions, and to open new avenues to study large amplitude motions. In one

approach, Groner [24] has tried to identify pinwheel rotations with crystal space groups, that is, point groups with translations. Harter and Patterson pointed out that the eigenstates of the system must be based off of the irreducible representation of S_m . This means that we can use the Young tableau labeling schemes to label the eigenstates of the systems.

In this paper, we will apply the techniques developed and used by Harter and Patterson [17–22]. The main goal is to attempt to answer the question of how molecular species come about due to some kind of coupling interactions in the strongly and weakly coupled regimes, which allows for insight into large amplitude motion of molecular systems. In this paper, Section 2 extends the frame transformation relation for couple rotor systems by Crogman and Harter [25] to include symmetry. A procedure for finding the induced symmetry that resulted from various coupled states is there developed; Section 3 shows how to Young Tableau algebra to compute spin weight for the permutation group; Section 4 provides a concrete example to illustrate our technique, and Section 5 discusses the various ways to couple two diatomics and compute the resulting tunneling matrix. It is hoped that the present work will be another step in this direction for a large number of molecular systems, which have some sort of geometrical symmetry in their structure. This work is meant to expose the simplest and most powerful symmetry analysis methods available.

2. Symmetry Relation for a Coupled System

Fano and Chang [26], and Junger *et al.* [27,28] have worked out a theory that describes the phenomenon that comes about for a diatomic molecule coupled to an electron. Harter *et al.* [12] have extended their idea to polyatomic molecules. Their work went a step further by considering the symmetry induced as a result of a polyatomic molecule interacting with an electron [12]. By using Young tableau's mathematics, Harter *et al.* [12] reduced the mathematical complexity involved in the symmetry analysis. This provided a convenient way to calculate the spin weight and predict what symmetry species were excluded. Spin weights give spectral intensity ratios for different symmetry species. Here, we consider the permutation group along with Young tableau techniques [11,12,17–22,29,30] for the symmetry of two coupled diatomic molecules. Lemus [31] applied the theory of Chen [29] to generate what is known as vibrational symmetry adapted bases. This method works because of the isomorphism between the symmetric group and point group. Our exploitation of two coupled diatomic molecules allows us to utilize the power of Young tableau for the symmetry group and projector analysis as described by Harter *et al.* [11,12,23,29]. First, we imagine more general cases such as listed in tables of Figure 4.4 as illustrated in a previous article by Crogman and Harter [25].

2.1. Symmetry Effects for Two Coupled Rotors

The symmetry of any rotor can be written in terms of the \mathfrak{S} (external) $\times \bar{\mathcal{G}}$ (internal). If the rotor is free, then \mathfrak{S} is R_3 . However, if it is confined to some crystal matrix, then \mathfrak{S} is the group site symmetry of the lattice. When the rotor experiences a homogeneous electric field, the external symmetry \mathfrak{S} is O_2 . On the other hand, $\bar{\mathcal{G}}$ is the internal symmetry of the rotor structure. The fact that a rotor is made up of nuclear points means that the real internal symmetry is finite and related to permutations groups. An

external field applied to the orbitals will split into various finite symmetry species according to the molecular version of the crystal field theory.

We will start with the following expressions and develop a base transformation going from a lab weakly correlated (LWC) momentum states to a Body Oriented Angular (BOA) momentum constricted states of two strongly coupled rotors [25,32]. A bare rotor with finite symmetry (B, b) is given in [25] is defined in terms of symmetric rotor waves $D_{m n}^N$ as follows:

$$r_{m b}^{NB*}(\phi\theta\chi) = \sum_n \binom{N}{n} \binom{B}{b}^* D_{m n}^{N*}(\phi\theta\chi) \sqrt{[N]} \quad (1a)$$

where N is the total angular momentum label of rotational group R_3 in three dimensions and $[N] = \sqrt{2N+1}$.

The inverse relation as defined in [25] is given by the unitarity of the internal “crystal field” coefficient $\binom{N}{n} \binom{B}{b}$.

$$\sqrt{[N]} D_{m n}^{N*}(\phi\theta\chi) = \sum_{Bb} \binom{N}{n} \binom{B}{b} r_{m b}^{NB}(\phi\theta\chi)^* \quad (1b)$$

Let us now consider two rotors of finite symmetry labels (D^*, d) for “little” rotor D^ℓ and (B^*, b) for “main” rotor D^R . Consequently we can use Equations (1a) and (1b) to write the following (LWC) coupled wave function.

$$\begin{aligned} \langle [R\ell]_d^{D^* B^* J} \rangle &= \sqrt{[R][\ell]} \sum_{m_\ell m_R} C_{m_\ell m_R}^{\ell R J} r_{m_\ell d}^{\ell D^*}(\phi\theta\chi) r_{m_R b}^{R B^*}(\alpha\beta\gamma) \\ &= \sqrt{[R][\ell]} \sum_{n_\ell n_R} \binom{\ell}{n_\ell} \binom{D}{d}^* \binom{R}{n_R} \binom{B}{b}^* \sum_{m_\ell m_R} C_{m_\ell m_R}^{\ell R J} D_{m_\ell n_\ell}^{\ell D^*}(\phi\theta\chi) D_{m_R n_R}^{R B^*}(\alpha\beta\gamma) \end{aligned} \quad (2)$$

Using the relationship

$$\begin{aligned} \langle \alpha\beta\gamma; \phi\theta\chi |_{n_R n_\ell M}^{R \ell J} \rangle &= \sum_{m_\ell m_R} C_{m_\ell m_R}^{\ell R J} D_{m_\ell n_\ell}^{\ell D^*}(\phi\theta\chi) D_{m_R n_R}^{R B^*}(\alpha\beta\gamma) \\ &= \sum_{K\Lambda} C_{\Lambda}^{\ell R J} D_{\Lambda n_\ell}^{\ell D^*}(\Phi\Theta\Xi) D_{M K}^{J B^*}(\alpha\beta\gamma) \\ &= \left\langle \Phi\Theta\Xi(\alpha\beta\gamma) \left| \begin{matrix} \ell \\ m_\ell, n_\ell M, K = n_R + m_\ell \end{matrix} \right. \right\rangle \end{aligned} \quad (3)$$

derived in a previous work [25] to have transformation the BOA

$$\langle [R\ell]_d^{D^* B^* J} \rangle = \sqrt{[R][\ell]} \sum_{n_\ell n_R} \binom{\ell}{n_\ell} \binom{D}{d}^* \binom{R}{n_R} \binom{B}{b}^* \sum_{K\Lambda} C_{\Lambda}^{\ell R J} D_{\Lambda n_\ell}^{\ell D^*}(\Phi\Theta\Xi) D_{M K}^{J B^*}(\alpha\beta\gamma) \quad (4)$$

Following the crystal field relation, Equation (1b) gives the following:

$$\begin{aligned} \langle [R\ell]_d^{D^* B^* J} \rangle &= \sqrt{\frac{[R]}{[J]}} \sum_{Cc} \sum_{Aa} \binom{\ell}{n_\ell} \binom{D}{d}^* \sum_{K\Lambda n_R} C_{\Lambda}^{\ell R J} \binom{\ell}{n_\ell} \binom{A}{a} \binom{R}{n_R} \binom{B}{b}^* \binom{J}{K} \binom{C}{c} r_{\Lambda a}^{\ell A^*}(\Phi\Theta\Xi) r_{M c}^{J C^*}(\alpha\beta\gamma) \end{aligned} \quad (5)$$

Now, an isoscalar coefficient $\langle AB|C \rangle$ is defined using the finite symmetry Wigner Eckart theorem and coupling coefficient $C_a^A \begin{smallmatrix} B \\ b \end{smallmatrix} \begin{smallmatrix} C \\ c \end{smallmatrix}$.

$$\sqrt{\frac{[N]}{[J]}} \sum_{mnM} C_m^{\ell \begin{smallmatrix} N \\ n \end{smallmatrix} \begin{smallmatrix} J \\ M \end{smallmatrix}} \begin{pmatrix} \ell \\ m \end{pmatrix} \begin{pmatrix} A \\ a \end{pmatrix} \begin{pmatrix} N \\ n \end{pmatrix} \begin{pmatrix} B \\ b \end{pmatrix}^* \begin{pmatrix} J \\ M \end{pmatrix} \begin{pmatrix} C \\ c \end{pmatrix} = \sqrt{\frac{[B]}{[C]}} C_a^A \begin{smallmatrix} B \\ b \end{smallmatrix} \begin{smallmatrix} C \\ c \end{smallmatrix} \langle (\ell)A(R)B|(J)C \rangle \quad (6)$$

Before moving on we will define what we mean by (ABC) . As in reference [12], A labels the internal $\bar{\mathcal{G}}$ symmetry of vibronic activity or excitation" (electronic Σ, Π, Δ , or vibrational δ, π, σ , symmetry). When relating the BOA states to the LWC or scattering states of angular momentum ℓ , the labels considered will be those A correlates with ℓ : $\mathcal{D}^\ell \downarrow \bar{\mathcal{G}} = \dots \oplus A \oplus \dots$. B on the other hand is the internal $\bar{\mathcal{G}}$ symmetry of bare rotor such that there must exist at least one Young Tableau $\{\mu\}$ associated with each a IR(B) of \mathcal{G} . As for the label C , this labels the internal $\bar{\mathcal{G}}$ symmetry of constricted rotor by writing all IR contained in the tensor product $A \otimes B = \dots \oplus C \oplus \dots$. The total angular momentum levels J^p belonging to each (ABC) triad are just those correlated with C : $\mathcal{D}^{J^p} \downarrow \bar{\mathcal{G}} = \dots \oplus C \oplus \dots$.

The preceding (LWC) state is then expressed:

$$\langle [R\ell]_a^{D^* \begin{smallmatrix} B^* \\ b \end{smallmatrix} \begin{smallmatrix} J \\ M \end{smallmatrix}} \rangle = \sum_{ACn_\ell} \langle (\ell)A(R)B|(J)C \rangle \begin{pmatrix} \ell \\ n_\ell \end{pmatrix} \begin{pmatrix} D \\ d \end{pmatrix}^* \sum_{ac} \sqrt{\frac{[B]}{[C]}} C_a^A \begin{smallmatrix} B \\ b \end{smallmatrix} \begin{smallmatrix} C \\ c \end{smallmatrix} r_{n_\ell a}^{\ell A^*}(\Phi\Theta\Xi) r_{M c}^{J C^*}(\alpha\beta\gamma) \quad (7)$$

(Here we consider only the real irreps ($B^* \equiv B$, $D^* \equiv D$) and $[C]$ is the dimension of group irreducible representative (IR) of D^C).

This amounts to a transformation between the (LWC) and the (BOA) state. The BOA basis function is defined as follows:

$$|BOA \ell AC \begin{smallmatrix} B^* \\ b \end{smallmatrix} \rangle = \sum_{ac} \sqrt{\frac{[B]}{[C]}} C_a^A \begin{smallmatrix} B \\ b \end{smallmatrix} \begin{smallmatrix} C \\ c \end{smallmatrix} r_{n_\ell a}^{\ell A^*}(\Phi\Theta\Xi) r_{M c}^{J C^*}(\alpha\beta\gamma) \quad (8)$$

Here the ℓ -rotor A -wave is using the angles $(\Phi\Theta\Xi)$ relative to the other rotor's frame whose coordinates are lab-relative $(\alpha\beta\chi)$, and whose quantum labels are (J, C) with R no longer valid. However the internal B -symmetry of the big rotor is the same as before and related to its nuclear spin state.

Suppose we begin in the LWC where both rotors have a finite symmetry, and move to the BOA basis. Then the transformation Equation (7) takes us from the BOA basis to the LWC basis and an inverse transformation Equation (8) does the reverse. According to Equation (7), the symmetry coefficient $\begin{pmatrix} \ell \\ n_\ell \end{pmatrix} \begin{pmatrix} D \\ d \end{pmatrix}^*$ remains as part of the transformation. The ℓ -rotor locks itself as a passenger on the rotor whose frame is being observed from the lab, while B is preserved through the transformation matrix. In Equation (8), it appears that a new symmetry species A corresponding to the ℓ -passenger is defined relative to the rotor labeled by B , while the symmetry label B remains invariant. However, this may not be the whole story; it seems that there may be two ways of interpreting Equation (8).

To compare our result to that of Harter *et al.* [12], let us assume that the effect of the smaller rotor on the larger rotor symmetry is negligible. So the overriding symmetry then would be that of the large rotor as treated by Harter *et al.* [12], where the ℓ -rotor is treated as a symmetry perturbation. This then leads to one way to compute A and C symmetry labels as described previously by Harter *et al.* [12].

But, what if the rotors are comparable or identical? Then other choices of frames may become relevant, and each may involve another way of interpreting Equation (8). As in reference [12] the symmetry label B may label the “under all” symmetry of “bare” nuclei in the system. Then as we move from a LWC basis to the strongly coupled basis, the B label must correlate the permutation symmetry of all identical nuclei to the composite geometrical point symmetry (labeled C) for the whole of the strongly interacting system. From a mathematical perspective, there is no indication that the symmetry label B in the LWC is different in the BOA basis, but physically, the new geometry suggests that B cannot remain unchanged; it must correlate with the symmetry of the composite BOA rotor states. If the interpretation of B “under all” symmetry species B changes, then both A and C must be determined differently.




Two possible symmetries emerge; one of them specifically corresponds to the larger one of the rotors behaving like a miniature “Lab” for the other, so that the symmetry of the small rotor will correlate with the frame of the larger body. In other words, the BOA transformation has the larger body acting as a lab for the small body that is like a passenger in the frame of the larger body, and respects its symmetry. However, the coefficient $\left(\begin{smallmatrix} \ell \\ n_\ell \end{smallmatrix} \middle| \begin{smallmatrix} D \\ d \end{smallmatrix}\right)^*$ of the second rotor is transferred to a transformation equation such as the equation below:

$$T_d^{\ell RJ(ABC)} = \sum_{ACn_\ell} \langle (\ell)A(R)B|(J)C \rangle \left(\begin{smallmatrix} \ell \\ n_\ell \end{smallmatrix} \middle| \begin{smallmatrix} D \\ d \end{smallmatrix}\right)^* \quad (9)$$

Unlike the orbiting electron of reference [12], this rotor has a finite symmetry. Moreover, identical rotor molecules present the problem of which rotor is “lab” and which is “passenger”. In fact, there can be no way to tell. But once the molecule is locked in some particular geometry, then we can begin symmetry analysis based upon whatever is the composite symmetry.

In our recent work [25,33], we have seen the effects of coupling becoming extremely strong between the molecules so that it causes them to behave as a single rigid body [25]. Where symmetry is concerned, the locking of the two rotors now appears as one body with some new symmetry. Recall that B is the symmetry label for the nuclear points in the rigid R-body but now the B symmetry label may not be the same as it was in the LWC basis. That being the case, B may correlate to a new symmetry belonging to an intersection of the symmetry groups of the two rotor molecules, while the symmetry label A describes the active modes of this composite geometry.

We begin with the finite symmetries of O^D and O^B groups, as they became correlated, the finite symmetry O^D is lost but there is an induced symmetry O^A that is now correlated to the symmetry group O^B . $O^A \times O^B \supseteq O^C$ and O^C is the symmetry group that is correlated to that of O_3 and the total angular momentum J of the coupled rotor described by r_{Mc}^{JC*} . Also, it is possible that O^C is a higher

symmetry as when two diatomic molecules  and  form a T_d molecule .

Thus, the electron-rotor correlation must be generalized further by these possibilities. An electron has no finite internal symmetry only external, therefore, the sum $\sum_{n_\ell} \left(\begin{smallmatrix} \ell \\ n_\ell \end{smallmatrix} \middle| \begin{smallmatrix} D \\ d \end{smallmatrix}\right)^*$ disappears since $n_\ell = 0$. Consequently Equations (7) and (8) reduce to

$$\left| BOA \ell AC \begin{smallmatrix} B^* \\ b \end{smallmatrix} \right\rangle = \sum_{ac} \sqrt{\frac{[B]}{[C]}} C_{a \begin{smallmatrix} A \\ a \end{smallmatrix} b \begin{smallmatrix} C \\ c \end{smallmatrix}}^A Y_{a \begin{smallmatrix} A \\ a \end{smallmatrix}}^{A*}(\Theta \Xi) r_{M \begin{smallmatrix} J \\ c \end{smallmatrix}}^{J C*}(\alpha \beta \gamma) \quad (10a)$$

And

$$\left| [R \ell] \begin{smallmatrix} B^* \\ b \end{smallmatrix} \begin{smallmatrix} J \\ M \end{smallmatrix} \right\rangle = \sum_{AC} \langle (\ell) A(R) B(J) C \rangle \left| BOA \ell AC \begin{smallmatrix} B^* \\ b \end{smallmatrix} \begin{smallmatrix} J \\ M \end{smallmatrix} \right\rangle \quad (10b)$$

Thus, the results in Equations (10a) and (10b) are the same as first given by Harter and his colleagues [12].

2.2. (ABC) Operators and Nuclear Exchange Symmetry

For the BOA basis there are two groups of symmetry operators that are analogous to the LWC symmetry composed of the outer product of O^A for the ℓ -rotor, and O^B for the R-rotor. If we consider the case where the BOA composite symmetry is the inner product $O^A \times O^B = O^C$ of “rigid rotations” then we may apply Equations (10a) and (10b) of electron polyatomic molecule interaction based on the symmetry of the bare R-rotor. We have shown above that transformation preserves the finite symmetry of the bare rotor with labels (B, b) as seen by combining the symmetry properties to the rotor-rotor states.

$$\bar{\mathbf{R}}_{\ell} r_{n_{\ell} a}^{\ell A*}(\mathbf{R}) = \sum_{a'} D_{a' a}^A r_{n_{\ell} a'}^{\ell A*}(\mathbf{R}) \quad (11a)$$

$$\mathbf{R}_J r_{M c}^{J C*}(\mathbf{R}) = \sum_{c'} D_{c' c}^C r_{M c'}^{J C*}(\mathbf{R}) \quad (11b)$$

$\bar{\mathbf{R}} = (\bar{\mathbf{R}}_{\ell}, \mathbf{R}_J)$ is defined with respect to body coordinates as described in [12]. We consider a system in which A is weakly coupled to B , but as we turned on the coupling the symmetry is reduced from $\zeta_A \times \zeta_B$ to ζ_{AB} . The internal frame operator $\bar{\mathbf{R}}$ moves the whole universe except for constricted coupled systems. We must point out that only operators that move the two systems around together will commute with the coupling interaction. We discuss here the symmetry of the general symmetry properties of our general BOA states and compare that to reference [12]. Applying Equations (11a) and (11b) to the (BOA) state, Equation (7) gives the following:

$$\begin{aligned} \bar{\mathbf{R}} \left| BOA (AC) \begin{smallmatrix} B^* \\ b \end{smallmatrix} \begin{smallmatrix} \ell \\ n_{\ell} \end{smallmatrix} \begin{smallmatrix} J \\ M \end{smallmatrix} \right\rangle &= \sum_{ac} \sqrt{\frac{[B]}{[C]}} C_{a \begin{smallmatrix} A \\ a \end{smallmatrix} b \begin{smallmatrix} C \\ c \end{smallmatrix}}^A \bar{\mathbf{R}}_{\ell} r_{n_{\ell} a}^{\ell A*}(\Phi \Theta \Xi) \mathbf{R}_J r_{M c}^{J C*}(\alpha \beta \gamma) \\ &= \sum_{a'c'} \sqrt{\frac{[B]}{[C]}} C_{a' b' c'}^A D_{a' a}^A D_{c' c}^C \left(r_{n_{\ell} a'}^{\ell A*} r_{M c'}^{J C*} \right) \end{aligned} \quad (12)$$

$$\begin{aligned} &= \sum_{b'a'c'} \sqrt{\frac{[B]}{[C]}} C_{a' b' c'}^A D_{b' b}^B r_{n_{\ell} a'}^{\ell A*} r_{M c'}^{J C*} \\ \mathbf{R} \left| BOA (AC) \begin{smallmatrix} B^* \\ b \end{smallmatrix} \begin{smallmatrix} \ell \\ n_{\ell} \end{smallmatrix} \begin{smallmatrix} J \\ M \end{smallmatrix} \right\rangle &= \sum_{b'} D_{b' b}^B \left| BOA (AC) \begin{smallmatrix} B^* \\ b' \end{smallmatrix} \begin{smallmatrix} \ell \\ n_{\ell} \end{smallmatrix} \begin{smallmatrix} J \\ M \end{smallmatrix} \right\rangle \end{aligned} \quad (13)$$

Thus we conclude that the total internal molecular ς symmetry properties are given by the bare rotor symmetry label (B, b) . This proves that the Harter–Patterson and da Paixao relation [12] for an electron coupled with a polyatomic molecule is more general than previously understood. Moreover, the very fact that $\sum_{b'} D_{b', b}^B$ multiplies the BOA state vector, implies that the operator \bar{R} moves both molecules rigidly together. This indicates that the symmetry label B is the representation of a new symmetry or the intersection of $O^D \times O^B$.

2.3. Theory of ABC Symmetry Selection

The symmetry of \bar{G}^C is the overall symmetry happening when both rotor wavefunctions are strongly correlated. We must remember that the total symmetry is based on S_n group products $S_m \times S_n \times \dots$ regardless of BOA or LWC or whatever and comes about from fundamental nuclear identity. S_n is the permutation of all nuclei common to both \bar{G}^B and \bar{G}^D . However, the $S_m \times S_n \times \dots$ challenge come in determining the symmetry label A . This is the symmetry induced to the rotor with angular momentum ℓ by the second rotor to which it is strongly correlated.

If we consider the case where all nuclei are identical, the “under all” nuclear permutational symmetry group is S_n where n is the number of identical nuclei. But, when the rotor molecules are uncorrelated, the relevant permutation symmetry is that of each of the individual rotors, namely S_{n_1} and S_{n_2} . As they become weakly coupled, the permutation symmetry is $S_{n_1} \times S_{n_2}$ and corresponds to the number of identical nuclei with the coupled system. This symmetry is approximately $S_{n_1} \times S_{n_2} \subseteq S_n$ since we assume that bonds between the constituent molecules are unbroken. As a result, we start with the permutation group S_n and the outer product of the individual permutation groups of each molecule. To begin with, we write following for the two rotors:

$$\begin{aligned} S_{n_1} \supset \bar{G}_R, S_{n_2} \supset \bar{G}_\ell \\ (\rho, B), \quad (\lambda, D) \end{aligned} \quad (14)$$

(S_{n_1}) represents the Young tableau irreducible representations whose subduction $D^\rho(S_{n_1}) \downarrow \bar{\xi}_R = \dots \oplus D^B \oplus \dots$, to the group \bar{G}^B gives a B symmetry label. So by the Frobenius reciprocity, $\rho(S_{n_1})$ is in the induction to S_{n_1} by $D^B(\bar{\xi}_R) \uparrow S_{n_1} = \dots \oplus D^\rho \oplus \dots$, where the symbol in the parentheses describe a correlation table for each rotor constituent. Since the two constituent rotors are interacting then their geometry of the symmetry is described by the product of the external and internal symmetry of each rotor that is:

$$\begin{pmatrix} \zeta_R^{LAB}, \bar{G}_R \\ (R, B) \end{pmatrix} \times \begin{pmatrix} \zeta_\ell^{LAB}, \bar{G}_\ell \\ (\ell, D) \end{pmatrix} \quad (15)$$

However, at first, we will only concern ourselves with the internal point symmetry. Thus, the symmetry is given by $\bar{G}^D \times \bar{G}^B$. This new group is a subgroup of the permutation S_n where n is the number of identical nuclei common to both molecules.

$$\begin{aligned} \bar{G}^D \times \bar{G}^B = \bar{G}^B \equiv S_{n_1} \times S_{n_2} \subseteq S_n \\ \text{where } n_1 + n_2 = n \end{aligned} \quad (16)$$

The first step is to compute the character table of S_n . Next we must perform symmetry analysis in both of our bases, namely LWC and BOA basis. As mentioned before for simplicity, we assume the simpler case where all nuclei are identical.

We will start with uncoupled bases and symmetry analysis of the individual constituent rotors. The symmetry groups are S_{n_1} and S_{n_2} which are subgroups of S_n . In the weakly coupled basis it is assumed that the symmetry group is S_n for a group of identical nuclei. For example two H_2 molecules would have a symmetric group of S_4 . We also need a correlation table between S_n and O_3 for the rotation group of the entire rotor-rotor system.

At first, nature reduces the symmetry from S_n to a lower symmetry since initial chemical bonds between individual rotors prevent all $n!$ permutations from being allowed, that is, $S_n \supseteq S_{n_1} \times S_{n_2}$. Initially, we assume that the molecules will have some freedom to rotate individually, and subjected to the coupling between them, give rise to various geometrical structures that may become a final geometry if a very strong interaction takes effect. For example, two H_2 molecules might later find themselves locked into D_2 , D_{2d} or D_4 symmetry. While the system is weakly coupled, they move through all these geometries giving rise to various states that make up a generalized cluster basis, that is, an induced representation basis. In the LWC basis, we have permutation symmetry $S_{n_1} \times S_{n_2}$ giving to two classes of operators, those that leave the two H_2 molecules together that is bi-cyclic operators like (12)(34), and those that move the individual H_2 molecules that is, monocyclic operators like (12) which is the same as transposition in group theory.

Therefore, we compute the correlation $S_{n_1} \times S_{n_2} \uparrow S_n$: This seems like a difficult task, in general, but by using Young tableau the procedure becomes easier. To achieve this, we must find the outer product using the Frobenius Reciprocity Theorem. If $S_{n_1} \times S_{n_2}$ is isomorphic to a composite molecular point group, then the approach in determining the C label is very easily done.

The BOA basis is approached if the coupling becomes strong enough to lock our coupled system. At this point, we assume that the coupling is so strong that molecules can only vibrate in the planes to which they are confined so the angle between them is almost constant. Then we can describe the composite system by some geometrical point group of a rigid (or semi-rigid) molecule.

We now outline a procedure to find the symmetry labels A and C . We find the “under all” permutational symmetry in the BOA constricted situation. Observing the intersection of $\bar{G}^D \times \bar{G}^B$ which is some particular geometry point group gives us the B labels. We defined B to be the symmetry label of the identical nuclei within the new arrangement. The derivation of the symmetry species C comes from the irreducible modes that are active. When we move to the BOA basis, certain modes become active. These active modes have an A (“Activity”) symmetry label. This lets us find C . Once the particular point group is determined and the modes that can be excited are found, then C is just the irreducible representation of the outer product of A and B , that is $A \otimes B = C$.

This gives all the possible symmetry pieces of C . We will consider an example of diatomic–diatomic interaction. Since the overall external symmetry is O_3 , knowing C gives a correlation induced between the irrep C and that of O_3 . Similarly, the correlation between the irreducible representation of O_3 and $\bar{G}^D \times \bar{G}^B$ or $S_{n_1} \times S_{n_2}$ can be computed. The characters of R_3 is given by

$$race D^{\ell}(\omega 00) = \frac{\sin\left(\ell + \frac{1}{2}\right)\omega}{\sin \omega/2} \quad (17)$$

We derive the character of O_3 from its outer product relation to R_3 .

$$R_3 \times C_i = O_3 \quad (18a)$$

Trace D(w 0 0)										
O_3	1 $\omega = 0^\circ$	r, r^2 $\omega = 120^\circ$	R^2 $\omega = 180^\circ$	R, R^3 $\omega = 90^\circ$	i $\omega = 180^\circ$	I $\omega = -180^\circ$	Ir, Ir^2 $\omega = 120^\circ$	IR^2 $\omega = 180^\circ$	IR, IR^3 $\omega = 90^\circ$	Ii $\omega = 180^\circ$
0^+	1	1	1	1	1	1	1	1	1	1
0^-	1	1	1	1	1	-1	-1	-1	-1	-1
1^+	3	0	-1	1	-1	3	0	-1	1	-1
1^-	3	0	-1	1	-1	-3	0	1	-1	1
2^+	5	-1	1	-1	1	5	-1	1	-1	1
2^-	5	-1	1	-1	1	-5	1	-1	1	-1
3^+	7	1	-1	-1	-1	7	1	-1	-1	-1
3^-	7	1	-1	-1	-1	-7	-1	1	1	1
4^+	9	0	1	1	1	9	0	1	1	1
4^-	9	0	1	1	1	-9	0	-1	-1	-1
5^+	11	-1	-1	1	-1	11	-1	-1	1	-1
5^-	11	-1	-1	1	-1	-11	1	-1	1	1
6^+	13	1	1	-1	1	13	1	1	-1	1

(18b)

The frequency $f^C(J)$ of the irreducible representation of C subduced to O_3 is given by

$$f^C = \frac{1}{G^\circ} \sum_{\substack{\text{classes} \\ c_g \in C}} \chi_g^{C^*} c_g \frac{\sin\left(\ell + \frac{1}{2}\right)\omega}{\sin \omega/2} \quad (19)$$

where C is the irrep label of the point group of the composite rigid body.

3. Spin Statistical Weights

One cannot speak of statistical weight without mentioning the extensive review by Bunker and Jensen [23], which outlines a procedure to calculate statistical weight for molecular symmetry. Hougen [34] has used a classical approach to classify the various labeling schemes of molecular symmetry. Alvarez-Bajo *et al.* [35] revisited group computed spin weight for methane using symmetry projection. As pointed out earlier, large amplitude motions may result in configurations that can be described by molecular symmetry group as shown by Bunker *et al.* [23]. However, we consider the treatment of such with symmetry permutation groups, which sometimes reveal details not obvious from traditional approaches. The power of such approach can be seen in a comparison study done by Harter *et al.* [12] with old techniques used by Hougen [34] to showcase Young tableau as a powerful tool that can be used to compute statistical weight. In their work, Harter points out that the tableau method gives the same result as long as inversion does not play a dynamical role. In comparing this method with that of Hougen [34], Harter shows that it reveals the hidden structure of the inversion doublets for various molecular symmetries.

The symmetric classification for nuclear wavefunction of a diatomic molecule is done by considering its two nuclei. However, symmetry classification may appear ambiguous because it depends on the details of the permutation of nuclei. Harter and colleagues [12] have shown how to assign Young tableau to the point symmetry species B . Furthermore, n identical nuclei may have one or more orbital tableau corresponding to each B labels of the spatial point group, and for each tableau, there is just one nuclear spin tableau that describes the permutational properties of the nuclear spin wavefunction. This is due to the Bose–Einstein Principle demanding the spin tableau of the nuclear state to be identical to its orbital tableau while the spin tableau of the Fermi–Dirac nuclear state is conjugated to its orbital tableau. The Bose nuclear states are those with integral spins such as a photon, or deuterium nuclei, whereas the Fermi nuclear states are those with half-integer spins such as nuclei with odd numbers of nucleons.

To compute the spin weight one must do a little U_m algebra. The statistical weight found by using the Robinson formula along with the correlation frequencies $f_B^{\mu_s}$ give

$$W_B = \sum_{\mu_s} l^{\mu_s} f_B^{\mu_s} \quad (20a)$$

where l^{μ_s} is given in Figure 28 of reference [12] as

$$l^{\{\mu_s\}} = \frac{\text{Dimension of } \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline \end{array}}{\text{product of } \begin{array}{|c|c|c|c|c|} \hline m & m+1 & m+2 & m+3 & m+4 & \cdots \\ \hline m-1 & m & m+1 & \cdot & \cdot & \\ \hline m-2 & m-1 & \cdot & \cdot & \cdot & \\ \hline m-3 & \cdot & \cdot & \cdot & \cdot & \\ \hline \end{array}} \quad (20b)$$

$m=2I+1$

product of hooklengths

The hooklength of a box in the tableau is the number of boxes in a “hook” which includes that box and all boxes in the line to the right and in the column below it. This formula gives directly the number l^{μ_s} of nuclear spin states associated with each tableau $\{\mu_s\}$. The sum of these numbers taken over each tableau associated with “soul” label B will be the desired statistical weight.

These correlation frequencies are quite simple for point groups but not as easy for the permutation groups. We will give a few formulas for the permutation group. If for the individual molecules we have two nucleons ($spin\ I = 1/2$), the necessary U_m is U_2 , three would be U_4 since $I = 3/2$, that is:

$$m = 2I + 1 \quad (21)$$

where I is the spin angular momentum. For Bosons: $I = 1$ use U_3 , $I = 2$ nuclei use U_5 , and so on. For a more detailed approach and understanding of how to use tableau to compute statistical spin weight for molecular system, one should consult the work of Harter *et al.* [12].


4. Coupling between Two Diatomic Molecules

It is well known that diatomic molecules such as H_2 , O_2 , and D_2 have $D_{\infty h}$ symmetry. However, the nuclear spins of these molecules are different. H_2 has two protons with each spins of $I = 1/2$, O_2 nuclear spin is $I = 0$, and D_2 nuclei each have each nuclear spins of $I = 1$. Thus H_2 is fermionic, whereas O_2 and D_2 are bosonic. The nuclear spins of all elements and their isotopes are very well known and can be found in tables of nuclides [36].

The spin statistical weight determines the relative intensity of spectra belonging to each B label. H_2 contain two spin-1/2 protons where the state $B = \Sigma_g^+$ by its even parity correlates to the antisymmetric tableau $\{\mu_s\} = \{1^2\}$ of singlet ($I = 0$) spin tableau by Fermi-Dirac rules. The state $B = \Sigma_\mu^+$ belongs to the symmetric spin tableau $\{\mu_s\} = \{2\}$ that is, triplet ($I = 1$) total spin. One should note that for a diatomic molecule with symmetry $D_{\infty h}$ the only choices are $B = \Sigma_g^+$ or Σ_μ^+ . Deuterated hydrogen D_2 contains two spin-1 deuterons where, by Bose-Einstein rules, the state $B = \Sigma_\mu^+$ by its parity correlates to the $\{\mu_s\} = \{1^2\}$ antisymmetric tableau and Σ_g^+ belongs to the tableau $\{\mu_s\} = \{2\}$. The state Σ_g^+ has spin states of total nuclear spin angular momentum of $S = 2$ and $S = 1$, while Σ_μ^+ is the deuterium triplet with $S = 1$. However, O_2 only has ortho-states species $\Sigma_g^+ = \{2\}$. The para species Σ_μ^+ are Pauli excluded because spin-0 states cannot be antisymmetric.

We now investigate symmetry structure and levels of two diatomic molecules. We have assumed that the interaction is between identical diatomic molecules. The energy levels for two uncoupled H_2 molecules are shown in Figure 1. As these molecules come into the vicinity of each other, we suppose they begin to interact in a weakly correlated state [25].

Figure 1. Rotational Energy levels of a H_2 molecule.



In the LWC basis, the symmetry of the molecular system is nearly the S_4 permutation group, which is isomorphic to that of the tetrahedral point T_d . The character table of S_4 is given in Equation (22) where $\{\mu\}$ represents the irreps of S_4 .

S_4	(1)	(132)	(14)(23)	(1234)	(13)
$\{4\}$	1	1	1	1	1
$\{1^4\}$	1	1	1	-1	-1
$\{2,2\}$	2	-1	2	0	0
$\{2,1^2\}$	3	0	1	1	-1
$\{3,1\}$	3	0	-1	-1	1

(22)

Since the overall symmetry of the R_3 group is O_3 then the correlation between S_4 and O_3 is given as follows. This uses the isomorphism of T_d and S_4 .

$T_d \uparrow O_3$	A_1	A_2	E	T_1	T_2
$S_4 \uparrow O_3$	$\{4\}$	$\{1^4\}$	$\{2^2\}$	$\{2,1^2\}$	$\{3,1\}$
$J^P = 0^+$	1
0^-	.	1	.	.	.
1^+	.	.	.	1	.
1^-	1
2^+	.	.	1	.	1
2^-	.	.	1	1	.
3^+	.	1	.	1	1
3^-	1	.	.	1	1
4^+	1	.	1	2	1
4^-	.	1	1	1	2
5^+	.	.	1	2	1
5^-	.	.	1	1	2
6^+	1	1	1	1	2
6^-	1	1	1	2	1
7^+	1	.	1	2	2
7^-	.	1	1	2	2

(23)

Equation (23) tells us how the angular momentum energy states splits according to S_4 symmetry group. As the angular momentum increases the effects of clustering begin to be manifested [11,12]. This is more noticeable in the irreps $\{2,1^2\}$ and $\{3,1\}$ of S_4 . But as we go from the LWC basis to the BOA basis by increasing the coupling strength, the nuclei is locked into a particular geometry. If the energy between the molecules is not high enough to cause bond breaking in the individual diatomic molecules we may see clustering of the energy levels.

4.1. $S_2 \times S_2$ Point Group

X_2 molecule has two identical nuclei, which means it has a permutation symmetry group of S_2 . When the molecules are weakly coupled, constituents allow full S_4 permutations, but as the coupling increases the permutation symmetry may be reduced to $S_2 \times S_2$, the outer product between the symmetry groups of the two diatomic molecules. The characters of $S_2 \times S_2$ are listed below:

S_2	<div style="display: flex; justify-content: space-around;"> (1) (12) </div> <div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div> <div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div>	\otimes	S_2	<div style="display: flex; justify-content: space-around;"> (1') (12') </div> <div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div> <div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div>
	<div style="display: flex; justify-content: space-around;"> 1 1 </div> <div style="display: flex; justify-content: space-around;"> 1 -1 </div>		<div style="display: flex; justify-content: space-around;"> 1 1 </div> <div style="display: flex; justify-content: space-around;"> 1 -1 </div>	

$S_2 \times S_2$	1	Ii_3	Ii_4	R_y
	1	(34)	(12)	(12)(34)
	(1)(1')	(1)(12')	(12)(1')	(12)(12')
<div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div>	1	1	1	1
<div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div>	1	-1	1	-1
<div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div>	1	1	-1	-1
<div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 20px; height: 20px; margin: 2px;"></div> </div>	1	-1	-1	1

(24)

The $S_2 \times S_2$ group is isomorphic to C_2 . $S_2 \times S_2$, however, it allows a 180° rotation about the center of each individual X_2 molecules, while the other stays fixed; for instance consider the configuration



where the numbers label the position of the atoms of the X_2 molecules. Under the $S_2 \times S_2$ operation (34), the previous configuration changes to



By using Frobenius Reciprocity theorem we obtain the $S_2 \times S_2 \uparrow S_4$ reduction table. In left column of Equation (25) we give the spin tableaus for fermions and Boson nuclei:

$S_4 \supset S_2 \times S_2$		$\{2\} \cdot \{2\}$	$\{1^2\} \cdot \{1^2\}$	$\{2\} \cdot \{1^2\}$	$\{1^2\} \cdot \{2\}$
<i>Fermi spin</i>	<i>Boson spin</i>				
$\{1^4\}$	$\{4\}$	1	.	.	.
$\{4\}$	$\{1^4\}$.	1	.	.
$\{2,2\}$	$\{2,2\}$	1	1	.	.
$\{3,1\}$	$\{2,1^2\}$.	1	1	1
$\{2,1^2\}$	$\{3,1\}$	1	.	1	1

(25)

In BOA basis the interaction is so strong that their position is locked in together preventing the large amplitude motion that is allowed in LWC basis. The only symmetry operators that are allowed are those that move both molecules rigidly together. If the molecules are locked in parallel to each other then they belong to a group with D_2 or D_{2h} symmetry but if oriented at right angles then the group form is one with D_4 or D_{2d} symmetry. We investigate X_2 molecules orientation with either D_{2h} or D_4 symmetry.

4.2. Statistical Weight $S_2 \times S_2$

The algebra needed is U_2 for Fermions of spin $I = 1/2$. From Equation (21) we have $m = 2$. This means that there are three possible choices for the numerator of l^{μ_s} by Equation (20b):

$\{4\} = \begin{array}{|c|c|c|c|} \hline 2 & 3 & 4 & 5 \\ \hline \end{array}$, $\{2,2\} = \begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline \end{array}$, and $\{3,1\} = \begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & & \\ \hline \end{array}$. The Frobenius Reciprocity theorem

gives the correlation frequencies and using Equations (20a) and (20b) we compute the statistical weight for $S_2 \times S_2$ to be,

$$W_{\{2\},\{2\}} = \frac{\begin{array}{|c|c|c|c|} \hline 2 & 3 & 4 & 5 \\ \hline 4 & 3 & 2 & 1 \\ \hline \end{array}}{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline \end{array}} + \frac{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline \end{array}}{\begin{array}{|c|c|} \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}} = 6 \quad (26a)$$

$$W_{\{2\},\{1^2\}} = W_{\{1^2\},\{2\}} = \frac{\begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & & \\ \hline \end{array}}{\begin{array}{|c|c|c|} \hline 4 & 2 & 1 \\ \hline 1 & & \\ \hline \end{array}} = 3 \quad (26b)$$

$$W_{\{1^2\},\{1^2\}} = \frac{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline \end{array}}{\begin{array}{|c|c|} \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}} + \frac{\begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & & \\ \hline \end{array}}{\begin{array}{|c|c|c|} \hline 4 & 2 & 1 \\ \hline 1 & & \\ \hline \end{array}} = 4 \quad (26c)$$

Equations (26a–c) shows how the various symmetry levels would cluster. The state corresponding to the symmetry species $\{2\}\{2\}$ would consist of four degenerate levels whereas the states with symmetry $\{2\}\{1^2\}$ would be triply degenerated. The use of Young tableau gives information about the clustering of the level. As we continue to reduce the symmetry we will compute the statistical weight using Young tableau.

4.3. D_{2h} Point Group

When two X_2 molecules are aligned parallel to each other, their arrangement gives a symmetry group of $D_2: \{1, R_x, R_y, R_z\}$ that is, there is a R 's rotation of 180° degrees around the x , y , and z axes. X_2 configuration contains a horizontal reflection plane $\sigma_h = \sigma_{xy} = IR_z$ where I is the inversion operator such that $\begin{array}{c} \textcircled{1} \text{---} \textcircled{2} \\ \textcircled{3} \text{---} \textcircled{4} \end{array} \longrightarrow \begin{array}{c} \textcircled{3} \text{---} \textcircled{4} \\ \textcircled{2} \text{---} \textcircled{1} \end{array}$. This operator works by inverting the spatial coordinates of all nuclei through the center of mass. The overall symmetry of X_2 configuration is the outer product of D_2 and $i = \{1, I\}$ that is $D_{2h} \equiv D_2 \otimes i$. D_{2h} contains three 180° degrees rotations about the xyz axes, an I inversion, a horizontal reflection σ_{xy} , and two vertical planes ($\sigma_{xz} = IR_x$), $\sigma_{yz} = IR_y$. The character table for D_{2h} symmetry group is given by Equation (27).


D_{2h}	1	R_x	R_y	R_z	I	IR_x	IR_y	IR_z
	(1)	(13)(24)	(12)(34)	(14)(23)	(14)(23)	(12)(34)	(13)(24)	(1)
A_{1g}	1	1	1	1	1	1	1	1
B_{1g}	1	1	-1	-1	1	1	-1	-1
A_{2g}	1	-1	-1	1	1	-1	-1	1
B_{2g}	1	-1	1	-1	1	-1	1	-1
A_{1u}	1	1	1	1	-1	-1	-1	-1
B_{1u}	1	1	-1	-1	-1	-1	1	1
A_{2u}	1	-1	-1	1	-1	1	1	-1
B_{2u}	1	-1	1	-1	-1	1	-1	1

(27)

Let us consider a spring model system such as shown in the Figure 2 in order to compute the vibrational levels in the plane in which both diatomic molecules lie. Since we assume that the D_{2h} symmetry, then the coordinates are described by following state vectors

$$x_i = \langle i|x \rangle \quad i = 1, 2 \dots 8 \quad \text{or} \quad |i \rangle \quad i = 1, \dots, 8 \quad (28)$$

Figure 2. Spring mass system for four nuclei.



$$\langle i|\ddot{x}\rangle = \sum_{j=1}^8 \langle i|a|j\rangle \langle j|x\rangle \quad (29a)$$

Where

$$\langle i|a|j\rangle = \begin{bmatrix} H & S & . & M & L & . & . & L \\ S & H & M & . & . & L & L & . \\ . & M & H & S & . & L & L & . \\ M & . & S & H & L & . & . & L \\ L & . & . & L & H' & . & S' & M' \\ . & L & L & . & . & H' & M' & S' \\ . & L & L & . & S' & M' & H' & . \\ L & . & . & L & M' & S' & . & H' \end{bmatrix} \quad (29b)$$

And $H = j\cos^2\theta + k$, $H' = j\sin^2\theta + m$, $S = -k$, $S' = -m$, $L = j\cos\theta\sin\theta$, $M = j\cos^2\theta$, and $M' = j\sin^2\theta$.

From Equation (27) we build our D_{2h} projectors; For example the projector $P^{A_{1g}} = 1/8[(1)1 + (1)R_z + (1)R_y + (1)R_x + (1)E^* + (1)\sigma_{xy} + (1)\sigma_{yz} + (1)\sigma_{xz}]$. When this acts state $|1\rangle$ we have $P^{A_{1g}}|1\rangle = 1/4[|1\rangle + |2\rangle + |3\rangle + |4\rangle]$. The application of $P^{A_{1g}}$ on $|1\rangle$ gives non-zero for states parallel it and give a similar result when applied to $|5\rangle$. Only irreps $\{A_{1g}, A_{2g}, B_{1u}, B_{2u}\}$ give a nonzero result.

Therefore, the application of D_{2h} projectors to the coordinates, gives the 8×8 transformation matrix Equation (30).

$$T = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 & . & . & . & . \\ 1 & -1 & 1 & -1 & . & . & . & . \\ 1 & -1 & -1 & 1 & . & . & . & . \\ 1 & 1 & -1 & -1 & . & . & . & . \\ . & . & . & . & 1 & 1 & 1 & 1 \\ . & . & . & . & 1 & 1 & -1 & -1 \\ . & . & . & . & 1 & -1 & -1 & 1 \\ . & . & . & . & 1 & -1 & 1 & -1 \end{pmatrix} \quad (30)$$

The transformation contains two 4×4 blocks, which block diagonalized the force matrix due to the presence of D_{2h} symmetry. As a result, the force matrix is reduced by this transformation to two 2×2 matrices and four 1×1 matrices or eigenvalues.

$$\begin{pmatrix} |B_{1u}^5\rangle & |A_{1g}^1\rangle \\ \hline 2j \cos^2 \theta + p & 2j \cos \theta \sin \theta \\ 2j \cos \theta \sin \theta & 2j \sin^2 \theta + p \end{pmatrix} \quad (31a)$$

$$\begin{pmatrix} |A_{1g}^1\rangle & |B_{1u}^5\rangle \\ \hline 2k + 2j \cos^2 \theta + p & 2j \cos \theta \sin \theta \\ 2j \cos \theta \sin \theta & 2m + 2j \sin^2 \theta + p \end{pmatrix} \quad (31b)$$

The parameter p is the spring constant if the spring system is connected to the springs along all eight coordinate axes, but we will assume that is not the situation there because $p = 0$. Therefore, the eigenvalues for Equation (31a) are 0 and $2j$, and eigenvectors $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$. But for Equation (31b) we have the following eigenvalue:

$$\eta_{\pm} = \alpha \pm \beta \quad (32)$$

where $\alpha = j + m + k$ and $\beta = \sqrt{(j - m + k)^2 + j(m - k)\sin^2 \theta}$.

The eigenvectors are listed below their eigenvalues

$$\frac{\begin{pmatrix} \frac{2}{\sqrt{3}} \eta_{\pm} \\ 1 \end{pmatrix}}{N} \quad (33)$$

Where $N = \sqrt{\frac{2\eta_{\pm}^2 + 3}{3}}$.

The other eigenvalues are $(2k$ and $0)$, $(0$ and $2m)$, which corresponds eigenvectors A_{2g} and B_{2u} modes of the D_{2h} respectively.

4.4. Statistical Weight for D_2

Bunker and Jensen [23] have given a detailed development of permutation and permutation inversion groups, which one should consult for a thorough understanding. The inversion group $i = \{1, I\}$ is synonymous to their $\varepsilon = \{E, E^*\}$. $D_{2h} \not\subseteq S_4 \equiv T_d$ since S_4 is absent of the inversion operation E^* . We use the out product between the groups S_4 and ε such that $D_{2h} \subseteq S_4 \otimes \varepsilon \equiv O_h$. The E^* operator commutes with all S_4 operations; thus the $S_4 \otimes \varepsilon$ group is called a complete nuclear permutation inversion (CNPI [23]) group. We are now in position to compute the correlation between D_{2h} and $S_4 \otimes \varepsilon$. For brevity we consider a much simpler correlation between the molecular symmetry group D_2 and S_4 since only the irreps $\{A_{1g}, A_{2g}, B_{1u}, B_{2u}\}$ give a non-result. The resulting correlation table is shown in Figure 3.

From Equations (20a) and (20b) we compute the spin weight for the D_{2h} point group by using Equation (27), along with the correlation table shown in Figure 3. As a result, the following spin state weights are found.


$$W_{A_{2g}} = W_{B_{1u}} = W_{B_{2u}} = \frac{\begin{array}{|c|} \hline 1 \\ \hline \end{array}}{\begin{array}{|c|c|c|} \hline 4 & 2 & 1 \\ \hline \end{array}} = 3 \quad (34a)$$

$$W_{A_{1g}} = \frac{\begin{array}{|c|c|c|c|} \hline 2 & 3 & 4 & 5 \\ \hline 4 & 3 & 2 & 1 \\ \hline \end{array}}{\begin{array}{|c|c|c|c|} \hline 4 & 3 & 2 & 1 \\ \hline \end{array}} + 2 \frac{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}}{\begin{array}{|c|c|} \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}} = 7 \quad (34b)$$

Figure 3. Energy spectrum and correlation diagram for D_2 . Structure of two diatomic molecules.

Duality: The “Flip Side” of Symmetry Analysis. (Continued)

LAB versus BODY, STATE versus PARTICLE,



The irreps $\{A_{1g}, A_{2g}, B_{1u}, B_{2u}\}$ of D_{2h} are the only active modes and are the B labels of the internal symmetry of the bare rotor. The modes of the constricted rotor come from the tensor product $A \otimes B = C$, as shown in the energy level diagram in Figure 3. As we move down the column of the correlation table we see clustering formation; for example the irreps $\{2, 1^2\}$ and $\{3, 1\}$ of S_4 comes together to form the B_{2u} cluster of D_{2h} as displayed in the right of Figure 3. On the other hand, moving across the row shows the splitting of the irreps of S_4 into various irreps of D_{2h} .

Figure 3 is an illustration of Lab-body duality. The rows represent “coerced” or forced symmetry breaking, this comes from applying some kind of external field, which would result in the energy level splitting. The columns describe what is happening in the body frame; in other words, the symmetry reduction brings about clustering. The energy level diagrams of Figure 3 were obtained from Equation (27) through Equations (34a) and (34b).

4.5. D_2 Point Group

Diatomic rotor-rotor BOA structures of D_4 or D_{2d} point group symmetry may come about when the two diatomic molecules are paired at right angles to each other. Equation (35) gives the character table for the D_4 point group. It is the same character table as its isomorphic symmetry D_{2d} for which the operators R_3, R_3^3, i_3 , and i_4 are paired with inversion E^* .

D_{2d}	1	R_3^2	$E^* R_3, E^* R_3^3$	R_1^2, R_2^2	$E^* i_3, E^* i_4$
A_1	1	1	1	1	1
B_1	1	1	−1	1	−1
A_2	1	1	1	−1	−1
B_2	1	1	−1	−1	1
E	2	−2	0	0	0

(35)

For a rotor-rotor locked in D_4 point symmetry, each of its S_4 tableau symmetry labeled levels must split according to a D_4 point symmetry Equation (36).

$S_4 \supset D_4$		A_1	A_2	B_1	B_2	E
<i>Fermi spin</i>	<i>Bose spin</i>					
$\{1^4\} \downarrow D_4$	$\{4\} \downarrow D_4$	1
$\{4\} \downarrow D_4$	$\{1^4\} \downarrow D_4$.	.	1	.	.
$\{2, 2\} \downarrow D_4$	$\{2, 2\} \downarrow D_4$	1	.	1	.	.
$\{3, 1\} \downarrow D_4$	$\{2, 1^2\} \downarrow D_4$.	1	.	.	1
$\{2, 1^2\} \downarrow D_4$	$\{3, 1\} \downarrow D_4$.	.	.	1	1

(36)

Now, reading the columns gives the induced representations such as $A_2 \uparrow S_4 = \{2, 1^2\}$ for a Boson. From this, we construct spin–tableau correlations with D_4 symmetry states, for either Bose or Fermi nuclei.

4.6. Statistical Weight for D_4

As with the D_2 analysis we calculate the spin weight for the irreps of D_4 . Since Fermions spin $I = 1/2$, the unitary algebra is U_2 . The B_2 spin weight is zero for D_4 symmetry since in the group correlation table between D_4 and S_4 , the B_2 irrep correspond only to $\{2, 1^2\}$, and using Robinson's hook length formula as defined in reference [11], we compute the numerator to be zero. The total sum of the spin weight is 13 and not 16. However, we need not be alarmed since the spin weight for doubly degenerated E must be double counted, hence, we still have 16 spin states in all.

$$W_E = W_{A_2} = \frac{\begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & & \\ \hline 4 & 2 & 1 \\ \hline 1 & & \\ \hline \end{array}}{\begin{array}{|c|c|c|} \hline 2 & 3 & 4 \\ \hline 1 & & \\ \hline 4 & 2 & 1 \\ \hline 1 & & \\ \hline \end{array}} = 3 \quad (37a)$$

$$W_{A_1} = \frac{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}}{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}} = 1 \quad (37b)$$

$$W_{B_1} = \frac{\begin{array}{|c|c|c|c|} \hline 2 & 3 & 4 & 5 \\ \hline 4 & 3 & 2 & 1 \\ \hline \end{array}}{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}} + \frac{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}}{\begin{array}{|c|c|} \hline 2 & 3 \\ \hline 1 & 2 \\ \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}} = 6 \quad (37c)$$

The energy level clusters shown in Figure 4 for a D_4 structure are quite different from that of D_2 . E-symmetry species indicate double degeneracy but Young Tableau assigns each of the two levels with a triple degeneracy. This is shown in the energy diagram of Figure 4.

Thus using (2) we can find BOA constricted wave function. Consider the angular momentum $\ell = 1^-$ state. From the correlation table we have for A labels $\{B_2, E\}$ and the bare rotor is D_4/D_{2d} symmetry, thus B labels are irreps of D_4/D_{2d} point group. We now computed the labels of constricted rotor from the tensor product $A \otimes B = C$ that is $\{1^-, \{A = \{B_2, E\} \otimes B = \{A_1, B_1, A_2, B_2, E\}\} = 1^-, B_2, EB_2, E, E, E, E \otimes E = A_1 \oplus B_1 \oplus A_2 \oplus B_2 \oplus E$. This shows in the energy level diagram in Figure 4.

5. Tunneling Hamiltonians

We can write down various tunneling Hamiltonian matrix for each of the symmetry above as we move from S_n symmetry to any of the others. We will start by giving the regular representation of the four particle states for each symmetry case $\{|1\rangle, |2\rangle, |3\rangle, |4\rangle\}$. From S_n we may have all the possible final point group geometry as given in Figure 5. However, $S_2 \times S_2$ and D_2 have only four possible arrangements whereas D_4 has eight. D_2 is a subset of D_4 , but $S_2 \times S_2$ although isomorphic to D_2 , is not a subset of D_4 . Figure 6 shows all the arrangements of $S_2 \times S_2$, D_2 and D_4 .

The reason why we have not done all 24 arrangements is due to the existence of the bonds between the two diatomic molecules, as a result, only certain feasible operations are allowed. To take advantage of the full S_4 symmetry, it would be necessary to break the bonds between the diatomic molecules. We

are ready to investigate each tunneling paths from S_4 to $S_2 \times S_2$, D_2 and D_4 . Here $|1\rangle$ is the argument (1234), and $|2\rangle$ is (2143), $|3\rangle$ is (2134), and $|4\rangle$ is (1243) as shown in Figure 6. Each one is obtained from $S_2 \times S_2$ operations. Similarly, we can derive the states for D_4 , D_{2d} , or D_{2h} by using their operations on state $|1\rangle$.

Figure 4. Energy spectrum and correlation diagram for D_4 structure of two diatomic molecules.






Figure 5. All possible paths from S_4 to a lower symmetry.**Figure 6.** All possible arrangement from S_4 to $S_2 \times S_2$, D_2 and D_4 . The state is shown in the kets, and the possible S_4 operator is represented in parentheses. The upper case letters represent the tunneling parameter between different configurations.

5.1. Tunneling Hamiltonian for $S_2 \times S_2$ Point Symmetry

In the LWC basis the molecular system have S_4 symmetry. As we increase the coupling or interaction, the symmetry is reduced. $S_2 \times S_2$ is one such symmetry reduction sub-group where the individual molecules still have some freedom to rotate in and out of the plane in which they are confined. In more recent work, Hougen *et al.* [37] have extended permutation-inversion group studying large amplitude motion acetylene. The earliest work in treating large amplitude motion in non-rigid molecules goes back to Longuet-Higgins [15]. $S_2 \times S_2$ has two configurations that normal symmetry operations cannot reproduce. Applying the permutation operations $\{1, (12), (34), (12)(34)\}$ to each of the base states $\{|1\rangle, |2\rangle, |3\rangle, |4\rangle\}$ gives the following regular representation of $S_2 \times S_2$.

$$\begin{pmatrix} 1 & . & . & . \\ . & 1 & . & . \\ . & . & 1 & . \\ . & . & . & 1 \end{pmatrix}, \begin{pmatrix} . & 1 & . & . \\ 1 & . & . & . \\ . & . & . & 1 \\ . & . & 1 & . \end{pmatrix}, \begin{pmatrix} . & 1 & . & . \\ 1 & . & . & . \\ . & . & 1 & . \\ . & . & . & 1 \end{pmatrix}, \begin{pmatrix} 1 & . & . & . \\ . & 1 & . & . \\ . & . & . & 1 \\ . & . & 1 & . \end{pmatrix} \quad (38)$$

$R(1) \quad R((12)(34)) \quad R(12) \quad R(34)$

These matrixes are for a group table that is isomorphic to D_2 .

$S_2 \times S_2$	1	(12)	(34)	(12)(34)
1	1	(12)	(34)	(12)(34)
(12)	(12)	1	(12)(34)	(34)
(34)	(34)	(12)(34)	1	(12)
(12)(34)	(12)(34)	(34)	(12)	1

(39)

Consider the symmetry operators g that commutes with the Hamiltonian H , *i.e.*, $gH = Hg$, but the non-commutative symmetry D_3 , D_4 , *etc.*, we build our Hamiltonian matrix by combining the dual symmetry operators as described by Harter [38]. Equation (39) shows that $S_2 \times S_2$ forms an Abelian group, thus we can construct the Hamiltonian matrix since it naturally commutes with the $S_2 \times S_2$ operators. The resulting parameters H , \bar{R} , M , and \bar{L} can be visualized as coupling or tunneling parameters related to the symmetry path. The bar is used as an indicator that only one of X_2 is being operated on; thus the outcome does not match a point group operation. \bar{L} is the tunneling parameter between $|1234\rangle$ and $|1243\rangle$, and \bar{R} is the tunneling parameter between $|1234\rangle$ and $|2134\rangle$, and M is between $|1234\rangle$, and $|2143\rangle$. Writing the tunneling Hamiltonian in the basis of the regular representation Equation (38): $H = H1 + M(12)(34) + \bar{L}(12) + \bar{R}(34)$ which gives:

$$R(H) = \begin{pmatrix} \langle 1 | \\ \langle 2 | \\ \langle 3 | \\ \langle 4 | \end{pmatrix} \begin{pmatrix} H + \bar{R} & M + \bar{L} & . & . \\ M + \bar{L} & H + \bar{R} & . & . \\ . & . & H + \bar{L} & \bar{R} + M \\ . & . & \bar{R} + M & H + \bar{L} \end{pmatrix} \quad (40)$$

Since $S_2 \times S_2$ is isomorphic to D_2 , one might expect that the transformation matrix created from the projection of D_2 would diagonalize Equation (40). But this is not the case. Instead, its block diagonalizes the $S_2 \times S_2$ Hamiltonian.

$$T^+R(H)T = \begin{pmatrix} \langle 1| \\ \langle 2| \\ \langle 3| \\ \langle 4| \end{pmatrix} \begin{pmatrix} H + \bar{L} + \bar{R} + M & . & . & . \\ . & H + \bar{L} + \bar{R} + M & . & . \\ . & . & H - M & \bar{R} - \bar{L} \\ . & . & R - \bar{L} & H - M \end{pmatrix} \quad (41)$$

We construct a transformation Equation (42) that diagonalizes the Hamiltonian Equation (40) using the fact that $\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ diagonalizes a bilaterally symmetric matrix $\begin{pmatrix} A & B \\ B & A \end{pmatrix}$.

$$T = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 & . & . \\ 1 & -1 & . & . \\ . & . & 1 & 1 \\ . & . & 1 & -1 \end{pmatrix} \quad (42)$$

As a result, we obtained the following eigenvalues:

$$\begin{aligned} E_1 &= H + \bar{L} + \bar{R} + M \\ E_2 &= H + \bar{L} + \bar{R} + M \\ E_3 &= H - M + \bar{R} - \bar{L} \\ E_4 &= H - M - \bar{R} + \bar{L} \end{aligned} \quad (43)$$

However, the eigenvectors are just the columns of the transformation matrix. There is degeneracy since two of our eigenvalues are the same. The trace of Equation (38) gives

1	(12)	(34)	(12)(34)
4	0	2	2

(44)

Therefore this splits into $(2A_1, B_1, B_2)$ of the D_2 irreps.

5.2. Tunneling Hamiltonian for D_2 Point Symmetry

As the coupling is increased so that the two diatomic molecules become locked, we can have any of the other three cases in Figure 5. At this point, we are in BOA basis. The Bunker and Jensen work [23] is paramount for molecular symmetry groups. Although one could easily solve a rigid rotor problem with molecular symmetry, we use here symmetry projector theory [38,39], which allows for a sort of elegance when solving the coupled rotor system that is locked. Previously we saw that the D_{2h} fundamental vibration modes are $(A_{1g}, A_{2g}, B_{1u}, B_{2u})$. Here, the physics of D_{2h} and D_2 are essentially the same. The group table of D_2 is given by

1	R_y	R_x	R_z
R_y	1	R_z	R_x
R_x	R_z	1	R_y
R_z	R_x	R_y	1

(45)

and its matrix representation as follow:

$$\begin{pmatrix} 1 & . & . & . \\ . & 1 & . & . \\ . & . & 1 & . \\ . & . & . & 1 \end{pmatrix}, \begin{pmatrix} . & 1 & . & . \\ 1 & . & . & . \\ . & . & . & 1 \\ . & . & 1 & . \end{pmatrix}, \begin{pmatrix} . & . & 1 & . \\ . & . & . & 1 \\ 1 & . & . & . \\ . & 1 & . & . \end{pmatrix}, \begin{pmatrix} . & . & . & 1 \\ . & . & 1 & . \\ . & 1 & . & . \\ 1 & . & . & . \end{pmatrix} \quad (46)$$

Next we have given the projectors of D_2

$$\begin{aligned} P^{A_1} &\equiv P^A \cdot P^1 = \frac{(\mathbf{1} + R_z) \cdot (\mathbf{1} + R_y)}{2 \cdot 2} = \frac{1}{4}(\mathbf{1} + R_z + R_y + R_x) \\ P^{B_1} &\equiv P^B \cdot P^1 = \frac{(\mathbf{1} - R_z) \cdot (\mathbf{1} + R_y)}{2 \cdot 2} = \frac{1}{4}(\mathbf{1} - R_z + R_y - R_x) \\ P^{A_2} &\equiv P^A \cdot P^2 = \frac{(\mathbf{1} + R_z) \cdot (\mathbf{1} - R_y)}{2 \cdot 2} = \frac{1}{4}(\mathbf{1} + R_z - R_y - R_x) \\ P^{B_2} &\equiv P^B \cdot P^2 = \frac{(\mathbf{1} - R_z) \cdot (\mathbf{1} - R_y)}{2 \cdot 2} = \frac{1}{4}(\mathbf{1} - R_z - R_y + R_x) \end{aligned} \quad (47)$$

The tunneling Hamiltonian in the basis of a D_2 regular presentation is given as follows:

$$R(H) = \begin{pmatrix} \langle 1 | \\ \langle 2 | \\ \langle 3 | \\ \langle 4 | \end{pmatrix} \begin{pmatrix} H & M & L & R \\ M & H & R & L \\ L & R & H & M \\ R & L & M & H \end{pmatrix} \quad (48)$$

From Equation (47) we have the following diagonalizing transformation:

$$T = \frac{1}{2} \begin{pmatrix} & |A_1\rangle & |B_1\rangle & |A_2\rangle & |B_2\rangle \\ \langle 1 | & 1 & 1 & 1 & 1 \\ \langle 2 | & 1 & -1 & 1 & -1 \\ \langle 3 | & 1 & 1 & -1 & -1 \\ \langle 4 | & 1 & -1 & -1 & 1 \end{pmatrix} \quad (49)$$

Applying Equation (49) to Equation (48) we obtain the eigenvalues and eigenvectors in Equation (50):

$$\begin{aligned} E_{A_1} &= H + R + L + M : \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \quad E_{B_1} = H - M + R - L : \frac{1}{2} \begin{pmatrix} 1 \\ -1 \\ -1 \\ 1 \end{pmatrix}, \\ E_{A_2} &= H + M - R - L : \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ -1 \\ -1 \end{pmatrix}, \quad E_{B_2} = H - M - R + L : \frac{1}{2} \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix} \end{aligned} \quad (50)$$

The D_2 tunneling matrix produces no degeneracy in A_1 mode, as it was the case with $S_2 \times S_2$. This must be due to the fact that for $S_2 \times S_2$ structures, that an individual diatomic molecule is allowed to rotate out of the plane and it is the same symmetry configuration whether the diatomic molecule has one of its rotors out of the plane or the other one.

5.3. Tunneling Hamiltonian for D_4 Point Symmetry

Unlike D_2 , D_4 is non-Abelian since its group elements do not commute and neither will they be any faithful to the representation of D_4 . D_4 point group contains an identity operation, a 180° rotation around each of the x , y and z axis $\{R_z, R_y, R_x\}$, 90° and -90° about z -axis $\{R, R^3\}$, and finally 180° about each of the diagonals $\{i_3, i_4\}$. Non-commutivity of D_4 means that we cannot write all eight of the operators as a spectral decomposition using commuting idempotents. By idempotents, we mean $\mathcal{P}^\alpha \cdot \mathcal{P}^\alpha = \mathcal{P}^\alpha$ and zero otherwise. We will use *projector theory* to give a description of its non-Abelian algebra. Since it is non-commutative it is not possible to spectral decompose all of its eight operators by using eight commuting idempotents.

1	R_z	R	R^3	R_x	R_y	i_3	i_4
R_z	1	R^3	R	R_y	R_x	i_4	i_3
R^3	R	1	R_z	i_3	i_4	R_y	R_x
R	R^3	R_z	1	i_4	i_3	R_x	R_y
R_x	R_y	i_3	i_4	1	R_z	R^3	R
R_y	R_x	i_4	i_3	R_z	1	R	R^3
i_3	i_4	R_y	R_x	R^3	R	1	R_z
i_4	i_3	R_x	R_y	R	R^3	R_z	1

(51)

We obtain the D_4 regular representation from a group table Equation (51) rearranged so that columns are labeled by kets $|g\rangle = g|1\rangle$, and rows are labeled by bras

$$\langle g| = |g\rangle^+ = \langle 1|g^+ = \langle 1|g^{-1} \quad (52)$$

(Note that the inverses of the column operators sit on the left end of the rows).

Here each $R(g)$ matrix component is 1 (or 0) if g is (or is not) at that spot in Equation (51).

$$\begin{aligned}
 & \begin{pmatrix} 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \end{pmatrix}, \begin{pmatrix} & 1 & & & & & & \\ 1 & & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \end{pmatrix}, \begin{pmatrix} & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \end{pmatrix}, \begin{pmatrix} & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \end{pmatrix} \\
 & \quad R(1) \quad \quad R(R_z) \quad \quad R(R) \quad \quad R(R^3) \\
 & \begin{pmatrix} & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \\ 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \end{pmatrix}, \begin{pmatrix} & & & & & & 1 & \\ & & & & & & & 1 \\ & & & & & & & & 1 \\ & & & & & & & & & 1 \\ 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \end{pmatrix}, \begin{pmatrix} & & & & & & & 1 \\ & & & & & & & & 1 \\ & & & & & & & & 1 \\ & & & & & & & & & 1 \\ 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \end{pmatrix}, \begin{pmatrix} & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & & 1 \\ 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \end{pmatrix} \\
 & \quad R(R_x) \quad \quad R(R_y) \quad \quad R(i_3) \quad \quad R(i_4)
 \end{aligned} \quad (53a)$$

Since these representations are non-commuting, we cannot simply write the Hamiltonian in terms of them. Using the idea of mutually commuting lab based and body based operators g and \bar{g} , we can find the dual space. Harter [38] has given a detailed description of how to do this in general. In this case,

we find a body matrix representation $\{..R(\bar{g})..\}$ in Equation (53a) that commutes with in Equation (53b).

$$\begin{aligned}
 & \begin{pmatrix} 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \end{pmatrix} \begin{pmatrix} & 1 & & & & & & \\ 1 & & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \end{pmatrix} \begin{pmatrix} & & 1 & & & & & \\ & & & 1 & & & & \\ & 1 & & & & & & \\ & & & & & & 1 & \\ & & & & & & & 1 \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \end{pmatrix} \begin{pmatrix} & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \end{pmatrix} \\
& \quad R(1) \quad R(R_z) \quad R(R) \quad R(R^3) \\
& \begin{pmatrix} & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ 1 & & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \end{pmatrix} \begin{pmatrix} & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \end{pmatrix} \begin{pmatrix} & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \end{pmatrix} \begin{pmatrix} & & & & & & 1 & \\ & & & & & & & 1 \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \\ & & & & & & & \end{pmatrix} \\
& \quad R(R_x) \quad R(R_y) \quad R(i_3) \quad R(i_4)
\end{aligned} \tag{53b}$$

The group-based $R(g) D_4$ cannot be diagonalized but reduced or block diagonalized to a combination of irreps A_1, B_1, A_2, B_2 , and E from the character Table (4.13). The character of $R(g)$ is found by taking its trace, which gives $\{8,0,0,0,0\}$. The trace of $R(g)$ is $R(g) = A_1 \oplus B_1 \oplus A_2 \oplus B_2 \oplus 2E$. All eight $R(g)$ matrices are reducible by some transformation T to a block diagonals form with A_1, B_1, A_2, B_2 , and E blocks. Thus the reduction is carried out by finding all eight of the generalized projectors. This approach is described in more detail in [23,29,38,39] but will not be approached here.

The D_4 Hamiltonian is made by combining body based or dual space $R(\bar{g})$ representations in Equation (53a) since they commute with matrices Equation (53b) of the regular D_4 representation as required by D_4 symmetry. Thus we have,

$$R^G(H) = H\bar{1} + R\bar{R}_2 + S\bar{R} + S^*\bar{R}^3 + L\bar{R}_x + M\bar{R}_y + P\bar{i}_3 + Q\bar{i}_4 \tag{54a}$$

$$R^G(H) = \begin{pmatrix} \langle 1 | \\ \langle R_z | \\ \langle R | \\ \langle R^3 | \\ \langle R_x | \\ \langle R_y | \\ \langle i_3 | \\ \langle i_4 | \end{pmatrix} \begin{pmatrix} H & R & S & S^* & L & M & P & Q \\ R & H & S^* & S & M & L & Q & P \\ S^* & S & H & R & Q & P & L & M \\ S & S^* & R & H & P & Q & M & L \\ L & M & Q & P & H & R & S^* & S \\ M & L & P & Q & R & H & S & S^* \\ P & Q & L & M & S & S^* & H & R \\ Q & P & M & L & S^* & S & R & H \end{pmatrix} \tag{54b}$$

The first step in finding a transformation that would block diagonalized Equations (54a) and (54b) is to find the largest number of mutually commuting operators. This is also referred to as the Maximal Set of Commuting Operators (MSOCO). As pointed out by Harter [38], the number of operators is unique and it is the rank of the group. The rank of D_4 is six and the number of orthogonal irreducible representation (irreps) of the algebra is five, which means that there must exist another member of the set. The choice of this member is not unique. We choose an operator diagonal so we can split \mathcal{P}^E

idempotent in a particular way in order to build a particular set of $E - irreps$. In case of D_4 there exist three subgroups of order 4, five of order 2, and one of order 1, of which we can choose one as a temporary member to complete the maximal sets. For our computation we choose the subgroup $C_2 = \{1, R_x\}$ to split \mathcal{P}^E idempotent. Harter [38] described a procedure that we use to split \mathcal{P}^E idempotent into four new projectors.

Equation (51) is divided into its five classes as seen by how the table is sectioned off in blocks that form sort of a commuting algebra. Harter [38] has gone in detailed description of non-Abelian algebra where D_3 algebra is given as an example. Since in non-Abelian algebra, not all the matrices can be diagonalized or reduced at once by a single transformation, we use a technique of class commuting algebra described by Harter [38] that simultaneously spectral decompose MSOCO. The class algebra does not form a group but allows us to derive the projectors of the group. From the projectors we are able to derive such properties as the character table Equations (34a) and (34b), eigenvectors, and the transformation matrices, *etc.* Since, we already know the D_4 character table we will not here follow the exact approach of reference [38] because we can derive its projectors from Equations (34a) and (34b) as was done in (Section 4). The irreps of Equations (34a) and (34b) give five projectors, however, because the rank of $R(g)$ is 6 we need to split \mathcal{P}^E idempotent as discussed before. Now that we have all eight projectors, we derive a transformation Equation (55), which allows us to reduce the Hamiltonian Equation (55) to a block diagonal form as shown in Equation (56). From the projectors of D_4 we found the transformation matrix Equation (55).

$$T = \frac{1}{2\sqrt{2}} \begin{pmatrix} 1 & 1 & 1 & 1 & \sqrt{2} & 0 & 0 & \sqrt{2} \\ 1 & 1 & 1 & 1 & -\sqrt{2} & 0 & 0 & -\sqrt{2} \\ 1 & -1 & 1 & -1 & 0 & -\sqrt{2} & \sqrt{2} & 0 \\ 1 & -1 & 1 & -1 & 0 & \sqrt{2} & -\sqrt{2} & 0 \\ 1 & 1 & -1 & -1 & \sqrt{2} & 0 & 0 & -\sqrt{2} \\ 1 & 1 & -1 & -1 & -\sqrt{2} & 0 & 0 & \sqrt{2} \\ 1 & -1 & -1 & 1 & 0 & \sqrt{2} & \sqrt{2} & 0 \\ 1 & -1 & -1 & 1 & 0 & -\sqrt{2} & -\sqrt{2} & 0 \end{pmatrix} \quad (55)$$

This block diagonalized Equations (53a) and (53b) to give the following:

$$\langle H \rangle = \begin{pmatrix} H^{A_1} & . & . & . & . & . & . & . \\ . & H^{B_1} & . & . & . & . & . & . \\ . & . & H^{A_2} & . & . & . & . & . \\ . & . & . & H^{B_2} & . & . & . & . \\ . & . & . & . & H_{11}^E & H_{12}^E & . & . \\ . & . & . & . & H_{21}^E & H_{22}^E & . & . \\ . & . & . & . & . & . & H_{11}^E & H_{12}^E \\ . & . & . & . & . & . & H_{21}^E & H_{22}^E \end{pmatrix} \quad (56a)$$

$$\langle H \rangle = \begin{pmatrix} H^{A_1} & . & . & . & . & . & . & . \\ . & H^{B_1} & . & . & . & . & . & . \\ . & . & H^{A_2} & . & . & . & . & . \\ . & . & . & H^{B_2} & . & . & . & . \\ . & . & . & . & H_{11}^E & . & H_{21}^E & . \\ . & . & . & . & . & H_{11}^E & . & H_{21}^E \\ . & . & . & . & H_{12}^E & . & H_{22}^E & . \\ . & . & . & . & . & H_{12}^E & . & H_{22}^E \end{pmatrix} \quad (56b)$$

The reduction of Equations (54a) and (54b) using the above transformation gives four singlet eigenvalues

$$\begin{aligned} H^{A_1} &= H + S^* + L + M + P + Q + R + S \\ H^{B_1} &= H - S^* + L + M - P - Q + R - S \\ H^{A_2} &= H + S^* - L - M - P - Q + R + S \\ H^{B_2} &= H - S^* - L - M + P + Q + R - S \end{aligned} \quad (57)$$

Finally, there are two pairs of identical two by two matrices each with the following matrix element:

$$\begin{aligned} H_{11}^E &= H + L - M - R; & H_{12}^E &= -S^* + P - Q + S \\ H_{22}^E &= H - L + M - R; & H_{21}^E &= S^* + P - Q - S. \end{aligned} \quad (58a)$$

These two by two matrices will have to be further diagonalized if Q , P or ImS are non-zero. Therefore, if we let $S = \rho + i\nu$ then we can represent these two-by-two matrices as

$$\begin{pmatrix} H_{11}^E & H_{12}^E \\ H_{21}^E & H_{22}^E \end{pmatrix} = \begin{pmatrix} H - R + L - M & -2\left(\frac{Q-P}{2} - i\nu\right) \\ -2\left(i\nu + \frac{Q-P}{2}\right) & H - R - (L - M); \end{pmatrix} \quad (58b)$$

If instead we apply the transformation Equation (55) using the regular representation $R(g)$ instead of $R(\bar{g})$ then it would be block diagonalized like in Equation (56b).

To interpret the eigensolutions of Equation (58b), we represent our two-by-two matrix in terms of the quasi-spin vectors (S_A, S_B, S_C) as defined in Figure 7.

$$\begin{pmatrix} H_{11}^E & H_{12}^E \\ H_{21}^E & H_{22}^E \end{pmatrix} = (H - R)\sigma_0 + 2(Q - P) \frac{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}}{2} + 4\nu \frac{\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}}{2} + 2(L - M) \frac{\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}}{2} \quad (59)$$

Equation (59) is analogous to the description of a two state system, therefore U_2 analysis may help to characterize the eigensolution. Harter [38,39] have given a general development U_2 analysis for any two states system. Suppose $\begin{pmatrix} H_{11}^E & H_{12}^E \\ H_{21}^E & H_{22}^E \end{pmatrix}$ has the form $\begin{pmatrix} A & B \\ B & D \end{pmatrix}$ with no component on the C axis.

Also let $A = D$ so both L and M are zero. Then there results a simple mixing between the E modes in the two by two blocks with eigenvectors $(\frac{1}{\sqrt{2}}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix})$ diagonalizing bilaterally symmetric $\begin{pmatrix} H-R & P-Q \\ P-Q & H-R \end{pmatrix}$ sub-Hamiltonians.

Thus the E-Hamiltonian under these conditions has a perfect bilateral symmetry. This means that if the quasi-spin vector starts out on the B axis as in Figure 7, it remains fixed. However, when $L \neq M$, we have an asymmetric two-state system that is very much like the one in reference [39]. This is plotted in Figure 8.

Figure 7. (ABC) symmetry coordinates. This diagram was published by Harter [38].





Figure 8. Two state system eigensolutions [39].



6. Conclusions

The frame relation derived in reference [25] was developed to include the finite symmetry of a molecular geometry. The B label was shown to be the symmetry species of the nascent geometry's point group. The transformation relation has a finite symmetry field coefficient of the small rotor included in it. It is not clear how one should choose a particular frame over. However, in a locked geometry, the choice of frame does not matter, but only the geometric point group that is formed. We investigated locked geometries for two diatomic molecules using Young tableau and unitary algebra to find the nuclear spin statistical weights. This gives approximates and preliminary insight for level structure.

Moreover, we used the frame transformation relations derived in reference [25] to calculate the transitions between angular momentum states. More general transition relations were found than those given in reference [12]. However, one could extend our relation to include symmetry for transitions in a two rotor coupled system, but this will be left for future work.

Acknowledgments

This work has been co-supported by the Research Corporation for Science Advancement.

References

1. Young, A. On quantitative substitutional analysis IX. *Proc. Lond. Math. Soc.* **1952**, *54*, 219–253.
2. Yamanouchi, T. On the construction of unitary irreducible representation of the symmetric group. *Proc. Phys. Math. Soc. Jpn.* **1937**, *19*, 436–450.
3. Yamanouchi, T. *The Application of the Rotational Group and Symmetric Group*; Beijing Science Press: Beijing, China, 1966.
4. Littlewood, D.E. *The Theory of Group Characters and Matrix Representations of Groups*; Oxford Press: Oxford, UK, 1950.
5. Gel'fand, M.I. The center of an infinitesimal group algebra. *Mat. Sb.* **1950**, *26*, 103–112.
6. Gel'fand, M.I. Matrix elements for the unitary groups. *Dokl. Akad. Nauk.* **1950**, *71*, 825–828.
7. Gel'fand, M.I. *Representations of the Rotation and Lorentz and Their Applications*; Pergamon Press: New York, NY, USA, 1963.
8. Shur, I. *Neue anwendungen der Integralrechnung auf Probleme der Invariantentheorie*; Sitzungsberichte Akademie der Wissenschaften: Berlin, Germany, 1924; pp. 189–208.
9. de Beauregard Robinson, G. *Representation Theory of the Symmetric Group*; University of Toronto Press: Toronto, Canada, 1960.
10. Coleman, A.J. *Induced Representations and Applications*; Queens University Press: Kingston, Ontario, Canada, 1971.
11. Harter, W. Theory of hyperfine and super levels in symmetric polyatomic molecules. II Elementary cases in octahedral hexafluoride molecules. *Phys. Rev. A* **1981**, *24*, 192–263.
12. Harter, W.G.; Patterson, C.W.; da Paixao, F.J. Frame transformation relations and multiple transitions in symmetric polyatomic molecules. *Rev. Mod. Phys.* **1978**, *50*, 37–83.

13. Harter, W.; Patterson, C.W. Theory of hyperfine and super levels in symmetric polyatomic molecules. Trigonal and tetrahedral: Elementary spin $\frac{1}{2}$ cases vibronic ground states. *Phys. Rev. A* **1979**, *19*, 2277–2303.
14. Streater, R.F.; Wightman, A.S. *PCT, Spin and Statistics, and All That*; Princeton University Press: Princeton, NJ, USA, 1964.
15. Longuet-Higgins, H.C. The symmetry groups of non-rigid molecules. *Mol. Phys.* **1963**, *6*, 445–460.
16. Hougen, J.T. Classification of rotational energy levels for symmetric—Top molecules. *J. Chem. Phys.* **1962**, *37*, 1433–1441.
17. Harter, W.G. Alternative basis for the theory of complex spectra. *Phys. Rev. A* **1973**, *8*, 2819–2827.
18. Harter, W.G.; Patterson, C.W. Alternative basis for the theory of complex spectra. II. *Phys. Rev. A* **1976**, *13*, 1067–1082.
19. Patterson, C.W.; Harter, W.G. Canonical symmetrization for unitary bases. I. Canonical Weyl bases. *J. Math. Phys.* **1976**, *17*, 1125–1136.
20. Harter, W.G.; Patterson, C.W. Simple model for asymptotic level clusters in SF₆ rotational spectra. *Phys. Rev. Lett.* **1977**, *5*, 224–227.
21. Harter, W.G.; Weeks, D.E. Rotation-vibration spectra of icosahedral molecules. I. Icosahedral symmetry analysis and fine structure. *J. Chem. Phys.* **1989**, *90*, 4727–4743.
22. Harter, W.G.; Patterson, C.W. Alternative basis for the theory of complex spectra. III. *Phys. Rev. A* **1976**, *15*, 2372–2379.
23. Bunker, P.R.; Jensen, P. *Molecular Symmetry and Spectroscopy*, 2nd ed.; NRC Research Press: Ottawa, Canada, 1998.
24. Groner, P. Effective rotational Hamiltonian for molecules with two periodic large-amplitude motions. *J. Chem. Phys.* **1997**, *107*, 4483–4498.
25. Croghan, H.T.; Harter, W.G. Frame transformation relations for fluxional symmetric rotor dimmers. *J. Chem. Phys.* **2004**, *121*, 9297–9312.
26. Chang, E.S.; Fano, U. Theory of electron-molecule collisions by frame transformations. *Phys. Rev. A* **1972**, *6*, 173–185.
27. Junger, C.H. Highly excited molecular states: Quantum defect and *ab initio* theory. In *Proceedings of International Symposium on Molecular and Spectroscopy*, Ohio State University, Columbus, OH, USA, 16–20 June 2003.
28. Junger, C.H.; Atabek, O. Rovibronic interactions in the photoabsorption spectrum of molecular hydrogen and deuterium: An application of multichannel quantum defect methods. *J. Chem. Phys.* **1976**, *66*, 5584–5609.
29. Chen, J.Q. *Group Representation Theory for Physicist*; World Scientific Publishing Co. Pte. Ltd.: Singapore, 1989.
30. Kung, P.S.J. *Young Tableaux in Combinatorics Invariant Theory and Algebra*; Academic Press: University of Michigan, Ann Arbor, MI, USA, 1982.
31. Lemus, R. A general method to obtain vibrational symmetry adapted bases in a local scheme. *Mol. Phys.* **2003**, *101*, 2511–2528.

32. Crogman, H.T. A rotational energy surface study for low polyad structures of the general rovibrational Hamiltonian. *Mol. Phys.* **2010**, *108*, 705–721.
33. Crogman, H.T.; Harter, W.G. The emergence of a single frame and effects on quantum states and levels for rotor-rotor interactions. In *Proceedings of 59th Ohio State University International Symposium on Molecular Spectroscopy*, Ohio State University, Columbus, OH, USA, 21–24 June 2004.
34. Hougen, J.T. Classification of rotational energy levels. II. *J. Chem. Phys.* **1971**, *39*, 358–365.
35. Alvarez-Bajo, O.; Lemus, R.; Carvajal, M.; Perez-Bernal, F. Equivalent rotations associated with the permutation inversion group revisited: Symmetry projection of the rotational functions of methane. *Mol. Phys.* **2011**, *109*, 797–812.
36. Lederer, C.M.; Hollander, J.M.; Perlman, I. *Table of Isotopes*, 6th ed.; Wiley: New York, NY, USA, 1967.
37. Hougen, J.T.; Merer, A.J. Extended permutation-inversion groups for simultaneous treatment of the rovibronic states of *trans*-acetylene, *cis*-acetylene, and vinylidene. *J. Mol. Spectrosc.* **2011**, *267*, 200–221.
38. Harter, W.G. *The Quantum Computer Age*; University of Arkansas: Fayetteville, NC, USA, 2004.
39. Harter, W.G. *Principle of Symmetry, Dynamics & Spectroscopy*; Wiley: New York, NY, USA, 1993.

© 2013 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).