

## Interesting Images

# Fatal Subarachnoid Hemorrhage in a Deep Brain Stimulation Patient: Displacement of Stimulation Leads for Deep Brain Stimulation Indicate Subarachnoid Hemorrhage on X-ray

Gregor Bara <sup>\*</sup>, Valeri Borger  and Jaroslaw Maciaczyk 

Department of Neurosurgery, University Hospital Bonn, Venusberg-Campus 1, Building 81, 53127 Bonn, Germany  
\* Correspondence: gregor.bara@ukbonn.de; Tel.: +49-228-287-16518

**Abstract:** We depict the rare case of a patient with aneurysmatic subarachnoid hemorrhage previously treated with deep brain stimulation for Parkinson's disease. Initial CT scans showed a Fisher grade 4 subarachnoid hemorrhage with lead displacement due to midline-shift. CT angiogram revealed a supra-ophthalmic aneurysm of the internal carotid artery. The patient subsequently underwent clipping of the aneurysm and decompressive hemicraniectomy.

**Keywords:** deep brain stimulation; subarachnoid hemorrhage; intracerebral hemorrhage



**Citation:** Bara, G.; Borger, V.; Maciaczyk, J. Fatal Subarachnoid Hemorrhage in a Deep Brain Stimulation Patient: Displacement of Stimulation Leads for Deep Brain Stimulation Indicate Subarachnoid Hemorrhage on X-ray. *Diagnostics* **2024**, *14*, 222. <https://doi.org/10.3390/diagnostics14020222>

Academic Editor: Andreas Kjaer

Received: 18 December 2023

Revised: 16 January 2024

Accepted: 18 January 2024

Published: 19 January 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

A 70-year-old patient was admitted to our emergency room with a history of sudden loss of consciousness and epileptic seizure. An initial GCS (Glasgow Coma Scale) was 3, and pupils were isochoric with intact direct and consensual light reflex. Sixteen years prior, he had been implanted with bilateral stimulation leads for deep brain stimulation of the subthalamic nucleus due to Parkinson's disease. An initial head CT scan (Philips Icon, 1mm head scan) showed a predominantly left-sided subarachnoid hemorrhage (SAH) Fisher grade 4 with intra-ventricular hemorrhage, intra-parenchymal hemorrhage within the left frontal lobe, subdural hematoma, and midline-shift of 12 mm. The SAH was classified as WFNS (World Federation of Neurological Surgeons) grade 5. The CT angiogram (Philips Icon, 1mm head scan) of the supra-aortal vessels revealed a left-sided supra-ophthalmic aneurysm of the internal carotid artery as the supposed source of hemorrhage. An external ventricular drain was placed and the patient subsequently brought to the operating room for clipping of the aneurysm and decompressive hemicraniectomy. During the latter, the left-sided DBS stimulation lead had to be removed. Eventually, the therapy was switched to best supportive care due to massive parenchyma destruction.

A subarachnoid hemorrhage (SAH) is an extravasation of blood into the space between the arachnoid membrane and the pia mater filled with cerebrospinal fluid. The most common cause of a non-traumatic SAH is due to a ruptured intracranial aneurysm [1].

Whilst aneurysmal SAH can be a devastating condition associated with neurological complications up to death, nonaneurysmal SAH is associated with a good prognosis and is rarely associated with neurological complications [2].

Diagnosis of a SAH is primarily performed by computed tomography (CT) of the head, in which the SAH appears characteristically with hyperdense extravasated blood sometimes associated with intraparenchymal hematoma and hydrocephalus [3,4].

In this case, however, first signs of the SAH could be identified in the initial head X-ray of the CT scout image due to the patient having been implanted with bilateral stimulation leads for deep brain stimulation (Figure 1). The differential diagnosis of an intracranial mass affect includes pathologies of traumatic (epidural hematoma, acute subdural hematoma, chronic subdural hematoma, traumatic intracerebral hemorrhage), tumorous (brain derived tumors such as glioblastoma multiforma or meningioma, metastases) as well as vascular origin (subarachnoid hemorrhage, intracranial hemorrhage). Table 1 depicts the most

common differential diagnoses in respect to pathological origin, entity, CT morphology and clinical dynamic. In this particular case, the CT scan revealed the brain shift to be due to the subarachnoid hemorrhage with an associated hematoma which displaced the stimulation leads to the contralateral side (Figure 2). The CT angiogram showed a supra-ophthalmic aneurysm of the internal carotid artery to be the source of hemorrhage (Figure 3).



**Figure 1.** Initial scout image of the head CT scan, essentially mimicking an AP cranial X-ray. Note the implanted stimulations leads for deep brain stimulation, which are connected to an infra-clavicular neurostimulator. The DBS stimulation leads are displaced to the right side.

The amount of blood within the subarachnoid space and the ventricles, as well as the parenchyma, can be easily graded on a head CT scan and is correlated with poor outcome [5,6]. Further, the prognosis is correlated to the initial clinical exam which is most widely measured on the clinical scale of Hunt and Hess and the World Federation of Neurological Surgeons [7,8]. Given the clinical and radiological presentation of this patient, this was defined as poor grade.

**Table 1.** Differential diagnosis of mass effect.


Pathological Class	Pathology	CT Morphology	Clinical Dynamic
Traumatic	Epidural hematoma	Hyperdense biconvex extra-axial	Minutes to hours
	Acute subdural hematoma	Hyperdense crescent-shaped extra-axial spreading diffusely over the affected hemisphere	Minutes to hours
	Chronic subdural hematoma	Hypodense crescent-shaped extra-axial spreading diffusely over the affected hemisphere septation and sediment effect	Weeks
	Traumatic intracerebral hemorrhage	Hyperdense Intra-axial Usually in combination with traumatic subdural hematoma and/or acute subdural hematoma	Minutes to hours
Tumorous	Brain derived such as glioblastoma multiforme, meningioma	GBM: intra-axial mass with thick, irregularly enhancing margins with a central necrotic core, possibly hemorrhagic Meningeoma: extra-axial mass, well-circumscribed, contract to dura	Weeks to months
	Metastasis	Intra-axial Subcortical Usually multiple	Weeks to months
Vascular	Subarachnoid hemorrhage	Hyperdense Usually extra-axial within the subarachnoid space and cisterns, but intracerebral hemorrhage may occur Hydrocephalus may occur	Sudden onset
	Intracerebral hemorrhage	Hyperdense Intra-axial Usually thalamus, caudate nucleus and pons Hydrocephalus may occur	Sudden onset

Once an aneurysm has been identified by CT angiogram and/or digital subtraction angiography, treatment can be performed either with endovascular coiling or microsurgical clipping [9].

Further complications can be managed by placement of an external ventricular drain for hydrocephalus and decompressive hemicraniectomy in case of otherwise uncontrollable intracranial hypertension, as performed in this case report (Figure 4) [10].



**Figure 2.** Initial head CT scan shows the predominantly left-sided subarachnoid hemorrhage with intra-ventricular hemorrhage, subdural hematoma, and midline-shift. The DBS stimulation leads are displaced to the contralateral side.



**Figure 3.** A 3D reconstruction of the CT angiogram showing a supra-ophthalmic aneurysm of the internal carotid artery as the source of hemorrhage (marked by the arrow). DBS leads are displaced to the contralateral side.



**Figure 4.** Postoperative head CT scan after clipping of the supra-ophthalmic aneurysm, left-sided DBS lead removal and decompressive hemicraniectomy. Eventually, therapy was switched to best supportive care due to massive parenchyma destruction.

**Author Contributions:** Conceptualization, G.B.; methodology, G.B.; investigation, G.B.; data curation, G.B.; writing—original draft preparation, G.B.; writing—review and editing, V.B. and J.M.; supervision, J.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board of the University of Bonn (protocol code 204/21 and date of 17 August 2021).

**Informed Consent Statement:** Due to fatality, patient could not give his consent. Written informed consent has been obtained from patient's next of kin to publish this paper.

**Data Availability Statement:** No new data was created.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. van Gijn, J.; Rinkel, G.J. Subarachnoid haemorrhage: Diagnosis, causes and management. *Brain* **2001**, *124*, 249–278. [[CrossRef](#)] [[PubMed](#)]
2. Adams, H.P., Jr.; Gordon, D.L. Nonaneurysmal subarachnoid hemorrhage. *Ann. Neurol.* **1991**, *29*, 461–462. [[CrossRef](#)] [[PubMed](#)]
3. Latchaw, R.E.; Silva, P.; Falcone, S.F. The role of CT following aneurysmal rupture. *Neuroimaging Clin. N. Am.* **1997**, *7*, 693–708. [[PubMed](#)]
4. Hijdra, A.; van Gijn, J.; Nagelkerke, N.J.; Vermeulen, M.; van Crevel, H. Prediction of delayed cerebral ischemia, rebleeding, and outcome after aneurysmal subarachnoid hemorrhage. *Stroke* **1988**, *19*, 1250–1256. [[CrossRef](#)] [[PubMed](#)]
5. Claassen, J.; Bernardini, G.L.; Kreiter, K.; Bates, J.; Du, Y.E.; Copeland, D.; Connolly, E.S.; Mayer, S.A. Effect of cisternal and ventricular blood on risk of delayed cerebral ischemia after subarachnoid hemorrhage: The Fisher scale revisited. *Stroke* **2001**, *32*, 2012–2020. [[CrossRef](#)] [[PubMed](#)]
6. Fisher, C.M.; Kistler, J.P.; Davis, J.M. Relation of cerebral vasospasm to subarachnoid hemorrhage visualized by computed tomographic scanning. *Neurosurgery* **1980**, *6*, 1–9. [[CrossRef](#)] [[PubMed](#)]
7. Hunt, W.E.; Hess, R.M. Surgical risk as related to time of intervention in the repair of intracranial aneurysms. *J. Neurosurg.* **1968**, *28*, 14–20. [[CrossRef](#)] [[PubMed](#)]
8. Drake, C.G. Report of World Federation of Neurological Surgeons committee on a universal subarachnoid hemorrhage grading scale. *J. Neurosurg.* **1988**, *68*, 985–986.
9. Molyneux, A.; Kerr, R.; Yu, L.-M.; Clarke, M.; Sneade, M.; Yarnold, J.; Stratton, I. International Subarachnoid Aneurysm Trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: A randomised trial. *Lancet* **2002**, *360*, 1267–1274. [[CrossRef](#)] [[PubMed](#)]
10. Dorfer, C.; Frick, A.; Knosp, E.; Gruber, A. Decompressive Hemicraniectomy after Aneurysmal Subarachnoid Hemorrhage. *World Neurosurg.* **2010**, *74*, 465–471. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.