

Article

The Myrmecofauna (Hymenoptera: Formicidae) of Hungary: Survey of Ant Species with an Annotated Synonymic Inventory

Sándor Csősz ^{1,2}, Ferenc Báthori ², László Gallé ³, Gábor Lőrinczi ⁴, István Maák ^{4,5}, András Tartally ^{6,*} , Éva Kovács ⁷, Anna Ágnes Somogyi ⁶  and Bálint Markó ^{8,9} 

¹ MTA-ELTE-MTM Ecology Research Group, Pázmány Péter sétány 1/C, 1117 Budapest, Hungary; sandorcsosz2@gmail.com

² Evolutionary Ecology Research Group, Centre for Ecological Research, Institute of Ecology and Botany, 2163 Vácrátót, Hungary; ferenc.bathori@gmail.com

³ Department of Ecology and Natural History Collection, University of Szeged, Szeged Boldogasszony sgt. 17., 6722 Szeged, Hungary; galle@bio.u-szeged.hu

⁴ Department of Ecology, University of Szeged, Közép fasor 52, 6726 Szeged, Hungary; lorinczig@gmail.com (G.L.); bikmakk@gmail.com (I.M.)

⁵ Museum and Institute of Zoology, Polish Academy of Sciences, ul. Wilcza 64, 00-679 Warsaw, Poland

⁶ Department of Evolutionary Zoology and Human Biology, University of Debrecen, Egyetem tér 1, 4032 Debrecen, Hungary; panka.somogyi@gmail.com

⁷ Kiskunság National Park Directorate, Liszt F. u. 19, 6000 Kecskemét, Hungary; kovacse@knp.hu

⁸ Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Clinicilor 5-7, 400006 Cluj-Napoca, Romania; balintm@gmail.com

⁹ Centre for Systems Biology, Biodiversity and Bioresources, Babeş-Bolyai University, Clinicilor 5-7, 400006 Cluj-Napoca, Romania

* Correspondence: tartally.andras@science.unideb.hu; Tel.: +36-52-512-900 (ext. 62349)



Citation: Csősz, S.; Báthori, F.; Gallé, L.; Lőrinczi, G.; Maák, I.; Tartally, A.; Kovács, É.; Somogyi, A.Á.; Markó, B. The Myrmecofauna (Hymenoptera: Formicidae) of Hungary: Survey of Ant Species with an Annotated Synonymic Inventory. *Insects* **2021**, *12*, 78. <https://doi.org/10.3390/insects12010078>

Received: 4 December 2020

Accepted: 13 January 2021

Published: 16 January 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Simple Summary: Abundance is a hallmark of ants (Hymenoptera: Formicidae). They are exceedingly common in both natural and artificial environments and they constitute a conspicuous part of the terrestrial ecosystem; every 3 to 4 out of 10 kg of insects are given by ants. Due to their key role in natural habitats, they are at the basis of any nature conservation and pest management policy. Thus, the first step in developing adequate management strategies is to build a precise faunistic inventory. More than 16,000 valid ant species are registered worldwide, of which 126 are known to occur in Hungary. Thanks to the last decade's efforts in the Hungarian myrmecological research, and because of the constantly changing taxonomy of several problematic ant genera, a new checklist of the Hungarian ants is presented here. A comparison of the Hungarian myrmecofauna to other European countries' ant fauna is also provided in this paper. The current dataset is a result of ongoing work on inventorying the Hungarian ant fauna, therefore it is expected to change over time and will be updated once the ongoing taxonomic projects are completed.

Abstract: Ants (Hymenoptera: Formicidae) are exceedingly common in nature. They constitute a conspicuous part of the terrestrial animal biomass and are also considered common ecosystem engineers. Due to their key role in natural habitats, they are at the basis of any nature conservation policy. Thus, the first step in developing adequate conservation and management policies is to build a precise faunistic inventory. More than 16,000 valid ant species are registered worldwide, of which 126 are known to occur in Hungary. Thanks to the last decade's efforts in the Hungarian myrmecological research, and because of the constantly changing taxonomy of several problematic ant genera, a new checklist of the Hungarian ants is presented here. The state of the Hungarian myrmecofauna is also discussed in the context of other European countries' ant fauna. Six species (*Formica lemani*, *Lasius nitidigaster*, *Tetramorium immigrans*, *T. staerckeii*, *T. indocile* and *Temnothorax turcicus*) have been reported for the first time in the Hungarian literature, nine taxon names were changed after systematic replacements, nomenclatorial act, or as a result of splitting formerly considered continuous populations into more taxa. Two species formerly believed to occur in Hungary are now excluded from the updated list. All names are nomenclaturally assessed, and complete synonymies applied in the Hungarian literature for a certain taxon are provided. Wherever it is not self-evident, comments

are added, especially to explain replacements of taxon names. Finally, we present a brief descriptive comparison of the Hungarian myrmecofauna with the ant fauna of the surrounding countries. The current dataset is a result of ongoing work on inventorying the Hungarian ant fauna, therefore it is expected to change over time and will be updated once the ongoing taxonomic projects are completed.

Keywords: ants; biogeography; faunistics; checklist; Europe

1. Introduction

Faunistic papers and regional checklists are cornerstones of nature conservation. Without up-to-date faunistic information, it is almost impossible to formulate adequate conservation strategies for species or specific habitats [1–3]. As a drastic decline in species number in the past decades is widely acknowledged [4,5], public interest for biodiversity is constantly growing, and there is an increasing demand for accurate faunistic information also from non-academics. This is proven by the existence of numerous specific interest groups on social media bringing together specialists and amateurs alike. Citizen science, the implication of volunteering non-academics in scientific research, could be essential for inventorying biota [6], saving habitats, protecting species [7,8], describing new taxa [9], and detecting new fauna elements [10,11]. It could also help overcome to a certain degree the ongoing crisis of taxonomic impediment, namely the lack of specialists [12]. Thus, a periodical update of any checklist should be considered as almost mandatory for assisting conservation policies. In the frame of the current study, we offer an updated version of the Hungarian ant fauna with necessary corrections and also additions due to novel findings in the past years.

The myrmecofauna of central and eastern Europe is relatively well-known as over the course of the past few decades several new checklists, faunistic monographs and keys have been published for this region and generally for Europe [13–23]. The last checklist of Hungarian ants was compiled in 2011 and reported 126 species [24].

The myrmecofauna of present-day Hungary has been studied in considerable detail over the course of the last few centuries, and several checklists have been published since the mid-1800s. The first detailed checklist containing locations was published by Mayr [25]. He listed 40 species for the territory of present-day Hungary. His work was later updated toward the end of the 19th century [26], resulting in a checklist that contained 56 species for the territory of present-day Hungary. More than 50 years later the monograph of Somfai [27] listed still just 66 species, which could be considered quite low for this region even by the standards of that time. The first checklist that, indeed, passed the threshold of 100 species, was published in Gallé et al. [28] that already contained 101 species, and also resolved many taxonomic and faunistic inconsistencies. The number of species increased considerably in the last checklist of Hungarian ants published in Csósz et al. [24], containing 122 ant species, mostly due to the increase in the number of myrmecologist scholars and also their faunistic and taxonomical works [29].

Despite this improvement over the last few decades, since the publication of the latest Hungarian checklist, many new taxa have been described [30,31], revived from synonymy [32], or the validity of their status has been confirmed or even dismissed on the basis of new findings [15,23]. Furthermore, several new myrmecological studies have been carried out reporting new ant species for the Hungarian fauna and new data regarding species already known [29,33]. Consequently, there is a need for an updated checklist of the Hungarian myrmecofauna, based on the most up to date taxonomic knowledge.

2. Materials and Methods

The current list of species was prepared by using every major faunistic paper concerning the territory of current Hungary, as well as other papers handling Hungarian data.

For the up-to-date nomenclature, taxonomy, and systematics of ants, we followed Bolton's Catalogue and Synopsis [34]. Most species were collected over the course of the recent decade, but there are some records that could not be verified, neither by checking voucher specimens nor by collecting from the sample sites.

3. Results

The total number of species is 126 belonging to 33 genera, out of which six are new elements for the Hungarian fauna: *Formica lemani*, *Lasius nitidigaster*, *Tetramorium immigrans*, *T. staerkei*, *T. indocile* and *Temnothorax turcicus* have been identified after the last check-list [24]. Five species of the exotic origin occur exclusively indoors: *Hypoponera punctatissima*, *Monomorium pharaonis*, *Tetramorium bicarinatum*, *T. insolens* and *Tapinoma melanocephalum*. Two species, *Temnothorax rabaudi* and *Tetramorium impurum*, are considered to be of uncertain occurrence, hence these species are now excluded from the Hungarian myrmecofauna, as voucher specimens or recent investigations do not support their presence.

3.1. Subfamily: Proceratiinae

3.1.1. Tribe: Proceratiini

Genus: *Proceratium*

P. melinum (ROGER, 1860)

= *Sysphincta fialai* KRATOCHVIL, 1944: [27]

3.2. Subfamily: Ponerinae

3.2.1. Tribe: Ponerini

Genus: *Cryptopone*

C. ochracea (MAYR, 1855)

= *Cryptopone ochraceum*: [35]

Genus: *Hypoponera*

H. punctatissima (ROGER, 1859): [28] (Almost exclusively an indoor tramp species)

= *Ponera punctatissima*: [27]

Genus: *Ponera*

P. coarctata (LATREILLE, 1802): [25]

P. testacea EMERY, 1895: [36]

3.3. Subfamily: Myrmicinae

3.3.1. Tribe: Attini

Genus: *Strumigenys*

S. argiola (EMERY, 1869)

= *Pyramica argiola*: [24]

= *Epitritus argiolus*: [26]

S. baudueri (EMERY, 1875)

= *Pyramica baudueri*: [24]

= *Strumigenys baudueri*: [27]

= *Smithistruma baudueri*: [28]

3.3.2. Tribe: Crematogastrini

Genus: *Cardiocondyla*

C. dalmatica SOUDEK, 1925: [23]

= *Cardiocondyla sahlbergi* FOREL, 1913: [28] (erroneous determination)

= *Cardiocondyla elegans* EMERY, 1869: [24] (erroneous determination)

Genus: *Crematogaster*

- C. schmidti* (MAYR, 1853): [28]
C. scutellaris (OLIVIER, 1792): [37]
C. sordidula (NYLANDER, 1849): [37]

Genus: *Formicoxenus*

- F. nitidulus* (NYLANDER, 1846): [27]

Genus: *Harpagoxenus*

- H. sublaevis* (NYLANDER, 1849): [27]

Genus: *Leptothorax*

- L. acervorum* (FABRICIUS, 1783): [27]
L. gredleri MAYR, 1855: [27]
L. muscorum (NYLANDER, 1846): [27]

Genus: *Myrmecina*

- M. graminicola* (LATREILLE, 1802): [27]
= *Myrmecina latreillei* CURTIS, 1829: [25]

Genus: *Strongylognathus*

- S. testaceus* (SCHENCK, 1852): [25]

Genus: *Temnothorax*

- T. affinis* (MAYR, 1855)
= *Leptothorax affinis* MAYR, 1855: [27]
T. albipennis (CURTIS, 1854): [24]
T. clypeatus (MAYR, 1853)
= *Leptothorax clypeatus*: [27]
T. corticalis (SCHENCK, 1852)
= *Leptothorax corticalis*: [27]
T. crassispinus (KARAVAJEV, 1926)
= *Leptothorax nylanderi*: [25]
= *Leptothorax slavonicus* SEIFERT, 1995: [38]
T. interruptus (SCHENCK, 1852)
= *Leptothorax interruptus* MAYR, 1855: [26]
= *Leptothorax tuberum* var. *interruptus* ANDRÉ, 1881: [27]
T. jailensis (ARNOLDI, 1977): [24]
T. nigriceps (MAYR, 1855)
= *Leptothorax nigriceps* MAYR, 1855: [39]
T. parvulus (SCHENCK, 1852)
= *Leptothorax parvulus* (SCHENCK, 1852): [27]
T. sordidulus (MÜLLER, 1923)
= *Leptothorax sordidulus*: [39]
T. tuberum (FABRICIUS, 1775)
= *Leptothorax tuberum* MAYR, 1855: [37]
T. turcicus (SANTSCHI, 1934): new record, Iszkaszentgyörgy, 47.236382 N, 18.284685 E, 187 m, leg. S. Csósz, 06.06.2020 (3 workers), Budapest, Rupp-hegy, 47.4730 N, 18.9794 E, 197 m, leg. S. Csósz, 04.05.2018 (6 workers, 2 gynes), Budapest, Rupp-hegy, 47.4730 N, 18.9794 E, 197 m, leg. S. Csósz, 17.06.2019 (3 workers), Mátraháza, 47.850518 N, 19.959551 E, 584 m, leg. S. Csósz, 04.08.2019 (3 workers)
T. unifasciatus (LATREILLE, 1798): [40]
= *Leptothorax unifasciatus*: [25]
= *Leptothorax tuberum* var. *unifasciata* ANDRÉ, 1881: [40]

T. zaleskyi (SADIL 1953)
 = *Myrmoxenus ravouxi* (ANDRÉ, 1896)
 = *Epimyrma goesswaldi* MENOZZI, 1930: [41]
 = *Epimyrma ravouxi*: [38]

Genus: *Tetramorium*

T. atratum: [25]
 = *Anergates atratus* (SCHENCK, 1952): [26]
T. bicarinatum (NYLANDER, 1846): [26] (Indoor tramp species)
T. caespitum (LINNAEUS, 1758): [42]
T. ferox Ruzsky, 1903: [43]
T. hungaricum RÖSZLER, 1935
 = *Tetramorium caespitum hungaricum* RÖSZLER, 1935: [44]
 = *Tetramorium hungaricum*: [45,46]
T. immigrans SANTSCHI, 1927
 = *Tetramorium* sp. E: [24,47]
T. indocile SANTSCHI 1927: [48]
 = *Tetramorium* sp. C: [24,47]
T. insolens (SMITH, 1861): [27] (Indoor tramp species)
T. moravicum KRATOCHVÍL, 1941
 = *Tetramorium rhenanum* SCHULZ 1996: [28]
 = *Tetramorium moravicum*: [49]
T. semilaeve (ANDRÉ, 1883): [28]
 = *Tetramorium simillimum* (SMITH, 1851): [28] (erroneous determination)
T. staercke KRATOCHVÍL, 1944
 = *Tetramorium* sp. D: [24,47]

3.3.3. Tribe: Myrmicini

Genus: *Manica*

M. rubida (LATREILLE, 1802): [28]
 = *Myrmica rubida* (LATREILLE, 1802): [42]
 = *Myrmica* (*Neomyrma*) *rubida*: [27]

Genus: *Myrmica*

M. constricta KARAVAJEV, 1934: based on findings of Seifert et al. [50], *M. constricta* is an eastern European congener of *M. hellenica* that has long been considered to occur in Hungary. The formerly believed continuous population of “*M. hellenica*” is therefore split into two geographically distinct species and the Hungarian population belongs to *M. constricta*.

= *Myrmica hellenica* FINZI, 1926: [51]
M. curvithorax BONDROIT, 1920
 = *Myrmica slovaca* SADIL, 1952: [24]
 = *Myrmica salina* Ruzsky, 1905: [38]
M. deplanata Ruzsky, 1905: [27]
M. gallienii BONDROIT, 1920: [38]
M. karavajevi (ARNOLDI, 1930)
 = *Sifolinia karavajevi*: [28]
 = *Sifolinia faniensis* (VAN BOVEN, 1970): [52]
M. lobicornis NYLANDER, 1846: [26]
M. lonae FINZI, 1926: [53]
M. rubra (LINNAEUS, 1758): [38]
 = *Myrmica laevinodis* NYLANDER, 1846: [25]
 = *Myrmica microrubra* SEIFERT, 1993: [54]
M. ruginodis NYLANDER, 1846: [42]

- M. rugulosa* NYLANDER, 1849: [26]
M. sabuleti MEINERT, 1861: [27]
M. scabrinodis NYLANDER, 1846: [42]
= *Myrmica rugulosoides* FOREL, 1915: [55]
M. schencki VIERECK, 1903: [56]
M. specioides BONDROIT, 1918: [57]
= *Myrmica sancta* KARAVAIEV, 1926: [58]
M. vandeli BONDROIT, 1920: [53]

3.3.4. Tribe: Solenopsidini

Genus: *Monomorium*

- M. pharaonis* (LINNAEUS, 1758): [27] (Indoor tramp species)

Genus: *Solenopsis*

- S. fugax* (LATREILLE, 1798): [25]

3.3.5. Tribe: Stenammini

Genus: *Aphaenogaster*

- A. subterranea* (LATREILLE, 1798): [26]

Genus: *Messor*

- M. structor* (LATREILLE, 1798): [27]
= *Atta structor*: [42]
= *Aphaenogaster structor*: [26]
= *Messor rufitarsus* (FABRICIUS 1804): [59]

Genus: *Stenamma*

- S. debile* (FÖRSTER, 1850): [60]
= *Stenamma westwoodii* WESTWOOD, 1839: [27]

3.4. Subfamily: Dolichoderinae

3.4.1. Tribe: Bothriomyrmecini

Genus: *Bothriomyrmex*

- B. communista* SANTSCHI, 1919
= *Bothriomyrmex meridionalis* (ROGER, 1863): [27]
B. corsicus SANTSCHI, 1923
= *Bothriomyrmex menozzii* EMERY, 1925: [61]

3.4.2. Tribe: Dolichoderini

Genus: *Dolichoderus*

- D. quadripunctatus* (LINNAEUS, 1771): [26]
= *Hypoclinea quadripunctata*: [42]

3.4.3. Tribe: Tapinomini

Genus: *Liometopum*

- L. microcephalum* (PANZER, 1798): [26]
= *Formica austriaca* MAYR, 1853: [42]

Genus: *Tapinoma*

- T. erraticum* (LATREILLE, 1798): [42]
T. melanocephalum (FABRICIUS, 1793): [24] (Indoor tramp species)
T. subboreale SEIFERT, 2012: [62]
= *Tapinoma madeirense* FOREL, 1895: [24]
= *Tapinoma ambiguum* EMERY, 1925: [56]

3.5. Subfamily: Formicinae

3.5.1. Tribe: Camponotini

Genus: *Camponotus*

C. aethiops (LATREILLE, 1798):

= *Formica aethiops* LATREILLE, 1798: [42]

= *Camponotus marginatus* LATREILLE, 1798: [26]

C. atricolor (NYLANDER, 1849) (sensu SEIFERT, 1996): [63]

C. fallax (NYLANDER, 1856): [64]

= *Camponotus caryae* var. *fallax* (NYLANDER, 1856): [27]

= *Camponotus caryae* (FITCH, 1855): [65]

C. herculeanus (LINNAEUS, 1758): [26]

C. lateralis (OLIVIER, 1792): [26]

= *Formica lateralis* OLIVIER, 1792: [42]

C. ligniperda (LATREILLE, 1802): [26]

C. piceus (LEACH, 1825): [52]

= *Camponotus lateralis* var. *piceus* (LEACH, 1825): [27]

C. tergestinus MÜLLER, 1921: [66]

C. vagus (SCOPOLI, 1763): [26]

= *Formica pubescens* FABRICIUS, 1775: [25]

= *Formica ligniperda*: [42]

Genus: *Colobopsis*

C. truncata (SPINOLA, 1808): [26]

3.5.2. Tribe: Formicini

Genus: *Cataglyphis*

C. aenescens (NYLANDER, 1849): [28]

= *Formica cursor* FONSCOLOMBE, 1846: [25]

= *Myrmecocystus cursor*: [26]

= *Cataglyphis cursor aenescens*: [67]

C. nodus (BRULLÉ, 1832): [28]

= *Monocombus viaticus* (FABRICIUS, 1787): [42]

= *Formica viatica*: [25]

= *Cataglyphis viaticus* var. *orientalis* FOREL, 1895: [26]

= *Myrmecocystus bicolor* (FABRICIUS, 1793): [27]

= *Cataglyphis bicolor nodus* (BRULLÉ, 1832): [52]

Genus: *Formica*

F. cinerea MAYR, 1853: [27]

F. clara Forel, 1886

= *Formica glauca* RUZSKY, 1895: [51]

= *Formica lusatica* (SEIFERT 1997): [68]

F. cunicularia LATREILLE, 1798: [25]

= *Formica fusca glebaria* NYLANDER, 1846: [67]

F. exsecta NYLANDER, 1846: [25]

F. fusca LINNAEUS, 1758: [25]

F. fuscocinerea FOREL, 1874: [54]

F. gagates LATREILLE, 1798: [42]

F. lemani BONDROIT, 1917: new record, Veszprém m. Szigliget, 20-22.04.2019, leg. Z. Vas (3 workers)

Formica lemani might be hard to distinguish from its congeners, *F. fusca* and *F. gagates*, hence the identification made by SC was confirmed via multivariate analyses of numeric traits using a modern numeric morphology-based alpha taxonomic key provided by Seifert [23]. The pubescence on the first gastral tergite (sqPDG) is very dense, scores

vary between 3.46 and 3.87, which is in the lowest range of the *F. gagates* (3.6 to 10.8) and the highest scores of *F. lemani* and *F. fusca* (2.4 to 3.5), but the dull cuticular surface of the workers and highly dense pubescence rule out the possibility of *F. gagates*. The relatively high number of unilateral pronotal setae (4 to 15, nest sample mean 7.67) and the discriminant function provided by Seifert [23] place the nest sample in *F. lemani* with a very high certainty. The D5 scores vary between 6.03 and 8.62 (nest sample mean 7.15). Morphometric character recording and analyses were done by SC.

- F. polyctena* FÖRSTER, 1850: [52]
F. pratensis RETZIUS, 1783: [26]
= *Formica congerens* NYLANDER, 1846: [25]
= *Formica rufa pratensis*: [40]
F. pressilabris NYLANDER, 1846: [43]
F. rufa LINNAEUS, 1761: [26]
F. rufibarbis FABRICIUS, 1793: [26]
= *Formica cunicularia* LATREILLE, 1798: [26]
= *Formica fusca* var. *glebaria* NYLANDER, 1846: [27]
= *Formica fusca glebaria* NYLANDER, 1846: [67]
F. sanguinea LATREILLE, 1798: [25]
F. truncorum FABRICIUS, 1804: [27]
= *Formica truncicola* NYLANDER, 1846: [25]

Genus: *Polyergus*

- P. rufescens* (LATREILLE, 1798): [25]

3.5.3. Tribe: Lasiini

Genus: *Lasius*

- L. alienus* (FÖRSTER, 1850): [26]
= *Formica aliena* FÖRSTER, 1850: [42]
L. balcanicus SEIFERT, 1988: [38]
L. bicornis (FÖRSTER, 1850): [27]
L. bombycinus (SEIFERT and GALKOWSKI, 2016) based on current findings of Seifert and Galkowski [30] *Lasius bombycinus* is an eastern European congener of *L. paralienus* that has long been considered to occur in Hungary. The formerly believed continuous population of “*L. paralienus*” is therefore split into two geographically distinct species and the Hungarian population belongs to *L. bombycinus*.
= *Lasius paralienus* SEIFERT, 1992: [38]
L. brunneus (LATREILLE, 1798): [26]
L. carniolicus MAYR, 1861: [41]
L. citrinus EMERY, 1922: [69]
= *Lasius bicornis* var. *affinis* SCHENCK, 1852: [70]
= *Lasius affinis* SCHENCK, 1852: [27]
L. distinguendus (EMERY, 1916): [54]
L. emarginatus (OLIVIER, 1792): [26]
L. flavus (FABRICIUS, 1782): [26]
= *Formica flava* FABRICIUS, 1782: [25]
L. fuliginosus (LATREILLE, 1798): [26]
= *Formica fuliginosa* LATREILLE, 1798: [42]
L. jensi SEIFERT, 1982: [54]
L. meridionalis (BONDROIT, 1920): [38]
L. mixtus (NYLANDER, 1846): [26]
L. myops FOREL, 1894: [28]
= *Lasius flavus* var. *myops* FOREL, 1894: [27]
L. neglectus Van Loon, Boomsma et Andrásfalvy, 1990: [71]
L. niger (LINNAEUS, 1758): [26]

- = *Formica nigra* LINNAEUS, 1758: [42]
L. nitidigaster SEIFERT, 1996: [29]
L. platythorax SEIFERT, 1991: [38]
= *Formica brunnea* LATREILLE, 1798: [25]
= *Formica timida* FÖRSTER, 1950: [25]
L. psammophilus SEIFERT, 1992: [38]
L. umbratus (NYLANDER, 1846): [26]
= *Formica umbrata* NYLANDER, 1846: [25]

Genus: *Prenolepis*

- P. nitens* (MAYR, 1853): [28]
= *Tapinoma nitens*: [25]
= *Prenolepis imparis* var. *nitens*: [26]

3.5.4. Tribe: Plagiolepidini

Genus: *Plagiolepis*

- P. ampeloni* (FABER, 1969): [24]
P. pallescens FOREL, 1889: [24]
P. pygmaea (LATREILLE, 1798): [26]
= *Tapinoma pygmaeum*: [42]
P. taurica SANTSCHI, 1920: [62]
= *Plagiolepis vindobonensis* LOMNICKI, 1925: [27]
P. xene STAERCKE, 1936: [41]

3.6. Species of Doubtful Occurrence

3.6.1. *Temnothorax rabaudi* (BONDROIT, 1918)

This species was reported from Hungary by Barrett [72] based on a male individual (in combination with *Leptothorax*), but it must have been a misidentification. The identification of *Temnothorax* species based only on males is possible in exceptional cases, hence such determinations should be considered uncertain. Moreover, *T. rabaudi* is a West European species, which may not stretch its distribution to the Carpathian Basin. Since neither confirmed information is available, nor has the voucher specimen been deposited to any public collection, this taxon is eliminated from the Hungarian checklist.

3.6.2. *Tetramorium impurum* (FÖRSTER, 1850)

This species was identified by Csősz et al. [54] based on workers and sexual forms, but the samples definitely belong to *Tetramorium staerkei* KRATOCHVÍL 1944. *Tetramorium impurum* is a mountain species and is unlikely to occur in the Carpathian Basin. Therefore this taxon is eliminated from the current checklist.

4. Discussion

Checklists are more than just simple lists of taxa. Inventorying what we have forms the solid basis of any nature conservation policy, but this information also serves as an indispensable tool for scientific field experiments. Ant taxonomy is a highly dynamic field, where new taxa are described or old names are revised even in the otherwise seemingly well-known European fauna every year [32,73], thus there is a need for a periodical update of any checklist, as is the case of the Hungarian myrmecofauna.

The Hungarian fauna and flora are characterized by a mixture of Eastern European, Central European and Mediterranean elements; many species having here their westernmost, easternmost or even northernmost location of their range of distribution. Additionally, there are quite a number of endemic animal and plant species, characteristic to the Pannonian biogeographical region, which harbors higher biodiversity compared to surrounding biogeographical regions, being mostly restricted to Hungary [74–76]. While in ants there are no endemic species or even subspecies known to Hungary, the Hungarian

myrmecofauna is no exception to the above-mentioned characteristics. Submediterranean elements as *Cataglyphis*, *Cardiocondyla* and *Bothriomyrmex* species are present, along with typical European temperate climate species, as *Myrmica* and *Formica* s. str. species.

Since the publication of the last Hungarian checklist [24] several taxonomic novelties appeared starting from the thorough revision of the genus *Lasius* [30,77] to the clarification of the taxonomy of the extremely problematic genus *Tetramorium* [31,48], which is also very well reflected by the high species richness of both genera in the Hungarian myrmecofauna. Characteristic to the Hungarian myrmecofauna is the presence of some truly rare species, as *Lasius carniolicus* or parasitic *Plagiolepis* species. Moreover, there is quite a considerable number of indoor species as well confirming the fact that constant efforts have been put into the survey of the myrmecofauna in the past decades.

In order to reveal the characteristics of any fauna, it should be put in the context of data known from neighboring countries (Figure 1). The biogeographic specificity of Hungary is also revealed by the high number of species compared to other countries in temperate Europe relative to the country's size and geographical characteristics (Figure 1). Surprisingly, the number of Hungarian species is comparable to the myrmecofauna of the much more diverse Slovenia or Ukraine, while it surpasses the known number of species in Germany or Romania, which are larger countries with habitats ranging from high mountains to seashores. Certainly, the smaller number of known ant species in other countries could also be attributed to the relative lack of data as is, probably, the case of Romania (see [13] for arguments). However, such comparisons, and the fact that in the last decade only six new species were recorded, entitle us to say that the Hungarian myrmecofauna is quite well-known; many more new native species are not expected to occur, maybe with the exception of some parasitic ones with cryptic lifestyle. While the number of known species may not vary much in the forthcoming years, knowledge on the geographic distribution of Hungarian ant species is still incomplete. Thus, the implementation of further faunistic studies should be encouraged to complete the picture with the missing biogeographic data.




Figure 1. The number of known ant species in other European countries (Turkey: [17]; France 1: [20]; France 2: [19]; Bulgaria: [78]; Montenegro: [21]; Switzerland: [23]; Austria: [23]; Serbia: [79]; Ukraine: [14]; Slovenia: [80]; Germany: [23]; Romania: [13,16,32,81]; Czech Republic: [82]; Slovakia: [83]; Poland: [15]; Republic of North Macedonia: [84]; Andorra: [18]; Belgium: [85]).

5. Conclusions

We provide a detailed list of the Hungarian ant fauna by adding the new elements found in the last decade including the annotated synonymic list of taxa.

Inventorying what we have forms the solid basis of any nature conservation policy. Such lists can also help to cope with recent challenges posed by agricultural intensification and climate change generated biodiversity crisis and serve indispensable information for scientific field experiments.

Awareness of fauna composition is a must to assess changes in our environment. The Hungarian fauna and flora are characterized by a mixture of Eastern European, Central European and Mediterranean elements; many species having here their westernmost, easternmost or even northernmost location of their range of distribution. Thereby our fauna can be considered one of the indicator areas that are affected by climatic changes.

In order to better highlight the characteristics of the Hungarian ant fauna, it is put in the context of data known from neighboring countries (Figure 1). The biogeographic specificity of Hungary is also revealed by the relatively high number of species compared to other countries in temperate Europe (Figure 1). Surprisingly, the number of Hungarian species is comparable to the myrmecofauna of the much more diverse Slovenia or Ukraine, while it surpasses the known number of species in Germany or Romania, which are larger countries with habitats ranging from high mountains to seashores.

These comparisons make us say that the Hungarian myrmecofauna is quite well-known, and many more new native species are not expected to occur. However, some cryptic parasitic ones or invasive elements can be expected to show up in the near future.

Author Contributions: Conceptualization, S.C. and B.M.; methodology, S.C. and B.M.; validation, S.C., F.B., A.T., A.Á.S., and I.M.; formal analysis, B.M.; investigation, S.C., F.B., L.G., G.L., I.M., A.T., É.K., A.Á.S., and B.M.; resources, S.C., F.B., L.G., G.L., I.M., A.T., É.K., A.Á.S., and B.M.; data curation, S.C. and F.B.; writing—original draft preparation, S.C., B.M., F.B., A.T., A.Á.S., L.G., G.L., and I.M.; writing—review and editing, S.C., F.B., A.T., B.M., and A.Á.S.; visualization, B.M., A.T.; supervision, S.C.; project administration, S.C., B.M., and A.T.; funding acquisition, S.C., B.M., and A.T. All authors have read and agreed to the published version of the manuscript.

Funding: S.C. was supported by the National Research, Development, and Innovation Fund under Grant No. K 135795. A.T. was supported by a ‘János Bolyai’ Scholarship of the Hungarian Academy of Sciences, by the EFOP-3.6.1-16-2016-00022 project (co-financed by the European Union and the European Social Fund) and by the ÚNKP-20-5 New National Excellence Program of the Ministry of Human Capacities.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Acknowledgments: We are indebted to all the myrmecologists who have made essential contributions to our knowledge of Hungarian myrmecofauna over the course of the past decades, either by collecting new species for the fauna, such as was done by Csaba Nagy, or by offering valuable new records concerning species already identified, as done by Hajnalka Dürögő, Orsolya Kanizsai, Klára Benedek, Orsolya Szalárdy, and many others. Ádám Bakos provided his photo for the Graphical Abstract.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Causey, D.; Janzen, D.H.; Peterson, A.T.; Vieglais, D.; Krishtalka, L.; Beach, J.H.; Wiley, E.O. Museum collections and taxonomy. *Science* **2004**, *305*, 1106. [[CrossRef](#)] [[PubMed](#)]
- Wheeler, Q.D.; Raven, P.H.; Wilson, E.O. Taxonomy: Impediment or Expedient. *Science* **2004**, *303*, 285. [[CrossRef](#)] [[PubMed](#)]
- Carvalho, M.R.; Bockmann, F.A.; Amorim, D.S.; de Vivo, M.; de Toledo-Piza, M.; Menezes, N.A.; de Figueiredo, J.L.; Castro, R.M.C.; Gill, A.C.; McEachran, J.D.; et al. Revisiting the taxonomic impediment. *Science* **2005**, *307*, 353. [[CrossRef](#)] [[PubMed](#)]

4. Hallmann, C.A.; Sorg, M.; Jongejans, E.; Siepel, H.; Hofland, N.; Schwan, H.; Stenmans, W.; Müller, A.; Sumser, H.; Hörren, T.; et al. More than 75 percent decline over 27 years in total flying insect biomass in protected area. *PLoS ONE* **2017**, *12*, e0185809. [[CrossRef](#)]
5. Rhodes, C.J. Are insect species imperilled? Critical factors and prevailing evidence for a potential global loss of the entomofauna: A current commentary. *Sci. Prog.* **2017**, *102*, 181–196. [[CrossRef](#)]
6. Suprayitno, N.; Narakusumo, R.P.; von Rintelen, T.; Hendrich, L.; Balke, M. Taxonomy and Biogeography without frontiers—WhatsApp, Facebook and smartphone digital photography let citizen scientists in more remote localities step out of the dark. *Biodivers. Data J.* **2017**, *5*. [[CrossRef](#)]
7. Devictor, V.; Mouillot, D.; Meynard, C.; Jiguet, F.; Thuiller, W.; Mouquet, N. Spatial mismatch and congruence between taxonomic, phylogenetic and functional diversity: The need for integrative conservation strategies in a changing world. *Ecol. Lett.* **2010**, *13*, 1030–1040. [[CrossRef](#)]
8. McKinley, D.; Miller-Rushing, A.J.; Ballard, H.L.; Bonney, R.; Brown, H.; Cook-Patton, S.C.; Evans, D.M.; French, R.A.; Parrish, J.K.; Phillips, T.B.; et al. Citizen science can improve conservation science, natural resource management, and environmental protection. *Biol. Conserv.* **2017**, *208*, 15–28. [[CrossRef](#)]
9. Freitag, H.; Pangantihon, C.V.; Njunjić, I. Three new species of *Grouvellinus* Champion, 1923 from Maliau Basin, Sabah, Borneo, discovered by citizen scientists during the first Taxon Expedition (*Insecta, Coleoptera, Elmidae*). *ZooKeys* **2018**, *754*, 1–24. [[CrossRef](#)]
10. Moulin, N. When Citizen Science highlights alien invasive species in France: The case of Indochina mantis, *Hierodula patellifera* (*Insecta, Mantodea, Mantidae*). *Biodivers. Data J.* **2020**, *8*. [[CrossRef](#)]
11. Tanduo, V.; Golemaj, A.; Crocetta, F. Citizen-science detects the arrival and establishment of *Branchiomma luctuosum* (Grube, 1870) (*Annelida: Polychaeta: Sabellidae*) in Albania. *Biodivers. Data J.* **2020**, *8*. [[CrossRef](#)]
12. Vinarski, M.V. Roots of the taxonomic impediment: Is the “integrativeness” a remedy? *Integr. Zool.* **2020**, *15*, 2–15. [[CrossRef](#)] [[PubMed](#)]
13. Markó, B.; Sipos, B.; Csősz, S.; Kiss, K.; Boros, I.; Gallé, L. A comprehensive list of the ants of Romania (Hymenoptera: Formicidae). *Myrmecol. Nachr.* **2006**, *9*, 65–76.
14. Radchenko, A.G. Zonal and zoogeographic characteristic of the ant fauna (Hymenoptera, Formicidae) of Ukraine. *Vestn. Zool.* **2011**, *45*, 30–39. [[CrossRef](#)]
15. Czechowski, W.; Radchenko, A.; Czechowska, W.; Vepsäläinen, K. *The Ants of Poland with Reference to the Myrmecofauna of Europe-Fauna Poloniae Vol. 4 New Series*; Museum and Institute of Zoology of the Polish Academy of Sciences and Natura optima dux Foundation: Warszawa, Poland, 2012.
16. Czeker, Z.; Radchenko, A.G.; Csősz, S.; Szász-Len, A.-M.; Tăușan, I.; Benedek, K.; Markó, B. The genus *Myrmica* Latreille, 1804 (Hymenoptera: Formicidae) in Romania: Distribution of species and key for their identification. *Entomol. Romanica* **2012**, *17*, 29–50.
17. Kiran, K.; Karaman, C. First annotated checklist of the ant fauna of Turkey (Hymenoptera: Formicidae). *Zootaxa* **2012**, *3548*, 1–38. [[CrossRef](#)]
18. Bernadou, A.; Fourcassié, V.; Espadaler, X. A preliminary checklist of the ants (Hymenoptera, Formicidae) of Andorra. *ZooKeys* **2013**, *277*, 13–23. [[CrossRef](#)]
19. Blatrix, R.; Galkowski, C.; Lebas, C.; Wegnez, P. *Fournmis de France, de Belgique et du Luxembourg [The Ants of France, Belgium, and Luxembourg]*; Delachaux & Niestlé: Paris, France, 2013; ISBN 978-2-603-01899-6.
20. Monnin, T.; Espadaler, X.; Lenoir, A.; Peeters, C. *Guide des Fourmis de France*; Belin: Paris, France, 2013; ISBN 978-2-7011-6471-7.
21. Bračko, G.; Gomboc, M.; Lupša, B.; Marić, R.; Prstovšek, U. New faunistic data on ants (Hymenoptera: Formicidae) of the southern part of Montenegro. *Nat. Slov.* **2014**, *16*, 41–51.
22. Wagner, H.C. *Die Ameisen Kärntens: Verbreitung, Biologie, Ökologie und Gefährdung [The Ants of Carinthia: Distribution, Biology, Ecology and Threats]*; Naturwissenschaftlicher Verein für Kärnten: Klagenfurt, Austria, 2014; ISBN 9783853280669.
23. Seifert, B. *The Ants of Central and North Europe*; Lutra Verlags- und Vertriebsgesellschaft: Tauer, Germany, 2018; ISBN 9783936412079.
24. Csősz, S.; Markó, B.; Gallé, L. The myrmecofauna (Hymenoptera: Formicidae) of Hungary: An updated checklist. *North West. J. Zool.* **2011**, *7*, 55–62.
25. Mayr, G. Ungarn’s Ameisen. *Program. der Städtischen Oberrealschule Pesth* **1857**, *3*, 5–26.
26. Mocsáry, A. Ordo Hymenoptera. In *Fauna Regni Hungariae*; Mocsáry, A., Ed.; Királyi Magyar Természettudományi Társulat: Budapest, Hungary, 1897; pp. 7–116.
27. Somfai, E. Hangya alkatúák, Formicoidea. [Ants, Formicoidea]. In *Magyarország Állatvilága (Fauna Hungariae)*; Székessy, V., Ed.; Akadémiai Kiadó: Budapest, Hungary, 1959; Volume 13, pp. 1–79.
28. Gallé, L.; Csősz, S.; Tartally, A.; Kovács, É. A check-list of Hungarian ants (Hymenoptera: Formicidae). *Folia Entomol. Hung.* **1998**, *59*, 213–220.
29. Lőrinczi, G. *Lasius nitidigaster* Seifert, 1996—a new ant species (Hymenoptera: Formicidae) for the Hungarian fauna. *Nat. Som.* **2011**, *19*, 223–226.
30. Seifert, B.; Galkowski, C. The Westpalaearctic *Lasius paralienus* complex (Hymenoptera: Formicidae) contains three species. *Zootaxa* **2016**, *4132*, 44–58. [[CrossRef](#)] [[PubMed](#)]

31. Wagner, H.C.; Arthofer, W.; Seifert, B.; Muster, C.; Steiner, F.M.; Schlick-Steiner, B.C. Light at the end of the tunnel: Integrative taxonomy delimits cryptic species in the *Tetramorium caespitum* complex (Hymenoptera: Formicidae). *Myrmecol. News* **2017**, *25*, 95–129. [[CrossRef](#)]
32. Steiner, F.M.; Csósz, S.; Markó, B.; Gamisch, A.; Rinnhofer, L.; Folterbauer, C.; Hammerle, S.; Stauffer, C.; Arthofer, W.; Schlick-Steiner, B.C. Turning one into five: Integrative taxonomy uncovers complex evolution of cryptic species in the harvester ant *Messor "structor"*. *Mol. Phylogenet. Evol.* **2018**, *127*, 387–404. [[CrossRef](#)]
33. Wagner, H.C.; Seifert, B.; Aurenhammer, S.; Komposch, C. *Temnothorax turcicus* (Santschi, 1934)—eine arborikole Ameise (Hymenoptera: Formicidae) neu für Österreich. *Ber. Nat. Med. Ver. Innsbr.* **2011**, *97*, 59–71.
34. Bolton, B. AntCat: An Online Catalog of Ants of the World. California Academy of Sciences. Available online: <http://antcat.org/> (accessed on 21 May 2020).
35. Csósz, S. A key to the Ponerinae species of the Carpathian Basin (Hymenoptera: Formicidae). *Ann. Hist. Nat. Mus. Nat. Hung.* **2003**, *95*, 147–160.
36. Csósz, S.; Seifert, B. *Ponera testacea* Emery, 1895 stat n.—a sister species of *P. coarctata* (Latreille, 1802) (Hymenoptera, Formicidae). *Acta Zool. Acad. Sci. Hung.* **2003**, *49*, 201–214.
37. Mayr, G.L. *Die Europäischen Formiciden: Nach der Analytischen Methode Bearbeitet von Gustav L. Mayr*; C. Gerold's Sohn: Wien, Austria, 1861.
38. Csósz, S.; Tártally, A. Adatok a Körös-Maros Nemzeti Park hangyafaunájához [Data to the ant fauna of the Körös-Maros National Park]. *Crisicum* **1998**, *1*, 180–194.
39. Csósz, S. Taxonomic and distributional notes on two new and a rare *Leptothorax* Mayr, 1855 species for the Hungarian ant fauna (Hymenoptera, Formicidae). *Ann. Hist. Nat. Mus. Nat. Hung.* **2001**, *93*, 99–106.
40. Móczár, L. Bátoraliget hártányásszárnyú faunája, Hymenoptera [Hymenoptera fauna of Bátoraliget]. In *Bátoraliget Élővilága [Flora and Fauna of Bátoraliget]*; Székessy, Ed.; Akadémiai Kiadó: Budapest, Hungary, 1953; pp. 286–316.
41. Gallé, L.; Szönyi, G. The checklist of ants (Hymenoptera, Formicoida) of a sandy grassland in Kiskunság National Park (Hungary). *Acta Biol. Szeged.* **1988**, *34*, 167–168.
42. Mayr, G. Formicina austriaca. Beschreibung der bisher im österreichischen Kaiserstaate aufgefundenen Ameisen nebst Hinzufügung jener in Deutschland, in der Schweiz und in Italien vorkommenden Ameisen. *Verh. Des. Zool. Bot. Ver. Wien.* **1855**, *5*, 273–478.
43. Radchenko, A.G. The ant fauna of Őrség, western Hungary (Hymenoptera: Formicidae). In *Natural History of Őrség Landscape Conservation Area III*; Vig, K., Ed.; Savaria—a Vas Megyei Múzeumok Értesítője: Szombathely, Hungary, 1997; Volume 24, pp. 81–91.
44. Röszler, P. Beiträge zur Kenntnis der Ameisenfauna von Siebenbürgen und Ungarn. *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* **1935**, *83–84*, 72–83.
45. Röszler, P. Myrmecologisches aus dem Jahre 1938. *Zool. Anz.* **1951**, *146*, 88–96.
46. Csósz, S.; Markó, B. Redescription of *Tetramorium hungaricum* Röszler, 1935, a related species of *T. caespitum* (Linnaeus, 1758) (Hymenoptera: Formicidae). *Myrmecol. Nachr.* **2004**, *6*, 49–59.
47. Schlick-Steiner, B.C.; Steiner, F.M.; Moder, K.; Seifert, B.; Sanetra, M.; Dyreson, E.; Stauffer, C.; Christian, E. A multidisciplinary approach reveals cryptic diversity in Western Palearctic *Tetramorium* ants (Hymenoptera: Formicidae). *Mol. Phylogenet. Evol.* **2006**, *40*, 259–273. [[CrossRef](#)]
48. Csósz, S.; Wagner, H.C.; Bozsó, M.; Seifert, B.; Arthofer, W.; Schlick-Steiner, B.C.; Steiner, F.M.; Pénzes, Z. *Tetramorium indocile* Santschi, 1927 stat. rev. is the proposed scientific name for *Tetramorium* sp. C sensu based on combined molecular and morphological evidence (Hymenoptera: Formicidae). *Zool. Anz.* **2014**, *253*, 469–481. [[CrossRef](#)]
49. Csósz, S.; Radchenko, A.G.; Schulz, A. Taxonomic revision of the Palaearctic *Tetramorium chefketi* species complex (Hymenoptera: Formicidae). *Zootaxa* **2007**, *1405*, 1–38. [[CrossRef](#)]
50. Seifert, B.; Schlick-Steiner, B.C.; Steiner, F.M. *Myrmica constricta* Karavaiev, 1934—a cryptic sister species of *Myrmica hellenica* Finzi, 1926 (Hymenoptera: Formicidae). *Soil Org.* **2009**, *81*, 53–76.
51. Gallé, L.; Markó, B.; Kiss, K.; Kovács, É.; Dürögő, H.; Kőváry, K.; Csósz, S. Ant fauna of Tisza river basin (Hymenoptera: Formicidae). *Tisia Monogr. Ser.* **2005**, *7*, 149–197.
52. Gallé, L. Adatok a Bakony hegység hangyafaunájának ismeretéhez (Hymenoptera, Formicidae) [Contributions to the knowledge of the myrmecofauna (Hymenoptera, Formicidae) of the Bakony mountain]. *Veszprém M. Múzeumok Közleményei* **1979**, *14*, 239–244.
53. Tártally, A.; Csósz, S. Adatok a *Maculinea* boglárkalepkék (Lepidoptera: Lycaenidae) kárpát-medencei hangyagazdáiról [Data on the ant hosts of the *Maculinea* butterflies (Lepidoptera: Lycaenidae) of Hungary]. *Természetvédelmi Közlemények* **2004**, *11*, 309–317.
54. Csósz, S.; Markó, B.; Kiss, K.; Tártally, A.; Gallé, L. The ant fauna of the Fertő-Hanság National Park (Hymenoptera: Formicoidea). In *The Fauna of the Fertő-Hanság National Park*; Mahunka, S., Ed.; Hungarian Natural History Museum: Budapest, Hungary, 2002; pp. 617–629.
55. Gallé, L. Ecological and zoocoenological conditions of the Formicoidea fauna at Tiszakürt. *Tisia* **1967**, *3*, 67–72.
56. Gallé, L. The ant fauna of the Kiskunság National Park. In *The Fauna of Kiskunság National Park*; Mahunka, S., Ed.; Akadémiai Kiadó: Budapest, Hungary, 1986; pp. 427–434.
57. Gallé, L. Factors stabilizing the ant populations (Hymenoptera: Formicidae) in the grass associations of the Tisza Basin. *Tisia* **1975**, *10*, 61–66.
58. Gallé, L. Formicidae populations of the ecosystems in the environs of Tiszafüred. *Tisia* **1972**, *7*, 59–68.

59. Gallé, L. Habitat and niche analysis of grassland ants. *Entomol. Gen.* **1986**, *11*, 197–211. [[CrossRef](#)]
60. Csősz, S. Hangya-faunisztikai adatok a Körös-Maros Nemzeti Parkból: A mályvádi erdők [Ant-faunistic investigations in the Körös-Maros National Park: The Mályvád-Forests]. *Crisicum* **2000**, *3*, 183–187.
61. Markó, B.; Csősz, S. Die europäischen ameisenarten (Hymenoptera: Formicidae) des Herrmannstädter (Sibiu, Rumänien) Naturkundemuseums I.: Unterfamilien Ponerinae, Myrmicinae und Dolichoderinae. *Ann. Hist. Nat. Mus. Nat. Hung.* **2002**, *94*, 109–121.
62. Gallé, L.; Kanizsai, O.; Maák, I.; Lőrinczi, G. Close nesting association of two ant species in artificial shelters: Results from a long-term experiment. *Acta Zool. Acad. Sci. Hung.* **2014**, *60*, 359–370.
63. Seifert, B. *Ameisen, Beobachten, Bestimmen*; Naturbuch Verlag: Augsburg, Germany, 1996; ISBN 978-3894401702.
64. Gallé, L. The Formicoid fauna of the Hortobágy. In *The Fauna of the Hortobágy National Park*; Mahunka, S., Ed.; Akadémiai Kiadó: Budapest, Hungary, 1981; pp. 307–311.
65. Gallé, L. Ecological and zoocoenological investigation of the Formicoidea fauna of the flood area of the River Tisza. *Tiscia* **1966**, *2*, 113–118.
66. Ionescu-Hirsch, A.; Markó, B.; Csősz, S. *Camponotus tergestinus* Müller, 1921 (Hymenoptera: Formicidae): First record of a rare species for Romania and Hungary. *Entomol. Romanica* **2009**, *14*, 19–22.
67. Gallé, L. Study of ant populations in various grassland ecosystems. *Acta Biol. Szeged.* **1972**, *18*, 159–164.
68. Tartally, A.; Csősz, S. Blaskovics-puszta gyepek értékelése a rajtuk talált hangyaközösségek (Hymenoptera: Formicidae) alapján [The evaluation of the Blaskovics-puszta's grasslands and their ant (Hymenoptera: Formicidae) communities]. *Crisicum* **1999**, *2*, 133–140.
69. Csősz, S.; Markó, B. European Ant Species (Hymenoptera: Formicidae) in the ant collection of the natural history museum of Sibiu (Hermannstadt/Nagyszeben), Romania II. Subfamily Formicinae. *Ann. Hist. Nat. Mus. Nat. Hung.* **2005**, *97*, 225–240.
70. Mocsáry, A. Ordo Hymenoptera. In *Fauna Regni Hungariae. Arthropoda.*; Paszlawsky, J., Ed.; Királyi Magyar Természettudományi Társulat: Budapest, Hungary, 1918.
71. Van Loon, A.J.; Boomsma, A.; Andrásfalvy, A. A new polygynous *Lasius* species from Central Europe. *Insectes Soc.* **1990**, *37*, 348–362. [[CrossRef](#)]
72. Barrett, K.E. Ants from Hungary and Bulgaria. *Entomologist* **1970**, *103*, 139–140.
73. Csősz, S.; Salata, S.; Borowiec, L. Three Turano-European species of the *Temnothorax interruptus* group (Hymenoptera: Formicidae) demonstrated by quantitative morphology. *Myrmecol. News* **2018**, *26*, 101–119. [[CrossRef](#)]
74. Varga, Z. Geographical patterns of biological diversity in the Palaearctic Region and the Carpathian Basin. *Acta Zool. Acad. Sci. Hung.* **1995**, *41*, 71–92.
75. Borhidi, A. *Magyarország Növénytársulásai* [Plant. Communities of Hungary]; Akadémiai Kiadó: Budapest, Hungary, 2003; ISBN 963-05-7983-9.
76. Varga, Z. Biogeografía: Az élet földrajza [Biogeography: Geography of life]. In *Biogeografía: Az élet földrajza PARS Könyvek* (12); Peregovits, L., Ed.; PARS Kft: Nagykovácsi, Hungary, 2019; ISBN 978-963-88339-9-0.
77. Seifert, B. A taxonomic revision of the Palaearctic members of the subgenus *Lasius* s. str (Hymenoptera, Formicidae). *Soil Org.* **2020**, *92*, 15–86.
78. Borowiec, L.; Lapeva-Gjonova, A.; Salata, S. Three species of *Aphaenogaster* Mayr, 1853 (Hymenoptera: Formicidae) new to Bulgarian fauna. *Acta Zool. Bulg.* **2019**, *71*, 613–616.
79. Petrov, I.Z. *Mravi Srbije i Crne Gore* [The ants of Serbia and Montenegro]; SANU [Serbian Academy of Sciences and Arts]: Beograd, Serbia, 2006; ISBN 978-8670254084.
80. Bračko, G. Checklist of the ants of Slovenia (Hymenoptera: Formicidae). *Nat. Slov.* **2007**, *9*, 15–24.
81. Tăușan, I.; Lapeva-Gjonova, A. *Camponotus samius* Forel, 1889 (Hymenoptera: Formicidae)-at the north edge of its European distribution. *North. West. J. Zool.* **2017**, *13*, 352–354.
82. Werner, P.; Bezděčka, P.; Bezděčková, K.; Pech, P. An updated checklist of the ants (Hymenoptera: Formicidae) of the Czech Republic. *Acta Rerum Nat.* **2018**, *22*, 5–12.
83. Werner, P.; Wieszik, M. Vespoidea: Formicidae (mravencovití). *Acta Entomol. Mus. Nationalis Pragae* **2007**, *11*, 133–164.
84. Bračko, G.; Wagner, H.C.; Schulz, A.; Gioahin, E.; Matičić, J.; Tratnik, A. New investigation and a revised checklist of the ants (Hymenoptera: Formicidae) of the Republic of Macedonia. *North. West. J. Zool.* **2014**, *10*, 10–24.
85. Dekoninck, W.; Vankerkhoven, F. Checklist of the Belgian ant-fauna (Formicidae, Hymenoptera). *Bull. Inst. R. Sci. Natur. Belg. Entomol.* **2001**, *71*, 263–266.