

Table 1. Olfactory, clinical and epidemiological outcome correlations.

Variable	P-value
Gender	0.177
Age	0.362
Clinical severity	0.038
Diabetes	0.146
Hypertension	0.849
Gastroesophageal disorders	0.003
Asthma	0.855
Heart disorders	0.275
Smoker	0.258
Fever	0.011
Cough	0.223
Chest pain	0.002
Anorexia	0.159
Sticky mucus/phlegm	0.035
Arthralgia	0.219
Myalgia	0.230
Diarrhea	0.104
Abdominal pain	0.054
Nausea, vomiting	0.065
Headache	0.708
Asthenia, malaise or confusion	0.227
Dyspnea	0.027
Nasal Obstruction	0.184
Rhinorrhea	0.279
Postnasal drip	0.287
Sore throat	0.203

Ear pain	0.961
Face pain	0.355
Dysphagia	0.276
Dysphonia	0.014

Table S2: Cross-tab analysis results

Variable	Z	OR	95% Confidence interval	P-value
<u>Clinical severity</u>				0.038
Hyposmia	0.297	0.931	0.582 - 1.490	0.766
Anosmia	2.425	0.484	0.269 - 0.87	0.015
Anosmia or hyposmia	1.467	0.726	0.474 - 1.114	0.142
<u>Gastroesophageal disorders</u>				0.003
Hyposmia	2.426	1.86	1.127 - 3.072	0.015
Anosmia	3.345	2.425	1.443 - 4.975	< 0.001
Anosmia or hyposmia	2.228	1.67	1.064 - 2.623	0.003
<u>Fever</u>				0.011
Hyposmia	2.614	1.61	1.126 - 2.302	0.009
Anosmia	0.083	0.983	0.659 - 1.467	0.934
Anosmia or hyposmia	1.651	1.31	0.951 - 1.806	0.1
<u>Chest pain</u>				0.002
Hyposmia	2.141	1.516	1.036 - 2.22	0.032
Anosmia	1.453	0.705	0.441 - 1.129	0.934
Anosmia or hyposmia	0.752	1.144	0.806 - 1.625	0.452
<u>Sticky mucus/phlegm</u>				0.035

Hyposmia	2.535	1.701	1.128 - 2.566	0.011
Anosmia	0.865	1.229	0.77 - 1.96	0.387
Anosmia or hyposmia	2.099	1.496	1.027 - 2.178	0.036
<u>Dyspnea</u>				0.027
Hyposmia	0.114	1.02	0.720 - 1.446	0.909
Anosmia	2.373	0.609	0.404 - 0.917	0.018
Anosmia or hyposmia	1.134	0.834	0.609 - 1.141	0.036
<u>Dysphonia</u>				0.014
Hyposmia	0.990	1.215	0.826 - 1.789	0.322
Anosmia	2.058	0.603	0.373 - 0.976	0.034
Anosmia or hyposmia	0.348	0.939	0.659 - 1.338	0.728

Table S2 footnotes: Abbreviations: OR: odds ratio