

Table S6. Morphometric data of *Trichuris suis* by different authors, presented as mean \pm standard deviation, in millimeters

	This study (n=10)	Cutillas et al. [25]		Nissen et al. [26]
Host	Wild boar	Pig	Wild boar	Pig
Male				
Length	36.784 \pm 3.854	40.4 \pm 4.2	40.8 \pm 8.8	28.9
Esophagus	2.346 \pm 0.29	2.61 \pm 0.31	2.5 \pm 0.58	-
Width of the anterior end	0.184 \pm 0.07	0.20 \pm 0.02	0.20 \pm 0.03	0.16
Width at the esophagus-intestine junction	0.379 \pm 0.08	0.35 \pm 0.04	0.38 \pm 0.06	-
Width of the posterior end	0.725 \pm 0.098	0.87 \pm 0.06	0.83 \pm 0.13	0.58
Spicule	2.37 \pm 0.182	2.35 \pm 0.15	2.43 \pm 0.22	2.21
Spicule sheath	0.29 \pm 0.132	0.34 \pm 0.20	0.15 \pm 0.09	-
Female				
Length	44.77 \pm 3.8	44.3 \pm 2.8	46.3 \pm 5.3	28.5
Esophagus	3.268 \pm 0.201	3.12 \pm 0.16	3.28 \pm 0.39	-
Width of the anterior end	0.21 \pm 0.04	0.19 \pm 0.01	0.20 \pm 0.02	0.18
Width at the esophagus-intestine junction	0.34 \pm 0.08	0.36 \pm 0.03	0.30 \pm 0.08	-
Width of the posterior end	0.73 \pm 0.11	0.93 \pm 0.10	0.90 \pm 0.08	0.69
Vulva to the esophagus-intestine junction	0.25 \pm 0.05	0.23 \pm 0.08	0.25 \pm 0.022	-