

Supplementary Table S1: Statin use and in-hospital outcomes.

	No Statin Use	Statin Use	p-value
In-Hospital outcomes			
Need for ICU	43 (34.1%)	50 (40.7%)	0.29
Need for IMV	26 (20.6%)	19 (15.4%)	0.29
Death	15 (11.9%)	27 (22.0%)	0.034
ICU duration (days*)	7.0 (3.0-14.0)	5.5 (4.0-9.0)	0.34
IMV duration (days*)	5.5 (2.0-13.0)	9.0 (5.0-14.0)	0.099

* Reported as median (IQR)

Supplementary Table S2: ACEi/ARB use and in-hospital outcomes*

	No ACEi/ARB Use	ACEi/ARB Use	p-value
In-Hospital outcomes			
Need for ICU	47 (33.8%)	46 (41.8%)	0.19
Need for IMV	22 (15.8%)	23 (20.9%)	0.30
Death	16 (11.5%)	26 (23.6%)	0.011
ICU duration (days†)	6.00 (3.0-12.0)	6.0 (4.0-16.0)	0.67
IMV duration (days†)	6.00 (2.0-10.0)	7.0 (5.0-14.0)	0.17

* ACEi refers to angiotensin converting enzyme inhibitor. ARB refers to angiotensin II receptor blocker; † Reported as median (IQR)