

Arsenic Removal via the Biomineralization of Iron-Oxidizing Bacteria *Pseudarthrobacter* sp. Fe7

Xia Fan ¹, Hanxiao Zhang ¹, Qian Peng¹, Yongliang Zheng ¹, Kaixiang Shi ^{2*} and Xian Xia ^{3*}

¹ College of Biology and Agriculture Resources, Huanggang Normal University, Huanggang 438000, China; fanxia@hgnu.edu.cn (X.F.); 15038881601@163.com (H.Z.); m18040651471@163.com (Q.P.); zylgolden@126.com (Y.Z.)

² National Key Laboratory of Agricultural Microbiology, College of Life Science and Technology, Huazhong Agricultural University, Wuhan 430070, China

³ Hubei Key Laboratory of Edible Wild Plants Conservation & Utilization, Hubei Engineering Research Center of Characteristic Wild Vegetable Breeding and Comprehensive Utilization Technology, College of Life Science, Hubei Normal University, Huangshi 43500, China

* Correspondence: kaixiangshi@mail.hzau.edu.cn (K.S.); xianxia@hbnu.edu.cn (X.X.)




Figure S1. (A) Colony morphology and (B) SEM image of strain Fe7.




Figure S2. Growth status of strain Fe7 in a modified PYCM liquid medium and SEM image of the iron precipitate produced by strain Fe7. (A) Medium without strain Fe7. (B) Cell suspension. (C) Biological iron precipitates. (D) SEM pattern.