

Supporting Information

An engineered reporter phage for the fluorometric detection of *Escherichia coli* in ground beef

Anqi Chen,^a Danhui Wang,^{a, b} Sam R. Nugen^{*a} and Juhong Chen^{a,c}

^aDepartment of Food Science, Cornell University, Ithaca, NY 14853, United States

^bDepartment of Food Science and Technology, The Ohio State University, Columbus, OH 43210, United States

^cDepartment of Biological Systems Engineering, Virginia Tech, Blacksburg, Virginia 24061, United States

Corresponding author:

* (SRN). Email: snugen@cornell.edu




Figure S1 Fluorescence intensity measured from T7_{lacZ} phages infecting *E. coli* with different concentrations ($0, 10^1, 10^2, 10^3, 10^4$, and 10^5 CFU/mL) in buffer solution after (a) 1 hour (b) 2 hours (c) 3 hours (d) 4 hours (e) 5 hours (f) 6 hours (g) 7 hours (h) 8 hours of incubation at 37°C, respectively.




Figure S2 Fluorescence intensity measured from T7_{lacZ} phages infecting *E. coli* with different concentrations (0, 10, 10², 10³ CFU/mL) in ground beef after (a) 5 hour (b) 6 hours (c) 7 hours (d) 8 hours of incubation at 37°C, respectively.