

Table S2. Demographic and clinical characteristics of the patients infected by B.1.1.7 and non-B.1.1.7 lineages according to age group (≤ 65 and > 65 year). IQR: interquartile range, OR: odds ratio, 95%CI: 95% confidence interval, ICU: intensive care unit, IRCU: intensive respiratory care unit, sig: significance.

Variable	Patients ≤ 65 Years				Patients > 65 Years			
	B.1.1.7/SGTF (<i>n</i> = 191)	B.1.1.7 (<i>n</i> = 453)	<i>p</i>	OR	B.1.1.7/SGTF (<i>n</i> = 235)	B.1.1.7 (<i>n</i> = 676)	<i>p</i>	OR
Age (median)	54	54	0.62		78	82	0.001	
IQR	(45–60)	(47–60)			(71–86)	(74–88)		
Sex								
Women	36.1%	36.9%	0.86		45.1%	49.3%	0.27	
Men	63.9%	63.1%			54.9%	50.7%		
ICU admission	23.6%	18.1%	0.11	1.39 (0.92–2.1)	16.2%	5.0%	0.001	3.64 (2.23–5.94)
(95%CI)	(17.73–30.23)	(14.66–21.96)			(11.7–21.51)	(3.5–6.96)		
IRCU admission	4.2%	1.8%	0.07	2.43 (0.89–6.58)	12.3%	6.2%	0.03	2.12 (1.29–3.49)
(95%CI)	(1.82–8.09)	(0.76–3.45)			(8.42–17.24)	(4.51–8.3)		
Death	4.2%	3.5%	0.68	1.21 (0.48–3.02)	22.5%	23.6%	0.57	0.82 (0.64–1.38)
(95%CI)	(1.69–8.39)	(1.96–5.66)			(16.79–29.1)	(20.37–27.03)		