

Supplementary Material S5. Distribution of COVID-19 cases with regard to the WHO Clinical Progression Scale

Table S5. COVID-19 severity and symptoms of the cases in the efficacy analysis with regard to WHO Clinical Progression Scale (the highest score during the course of the disease is given here)

| Patient state | Descriptor | CoronaVac | TURKOVAC | Total |
|--------------------------------|---|--------------|--------------|-----------|
| | | arm n (%) | arm n (%) | n (%) |
| Ambulatory mild disease | 2 Symptomatic- independent | 60 (13.1) | 35 (7.7) | 95 (10.4) |
| | 3 Symptomatic- assistance needed | 0 (0) | 0 (0) | 0 (0) |
| | 4 Hospitalized (If hospitalization is for isolation only record the status with an outpatient)- no oxygen therapy | 0 (0) | 0 (0) | 0 (0) |
| Hospitalized: moderate disease | 5 Hospitalized; oxygen by mask or nasal prongs | 1 (0.2) | 0 (0) | 1 (0.1) |
| Hospitalized: severe diseases | 6 Hospitalized-oxygen by non-invasive or high flow ventilation | 0 (0) | 0 (0) | 0 (0) |

| | | | |
|-------|-----------|----------|-----------|
| Total | 61 (13.3) | 35 (7.7) | 96 (10.5) |
|-------|-----------|----------|-----------|