

Supplementary Table S1. Baseline clinical characteristics of the excluded cohorts

	Excluded after PSM n = 548	iSAVR < 75 years n = 532	TAVR < 75 years n = 16
Demographics			
Age, mean-(±SD)	68.8 (5.6)	62.8 (10.0)	69.6 (4.2)
Female, n-(%)	215 (39.2)	204 (38.3)	11 (68.8)
Body mass index kg/m ² , median-(IQR)	28.8 (5.8)	28.6 (5.5)	29.5 (5.3)
Risk profile			
EuroSCORE II, median (IQR)	4.6 (3.5)	1.5 (1.8)	13.4 (6.7)
Chronic health conditions and risk factors			
Hypertension, n (%)	423 (77.2)	411 (77.3)	12 (75.0)
Dyslipidaemia, n (%)	350 (63.9)	338 (63.5)	12 (75.0)
Diabetes mellitus, n (%)	43 (7.8)	39 (7.3)	4 (25.0)
Active smoker, n (%)	171 (31.2)	165 (31.0)	6 (37.5)
Serum creatinine mg/dL, mean (±SD)	1.3 (0.8)	1.0 (0.3)	2.4 (2.2)
Preoperative dialysis, n (%)	3 (0.5)	0 (0)	3 (18.8)
Chronic obstructive pulmonary disease, n (%)	139 (25.3)	127 (22.8)	12 (75.0)
Peripheral vascular disease, n (%)	39 (7.1)	29 (5.5)	10 (62.5)
Cerebrovascular disease, n (%)	83 (15.2)	78 (14.7)	5 (31.3)
Previous cerebrovascular accident, n (%)	4 (0.7)	3 (0.6)	1 (6.3)
Atrial fibrillation, n (%)	92 (16.8)	87 (16.4)	5 (31.3)
Previous myocardial infarction, n (%)	31 (5.7)	27 (5.1)	4 (25.0)
New York Heart Association class III/IV, n (%)	253 (46.2)	237 (44.6)	16 (100)
Previous PCI, n (%)	25 (4.7)	22 (4.1)	3 (18.8)

Previous pacemaker implantation, n (%)	16 (2.9)	12 (2.3)	4 (25.0)
Previous cardiac surgery, n (%)	22 (4.0)	10 (1.9)	12 (75.0)
Previous CABG, n (%)	8 (1.6)	2 (0.4)	6 (37.5)
Previous valve surgery, n (%)	14 (2.6)	6 (1.3)	8 (50.0)
Previous other cardiac surgery, n (%)	7 (58.3)	31 (0.4)	5 (31.3)
Preoperative echocardiographic data			
Mean pressure gradient, mean (\pm SD)	47.8 (19.0)	49.2 (24.3)	43.7 (17.1)
Left ventricular ejection fraction %, mean (\pm IQR)	45.8 (12.1)	53.8 (8.6)	41.1 (13.1)

CABG – Coronary Artery Bypass Graft; COPD – chronic obstructive pulmonary disease; EuroSCORE - European System for Cardiac Operative Risk Evaluation; IDDM – Insulin dependent diabetes mellitus; IQR – Interquartile range; LVEF - left ventricular ejection fraction; NYHA – New York Heart Association; PSM – propensity score matching; PCI - percutaneous coronary intervention; SD – standard deviation;

Supplementary Table S2. Procedural characteristics of the excluded cohorts

	Excluded after PSM n = 548	iSAVR < 75 years n = 532	TAVR < 75 years n = 16
Procedural variables			
Biological valve prosthesis, n (%)	461 (84.1)	445 (83.6)	16 (100)
Balloon-expandable THV, n (%)			10 (62.5)
Prosthesis size in mm, mean (\pm SD)	26.0 (2.5)	22.8 (3.2)	25.4 (2.6)
Full sternotomy, n (%)		428 (80.5)	
Cross-clamp time, mean (\pm SD)		67.3 (23.4)	
Perfusion time, mean (\pm SD)		99.2 (42.9)	
Transfemoral access, n (%)			13 (81.3)
Predilatation, n (%)			5 (31.3)
Postdilatation, n (%)			4 (25)
Paravalvular leak > mild, n (%)	0 (0)	0 (0)	0 (0)
Postoperative circulatory support, n (%)	4 (0.7)	3 (0.6)	1 (6.3)
Extubated in the OR, n (%)	4 (0.7)	0 (0)	4 (25)
Total hours ventilated, median (\pm IQR)	5 (9)	8 (8)	4 (8)
Re-intubated during hospital stay, n (%)	15 (2.7)	15 (2.8)	0 (0)
Number of red blood cell units, mean (\pm SD)	0.9 (2.6)	1.0 (3.0)	0 (0)
Length of stay, median (\pm IQR)	10 (8)	11 (5)	10 (13)

OR - operating room; THV - transcatheter heart valves; other abbreviations as in Supplementary table 1

Supplementary Table S3. Adverse events of the excluded cohorts

	Excluded after PSM n = 548	iSAVR < 75 years n = 532	TAVR < 75 years n = 16
Procedural variables			
Myocardial infarction, n (%)	1 (0.2)	1 (0.2)	0 (0)
Neurological adverse event, n (%)	4 (0.7)	4 (0.8)	0 (0)
Major vascular complication, n (%)	1 (0.2)	0 (0)	1 (6.3)
Major bleeding complication, n (%)	19 (3.5)	18 (3.4)	1 (6.3)
Postoperative renal replacement therapy, n (%)	6 (1.1)	6 (1.1)	0 (0)
New-onset atrial fibrillation, n (%)	65 (11.9)	65 (12.2)	0 (0)
AV-Block III, n (%)	16 (2.9)	13 (2.4)	3 (18.8)
Pacemaker implantation, n (%)	16 (2.9)	13 (2.4)	3 (18.8)
Reoperation for valvular dysfunction, n (%)	1 (0.2)	1 (0.2)	0 (0)
Reoperation for bleeding/tamponade, n (%)	11 (2.0)	11 (2.1)	0 (0)
Reoperation for other cardiac problem, n (%)	2 (0.4)	1 (0.2)	1 (6.3)
Reoperation for non-cardiac problem, n (%)	10 (1.8)	10 (1.9)	0 (0)
Postoperative sepsis, n (%)	2 (0.4)	2 (0.4)	0 (0)
Deep sternal wound infection, n (%)	5 (0.9)	5 (0.9)	0 (0)
Prolonged ventilation >6hrs, n (%)	44 (8.0)	43 (8.1)	1 (6.3)
Multi-system organ failure, n (%)	5 (0.9)	5 (0.9)	0 (0)
In-hospital death, n (%)	6 (1.1)	6 (1.3)	0 (0)
30-day all-cause mortality, n (%)	8 (1.6)	7 (1.3)	1 (6.3)

AV – atrioventricular; other abbreviations as in Supplementary table 1, 2 and 3