

Article

Exploring the Mechanisms of the Spatiotemporal Invasion of *Tuta absoluta* in Asia

Ritter A. Guimapi ^{1,2,*} , Ramasamy Srinivasan ³ , Henri E. Tonnang ¹ ,
Paola Sotelo-Cardona ³  and Samira A. Mohamed ¹

¹ ICIPE—International Centre of Insect Physiology and Ecology, Nairobi 30772-00100, Kenya; htonnang@gmail.com (H.E.T.); sfaris@icipe.org (S.A.M.)

² Department of computing, School of Computing & Information Technology, Jomo Kenyatta University of Agriculture and Technology (JKUAT), Nairobi 62000-00200, Kenya

³ World Vegetable Center, Shanhua, Tainan 74151, Taiwan; srini.ramasamy@worldveg.org (R.S.); paola.sotelo@worldveg.org (P.S.-C.)

* Correspondence: ritteryvan@gmail.com

Received: 8 March 2020; Accepted: 8 April 2020; Published: 13 April 2020



Abstract: International crop exchange always brings the risk of introducing pests to countries where they are not yet present. The invasive pest *Tuta absoluta* (Meyrick 1917), after taking just a decade (2008–2017) to invade the entire Africa continent, is now continuing its expansion in Asia. From its first detection in Turkey (2009), the pest has extended its range of invasion at a very high speed of progression to the southeast part of Asia. This study adopted the cellular automata modelling method used to successfully predict the spatiotemporal invasion of *T. absoluta* in Africa to find out if the invasive pest is propagating with a similar pattern of spread in Asia. Using land cover vegetation, temperature, relative humidity and the natural flight ability of *Tuta absoluta*, we simulated the spread pattern considering Turkey as the initial point in Asia. The model revealed that it would take about 20 years for the pest to reach the southeast part of Asia, unlike real life where it took just about 10 years (2009–2018). This can be explained by international crop trade, especially in tomatoes, and movement of people, suggesting that recommendations and advice from the previous invasion in Europe and Africa were not implemented or not seriously taken into account. Moreover, some countries like Taiwan and the Philippines with suitable environmental condition for the establishment of *T. absoluta* are not at risk of natural invasion by flight, but quarantine measure must be put in place to avoid invasion by crop transportation or people movement. The results can assist policy makers to better understand the different mechanisms of invasion of *T. absoluta* in Asia, and therefore adjust or adapt control measures that fit well with the dynamic of the invasive pest observed.

Keywords: cellular automata; *Tuta absoluta*; insect's pest invasion; dispersal pattern; international crop trade; integrated pest management

1. Introduction

Tuta absoluta (Meyrick 1917) (*Lepidoptera: Gelechiidae*), the South American tomato pinworm, is considered a serious lepidopteran pest that causes severe damage to tomatoes, leading to economic losses with a shortage of tomato supply in affected countries [1–4]. After its initial arrival from South America to Spain in 2006, other European countries including Italy (2008), France (2008), Albania (2009), Bulgaria (2009), Portugal (2009), the Netherlands (2009), United Kingdom (2009), and Serbia (2011) were also affected [5,6]. Later, *T. absoluta* also invaded Africa through Morocco (2007/2008), Algeria (2008), Tunisia, Libya, (2009), Senegal (2011), Sudan (2011), Ethiopia (2013), Kenya (2013), and South Africa in 2017 [7–13], as well as Benin in 2018 [14]. Turkey was the first country in Asia to detect the presence of *T. absoluta* in 2009 [15], and from there

it has spread and reached many countries in South and Central Asia including India (2015), the northwestern Himalayan region of India, Bangladesh (2016), Nepal (2016), Kyrgyzstan (2017), Tajikistan (2018) [16–22], and possibly in Myanmar (2018) in both greenhouse and open field conditions. However, despite the wide spread of *T. absoluta* in South Asia, the invasive pest has not yet been officially reported in many countries in Southeast or East Asia, including Laos, Thailand, Cambodia, Vietnam, Malaysia, Indonesia, Philippines, Taiwan, Korea and Japan.

While spreading from one location to another, *T. absoluta* causes crops damage, with losses which can reach up to 100% in tomato production [10,23]. However, the complex physiology and ecology of *T. absoluta* may be in part responsible for the lack of successful management strategies [10], in spite of multiple activities and efforts undertaken both in laboratory and in the field during the last decade to control this invasive pest [24–35]. *Tuta absoluta* has a high reproduction rate, with about 12 generations per year in optimal conditions, which range between 21 to 30 °C, and higher than 50% relative humidity. In addition, a mature *T. absoluta* female can lay up to 260 eggs with very high flight ability, reaching up to 100 km during its lifetime [36–38].

According to [39], species with a high growth rate, maturation, and reproduction on a wide range of environmental conditions are those with a great probability of invasion [39]. These attributes perfectly fit with the physiological and ecological description of *T. absoluta*. A combination of these factors with the constant interaction of *T. absoluta* with human activities such as crop transportation favors the expansion of its range to new regions where the invasive pest was not present before. The consequence of these interactions is that it becomes difficult and complex to implement efficient control measures against *T. absoluta* due to multiple paths of invasion. There are three main paths of insect invasion: it can occur naturally, by introduction, or accidentally, and is caused either by human activities or by natural dispersal [40]. Previous work has shown that from its first detection in Morocco in 2006, *T. absoluta* has invaded the entire African continent in ten years naturally, just by its flying ability [7]. However, there is not yet a report of any study exploring the mechanism of invasion of *T. absoluta* in Southeast Asia through modeling and simulation.

Many modelling concepts exist to simulate insect population dynamics and risk of invasion, including the use of the Species distribution modelling (SDM) approach, differential equations, individual-based models and cellular automata [41,42]. The success of one approach depends on selection of an appropriate complexity level for the modelling experiment objectives and the availability of data [42]. SDM is often used for predicting the suitable area for occurrence and the distribution of species without emphasis on the dispersal pattern across space and time. These modelling approaches cannot inform about the speed of spread and when the event will occur in a given location. Differential equations (DE) are frequently used to address the issues related to temporal predictions of insect population dynamics and abundance [43,44]. The inconvenience of using the DE approach is that they presume the space to be uniform and continuous, which is generally not the case in reality, especially for large scale areas of study like countries and continents. Indeed, models that attempt to integrate all existing knowledge about an insect species such as DE, and to serve all possible purposes, usually become obsolete before completion and have unexpectedly limited usage because of their lack of flexibility [45]. On the other hand, models with spatiotemporal approaches using cellular automata (CA) have been useful to understand the path of invasion and predict the timing of invasion of *T. absoluta* in Africa [7] and Asia [46]. The CA-based approach is spatially explicit and favors the consideration of the spatial heterogeneity of interactions in the dynamic process. The aim of this work is to use the model previously designed by [7] in order to predict the timing of invasion of *T. absoluta* in Southeast Asia, compare the path and mechanism of the invasion of the pest in Asia, and advise on suitable management strategies to be adopted.

2. Materials and Methods

2.1. Area of Study and Data Transformation

To process the simulations, temperature, relative humidity, and *T. absoluta* occurrence data from Asia were used (Figure 1). Previous works have highlighted the key role played by temperature and RH for the survival of *T. absoluta* both in the laboratory and in the field [37,47–49]. The vegetation variable, characterized by the normalized difference vegetation index (NDVI), was not taken into account in this study because it was demonstrated that NDVI did not really influence the choice of a location of *T. absoluta* during the process of invasion, unlike temperature and relative humidity [7]. Moreover, NDVI does not negatively affect the flying distance of *T. absoluta* during the invasion process. This is the same with the parameter quantity of tomato (main host plant) production per hectare, which mainly improves detection of potential invaded locations where environmental conditions are not suitable, without necessarily slowing down the flying distance [7]. Given that the main interest here was to explore if the path of invasion in Asia is the same as the one in Africa, only the climatic parameters were used. The temperature dataset was obtained from the WorldClim database (<http://www.worldclim.org/current>) (Hijmans et al., 2005); relative humidity data were downloaded on the Surface meteorology and Solar Energy (SSE) website (<http://eosweb.larc.nasa.gov/sse/>). However, *T. absoluta* occurrence data in Asia were collected from different scientific publications [16–21].




Figure 1. Area of study. The location in green represents the Asian continent; the red dot is the Asia location in Turkey where *T. absoluta* was discovered for the first time in 2009. This point is also the initial location for the model simulation.

The temperature datasets were organized into monthly mean values using a grid of 30 arc-seconds that correspond to about 1-km resolution. These processes were carried out using the functionalities offered by the R-packages *sp* (1.3.1) and *raster* (2.6.7). These allow reading, writing and manipulating of

the gridded spatial data. We further applied the *rgdal* (1.3.4) functionalities for bindings. The Geospatial Data Abstraction Library (GDAL) was useful to handle and convert different geospatial data in different formats [50–52]; the function *extractValue()* was implemented using the R programming language to extract temperature values of all geographic coordinates overlapping the area of study for a temperature raster file. Relative humidity data were obtained and organized into text file format containing coordinates of the Earth's surface spaced by 1×1 degree. The inverse distance weighting method (IDW) [53] functionality built in the Geographic Information System (GIS) software Quantum GIS (QGIS) was used to interpolate value and produce a raster file of relative humidity data. Hence, the method *extractValue()* was used again following the same process as described for the extraction of the temperature value.

2.2. Cellular Automata (CA) Model Thresholds, Rules and Implementation

The model used in the current study is based on the CA model designed by [7] to predict the invasion of *T. absoluta* in Africa, due to the accuracy provided to spatially mimic the timing of invasion of *T. absoluta* in Africa. cellular automata have been very useful to explore the dynamics of many agricultural systems and processes such as the prediction of land use and land cover change, or the vegetation and landscape dynamics [54–56]. CA modeling is characterized by a set of states and a set of rules that describe the dynamics of the studied system after every time step. It can be represented by a square lattice where each cell represents a location. Therefore, the state of a cell at any given time t depends on its state at the time $t-1$ and the transition rule applied from t to $t-1$. In the context of this study, a location can be found in two possible states, which are susceptible (S) and invaded (I). All the locations of our area of study at the beginning of the simulation are in the state susceptible, except the location in Turkey from which the pest invaded Asia. Therefore, that particular location was used in the state invaded (I).

For the CA model, a monthly time step was used for simulations. A temperature threshold was defined at 22 °C, while RH threshold was at 55%. As the natural ability of flight of *T. absoluta* is taken into account, 100 km was used, considering the Moore neighborhood [57]. From the transition time t to $t + 1$, cells in the neighborhood of an invaded area are updated based on the following rules:

Susceptible cell (S): As previously explained, all the locations within the area of study and at the beginning of the simulation are Susceptible to be invaded, except the first invaded location in Asia (i.e., Turkey). The location will remain susceptible (S to S) if the combination of environmental condition does not favor the settlement of the invasive pest. However, a susceptible location turns into invaded (S to I) if both temperature and relative humidity value in the location are greater than their respective threshold. Once it becomes invaded, the state of the location cannot change anymore and will remain invaded (I to I) as no control measures are taken into account during the simulation.

MATLAB (R2010a, The Math-Works) was used as a programming medium to implement the model. Therefore, simulations were made starting from Turkey, and were processed until the entire Asian continent was covered; from there, the number of years was recorded, and a comparison was made with the progression under real conditions.


2.3. Keys Assumptions

The developed model is based on the following assumptions: (1) the time scale of *T. absoluta* invasion process is driven by its natural flight ability like in Africa, without human assisted transportation; (2) for all locations where the environmental conditions are favorable for agricultural activities, there is sufficient host plant vegetation for sustaining the spread of this invasive pest. However, despite the presence of three main tomato producers of the world in Asia, there are many areas unsuitable for tomato cropping.

3. Results

The simulation of the invasion process of *T. absoluta* in Asia, considering Turkey (2009) as the entry location in Asia, is presented in the following maps (Figure 1), showing the progression of the invasion when temperature, relative humidity, and the natural flight ability of *T. absoluta* were taken into account.

From the start of invasion, and based on the simulation model, *T. absoluta* would require 25 years for natural dispersal into Southeast Asia.






Figure 1. The invasion of *Tuta absoluta* in Asia. The simulation is made for 25 years considering the temperature and relative humidity as parameters. Locations in green are those susceptible to being invaded; locations in red are those invaded by the spread of *T. absoluta*. (a) 1 Year; (b) 4 Years; (c) 6 years; (d) 8 years; (e) 10 years; (f) 12 years; (g) 15 years; (h) 17 years; (i) 19 years; (j) 21 years; (k) 23 years; (l) 25 years.

Comparison between the simulations and current progress showed a contrasting story, since the current invasive pest progress is much faster than the one predicted from the simulation model. More specifically, it took about ten years for *T. absoluta* to reach Myanmar (no official report yet, but the invasive pest presence has been suspected). These observations, therefore, reject the hypothesis that the invasion of *T. absoluta* in Asia is only due to its natural flight ability, as previously found under African conditions. Moreover, isolated regions or countries like the Philippines, Taiwan, parts of Indonesia and Malaysia seem to show a low risk of natural invasion, due to the large distance between continental Asia and these island regions, restricting natural flight arrival. In addition, and based on the simulation model, the northern part of Asia would be also out of risk of natural invasion mainly because of the environmental conditions, which seem to be unsuitable for the establishment and development of the invasive pest.

4. Discussion

The development and implementation of efficient control measures against insect invasive pests with complex biological and physiological characteristic such as *T. absoluta* require a good understanding of the mechanism used by the insect pest to disperse and establish. Here we used the CA model developed by [7], who successfully predicted the spatiotemporal invasion of *T. absoluta* in Africa to explore if the mechanism and path of dispersal being used by the invasive pest in Asia are the same. The details about the phenology of *T. absoluta* are not taken into account by the model. However, without minimizing the level of damage caused by the larvae of *T. absoluta* in crops [58], the simulations have provided a broad idea about when the invasive pest is supposed to reach certain location(s) in order to anticipate and adopt proper management strategies in place to slow down the arrival of the pest.

Regardless of the simulation model, an important observation is that *T. absoluta* has spread in Asia faster than initially supposed considering only its natural flight ability. Using the simulation model, *T. absoluta* was expected to invade Southeast Asia naturally in about 20 years after its first detection in Turkey. However, in the current real-time situation, the invasive pest reached this region less than 10 years from its first report in Turkey. International crop trade and movement of people have proven to be responsible for expediting the invasion of insect pests [40,59–61]. These factors can easily explain the high speed of invasion of *T. absoluta* in Asia, especially among the top five highest tomato producing regions in the world, including three Asian countries: China, India, and Turkey. Although it has not yet been reported from China, its presence in Myanmar, Laos or Vietnam can expedite the invasion into southern China. Likewise, it is worrying that a large quantity of tomato produced in these affected regions/countries is exported to other countries that have not yet reported the presence of the invasive pest. Therefore, it is expected and predictable that a rapid introduction and invasion of this pest to other neighboring countries in the short term is possible.

Many Asian countries are among the world top five producers of potatoes and eggplants, with about 40% of the world production for potatoes and more than 75% of world production for eggplants [62]. This includes China and India for potato production and China, India, Turkey, and Iran for eggplant [63]; moreover, these crops are also widely cultivated on the entire continent by other countries (FAO: <http://www.fao.org/faostat/en/#data/QC/visualize>). These factors make the Asian continent extremely vulnerable and more suitable for the invasion of *T. absoluta*. Considering that a huge number of these products are traded worldwide assisted by human transportation, this will speed up the spread of *T. absoluta* towards surrounding locations. Although damages due to *T. absoluta* on crops in Asia have been reported on all previous mentioned crops [64], recent studies reported that this invasive pest poses a major threat to tomatoes and less to potato, eggplant, and other solanaceous species [18,65]. With tomatoes as the main target and the preferred host plant of *T. absoluta*, it would be advisable to focus the development and implementation of control strategies on tomatoes, and also the improvement of regulation of human mediated international crop trade to slow down and limit the spread.

Apart from the contrast observed with the speed of invasion of *T. absoluta* in Asia, the simulations also showed that the northern part of Asia seems not to be at risk of natural invasion by *T. absoluta*, due to unfavorable climatic conditions for the establishment of this invasive pest. These results are in agreement with the previous suitability map produced for the same pest in Asia [18,66]. Moreover, most of the isolated countries in Southeast or East Asia such as Taiwan, the Philippines, Korea or Japan are also out of risk from natural invasion by flight, despite suitable environmental conditions for the establishment of *T. absoluta* [18,66]. However, the invasive pest can be easily introduced if these countries import tomatoes from those countries which have already reported the presence of *T. absoluta*. Therefore, proper quarantine measures are highly recommended in these countries to curtail or slow down the invasion by crop transportation or human movement.

According to [18], many countries in Asia, including both invaded and non-invaded ones, do not seem to take the economic and ecological damage due to *T. absoluta* seriously, because the invasive pest has not been included in their quarantine list [18]. Unfortunately, local farmers tend to adopt the use of chemical pesticides once the pest is detected, but it has already been proven to be inefficient against *T. absoluta* [67–71]. It is quite usual that farmers change and adopt IPM strategies including pheromone trapping, the use of natural enemies and microbial insecticides only after realizing the control failures even after applying these chemical pesticides [72–75]. Phytosanitary officers can play a key role in preventing or slowing down the progress of *T. absoluta* in Asia, especially by (i) advising government(s) of *T. absoluta*-free countries but with suitable environmental conditions for the invasive pest to adopt quarantine measures, and avoid or limit crop importation from already invaded countries; (ii) educating and training farmers in *T. absoluta*-invaded regions in the successful existing IPM strategies to control or reduce damage from *T. absoluta* intrusion; (iii) creating a platform for technical information exchange, where farmers, researchers, phytosanitary officers and governmental staff from different countries will interact for timely and rapid communication of advances in *T. absoluta* management.

5. Conclusions

Herein we investigate the process of the spatiotemporal invasion of *T. absoluta* in Asia using a CA model to predict its natural spread pattern. We realized that the speed of spread of *T. absoluta* in Asia is accelerated by international crop exchange and human movement across the continent. Therefore, including information about the trade of the main host plant (tomatoes) could help extend this study and help policy makers to adapt control measures that fit well with the observed dynamic of the pest.

Author Contributions: R.A.G., R.S., S.A.M. conceived and designed research. R.A.G. implemented and did the simulations. R.A.G., H.E.T., P.S.-C., wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Federal Ministry for Economic Cooperation and Development (GIZ), grant number 16.7860.6-001.00.

Acknowledgments: This work was supported by the Federal Ministry for Economic Cooperation and Development (GIZ Project number 16.7860.6-001.00), Germany and by core donors to the World Vegetable Center: Republic of China (Taiwan), UK aid from the UK Government, United States Agency for International Development (USAID), Australian Centre for International Agricultural Research (ACIAR), Germany, Thailand, Philippines, Korea, and Japan. We also thank Sopana Yule, WorldVeg East and Southeast Asia, Thailand for her support in data collection, and the German Academic Exchange Service (DAAD) and the STRIVE project funded by the German Federal Ministry of Education and Research.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Zekeya, N.; Chacha, M.; Ndakidemi, P.A.; Materu, C.; Chidege, M.; Mbega, E.R. Tomato Leafminer (*Tuta absoluta* Meyrick 1917): A Threat to Tomato Production in Africa. *J. Agric. Ecol. Res. Int.* **2017**, *10*, 1–10. [[CrossRef](#)]
2. Tadele, S.; Emanu, G. Determination of the economic threshold level of tomato leaf miner, *Tuta absoluta* Meyrick (*Lepidoptera: Gelechiidae*) on tomato plant under glasshouse conditions. *J. Hortic. For.* **2018**, *10*, 9–16. [[CrossRef](#)]
3. Gharekhani, G.; Salek-Ebrahimi, H. Evaluating the damage of *Tuta absoluta* (Meyrick) (*Lepidoptera: Gelechiidae*) on some cultivars of tomato under greenhouse condition. *Arch. Phytopathol. Plant Prot.* **2013**, *47*, 429–436. [[CrossRef](#)]
4. Negeri, T.S.A.M.; Getu, E. Experimental Analysis of Economic Action Level of Tomato Leafminer, *Tuta absoluta* Meyrick (*Lepidoptera: Gelechiidae*) on Tomato Plant under Open Field. *Adv. Crop Sci. Technol.* **2018**, *6*, 1–5. [[CrossRef](#)]
5. Desneux, N.; Luna, M.G.; Guillemaud, T.; Urbaneja, A. The invasive South American tomato pinworm, *Tuta absoluta*, continues to spread in Afro-Eurasia and beyond: The new threat to tomato world production. *J. Pest Sci.* **2011**, *84*, 403–408. [[CrossRef](#)]
6. Campos, M.R.; Biondi, A.; Adiga, A.; Guedes, R.N.C.; Desneux, N. From the Western Palaearctic region to beyond: *Tuta absoluta* 10 years after invading Europe. *J. Pest Sci.* **2017**, *90*, 787–796. [[CrossRef](#)]
7. Guimapi, R.Y.; Mohamed, S.A.; Okeyo, G.O.; Ndjomatchoua, F.; Ekesi, S.; Tonnang, H.E. Modeling the risk of invasion and spread of *Tuta absoluta* in Africa. *Ecol. Complex.* **2016**, *28*, 77–93. [[CrossRef](#)]
8. Visser, D.; Uys, V.; Nieuwenhuis, R.; Pieterse, W. First records of the tomato leaf miner *Tuta absoluta* (Meyrick, 1917) (*Lepidoptera: Gelechiidae*) in South Africa. *BiolInvasions Rec.* **2017**, *6*, 301–305. [[CrossRef](#)]
9. Abbes, K.; Harbi, A.; Chermiti, B. The tomato leafminer *Tuta absoluta* (Meyrick) in Tunisia: Current status and management strategies. *EPPPO Bull.* **2012**, *42*, 226–233. [[CrossRef](#)]
10. Desneux, N.; Wajnberg, E.; Wyckhuys, K.A.G.; Burgio, G.; Arpaia, S.; Narváez-Vasquez, C.A.; Gonzalez-Cabrera, J.; Ruescas, D.C.; Tabone, E.; Frandon, J.; et al. Biological invasion of European tomato crops by *Tuta absoluta*: Ecology, geographic expansion and prospects for biological control. *J. Pest Sci.* **2010**, *83*, 197–215. [[CrossRef](#)]
11. Ouardi, K.; Chouibani, M.; Rahel, M.A.; El Akel, M. Stratégie Nationale de lutte contre la mineuse de la tomate *Tuta absoluta* Meyrick. *EPPPO Bull.* **2012**, *42*, 281–290. [[CrossRef](#)]
12. Pfeiffer, D.G.; Muniappan, R.; Sall, D.; Diatta, P.; Diongue, A.; Dieng, E.O. First Record of *Tuta absoluta* (*Lepidoptera: Gelechiidae*) in Senegal. *Fla. Entomol.* **2013**, *96*, 661–662. [[CrossRef](#)]
13. Mohamed, E.S.I.; Mohamed, M.E.; Gamiel, S.A. First record of the tomato leafminer, *Tuta absoluta* (Meyrick) (*Lepidoptera: Gelechiidae*) in Sudan. *EPPPO Bull.* **2012**, *42*, 325–327. [[CrossRef](#)]

14. Karlsson, F. First report of *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) in the Republic of Benin. *BioInvasions Rec.* **2018**, *7*, 463–468. [[CrossRef](#)]
15. Kiliç, T. First record of *Tuta absoluta* in Turkey. *Phytoparasitica* **2010**, *38*, 243–244. [[CrossRef](#)]
16. Bajracharya, A.S.R.; Mainali, R.P.; Bhat, B.; Bista, S.; Shashank, P.R.; Meshram, N.M. The first record of South American tomato leaf miner, *Tuta absoluta* (Meyrick 1917) (Lepidoptera: Gelechiidae) in Nepal. *J. Entomol. Zool. Stud.* **2016**, *4*, 1359–1363. Available online: <http://www.entomoljournal.com/archives/?year=2016&vol=4&issue=4&ArticleId=1156> (accessed on 4 January 2019).
17. Uulu, T.E.; Ulusoy, M.R.; Çalışkan, A.F. First record of tomato leafminer *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) in Kyrgyzstan. *EPPO Bull.* **2017**, *47*, 285–287. [[CrossRef](#)]
18. Han, P.; Bayram, Y.; Shaltiel-Harpaz, L.; Sohrabi, F.; Saji, A.; Esenali, U.T.; Jalilov, A.; Ali, A.; Shashank, P.R.; Ismoilov, K.; et al. *Tuta absoluta* continues to disperse in Asia: Damage, ongoing management and future challenges. *J. Pest Sci.* **2018**, *92*, 1317–1327. [[CrossRef](#)]
19. Hossain, M.S.; Mian, M.Y.; Muniappan, R. First Record of *Tuta absoluta* (Lepidoptera: Gelechiidae) from Bangladesh. *J. Agric. Urban Entomol.* **2016**, *32*, 101–105. [[CrossRef](#)]
20. Sankarganesh, E.; Firake, D.M.; Sharma, B.; Verma, V.; Behere, G.T. Invasion of the South American Tomato Pinworm, *Tuta absoluta*, in northeastern India: A new challenge and biosecurity concerns. *Entomol. Gen.* **2017**, *36*, 335–345. [[CrossRef](#)]
21. Sharma, P.L.; Gavkare, O. New Distributional Record of Invasive Pest *Tuta absoluta* (Meyrick) in North-Western Himalayan Region of India. *Natl. Acad. Sci. Lett.* **2017**, *40*, 217–220. [[CrossRef](#)]
22. Saidov, N.; Srinivasan, R.; Mavlyanova, R.; Qurbonov, Z. First Report of Invasive South American Tomato Leaf Miner *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Tajikistan. *Fla. Entomol.* **2018**, *101*, 147–149. [[CrossRef](#)]
23. Gebremariam, G. *Tuta Absoluta*: A Global looming challenge in tomato production, Review Paper. *J. Biol. Agric. Healthc.* **2015**, *5*, 57–63. Available online: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.88.3.6592&rep=rep1&type=pdf> (accessed on 8 January 2019).
24. Terzidis, A.; Wilcockson, S.; Leifert, C. The tomato leaf miner (*Tuta absoluta*): Conventional pest problem, organic management solutions? *Org. Agric.* **2014**, *4*, 43–61. [[CrossRef](#)]
25. Gontijo, P.C.; Picanço, M.; Pereira, E.J.G.; Martins, J.; Chediak, M.; Guedes, R.N.C. Spatial and temporal variation in the control failure likelihood of the tomato leaf miner, *Tuta absoluta*. *Ann. Appl. Biol.* **2012**, *162*, 50–59. [[CrossRef](#)]
26. Ayalew, G. Efficacy of Selected Insecticides against the South American Tomato Moth, *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) on Tomato in the Central Rift Valley of Ethiopia. *Afr. Entomol.* **2015**, *23*, 410–417. [[CrossRef](#)]
27. Negeri, T.S.A.M. Evaluation of Some Insecticides against Tomato Leaf Miner, *Tuta absoluta* (Meyrick) (Gelechiidae: Lepidoptera) Under Laboratory and Glasshouse Conditions. *Agric. Res. Technol. Open Access J.* **2017**, *7*. [[CrossRef](#)]
28. Campos, M.R.; Rodrigues, A.R.S.; Silva, W.M.; Silva, T.B.M.; Silva, V.R.F.; Guedes, R.N.C.; Siqueira, H.A.A. Spinosad and the Tomato Borer *Tuta absoluta*: A Bioinsecticide, an Invasive Pest Threat, and High Insecticide Resistance. *PLoS ONE* **2014**, *9*, e103235. [[CrossRef](#)]
29. Guedes, R.N.C.; Picanço, M.C. The tomato borer *Tuta absoluta* in South America: Pest status, management and insecticide resistance. *EPPO Bull.* **2012**, *42*, 211–216. [[CrossRef](#)]
30. Abdel-Baky, N.F.; Al-Soqeer, A.A. Controlling the 2nd Instar Larvae of *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) by Simmondsin Extracted from Jojoba Seeds in KSA. *J. Entomol.* **2017**, *14*, 73–80. [[CrossRef](#)]
31. Balzan, M.V.; Moonen, A.-C. Management strategies for the control of *Tuta absoluta* (Lepidoptera: Gelechiidae) damage in open-field cultivations of processing tomato in Tuscany (Italy). *EPPO Bull.* **2012**, *42*, 217–225. [[CrossRef](#)]
32. Megido, R.C.; Brostaux, Y.; Haubruge, E.; Verheggen, F.J. Propensity of the Tomato Leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae), to Develop on Four Potato Plant Varieties. *Am. J. Potato Res.* **2013**, *90*, 255–260. [[CrossRef](#)]
33. El-Ghany, N.A.; Abdel-Razek, A.S.; Ebadah, I.; Mahmoud, Y. Evaluation of some microbial agents, natural and chemical compounds for controlling tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *J. Plant Prot. Res.* **2016**, *56*, 372–379. [[CrossRef](#)]

34. Abdelgaleil, S.A.; El-Bakary, A.S.; Shawir, M.S.; Ramadan, G.R. Efficacy of various insecticides against tomato leaf miner, *Tuta absoluta*, in Egypt. *Appl. Boil. Res.* **2015**, *17*, 297. [CrossRef]
35. Mahmoud, Y.; Salem, H.; Shalaby, S.; Abdel-Raza, A.; Ebadah, I. Effect of Certain Low Toxicity Insecticides Against Tomato Leaf Miner, *Tuta absoluta* (Lepidoptera: Gelechiidae) with Reference to Their Residues in Harvested Tomato Fruits. *Int. J. Agric. Res.* **2014**, *9*, 210–218. [CrossRef]
36. Cuthbertson, A.G.S.; Mathers, J.J.; Blackburn, L.F.; Korycinska, A.; Luo, W.; Jacobson, R.J.; Northing, P. Population Development of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under Simulated UK Glasshouse Conditions. *Insects* **2013**, *4*, 185–197. [CrossRef]
37. Miranda, M.M.M.; Picanço, M.; Zanoncio, J.C.; Guedes, R.N.C. Ecological Life Table of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Biocontrol Sci. Technol.* **1998**, *8*, 597–606. [CrossRef]
38. Erdoğan, P. Life Table of the Tomato Leaf Miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *J. Agric. Fac. Gaziosmanpasa Univ.* **2014**, *31*, 75. [CrossRef]
39. Venette, R. *Assessment of the Colonization Potential of Introduced Species during Biological Invasions*; University of California: Davis, CA, USA, 1997; Available online: https://scholar.google.com/scholar?hl=en&as_sdt=0,5&cluster=6697658320557623657 (accessed on 6 January 2019).
40. Govorushko, S. *Human–Insect Interactions*; CRC Press: Boca Raton, FL, USA, 2017. [CrossRef]
41. Hastings, A.; Cuddington, K.; Davies, K.; Dugaw, C.J.; Elmendorf, S.; Freestone, A.; Harrison, S.P.; Holland, M.; Lambrinos, J.; Malvadkar, U.; et al. The spatial spread of invasions: New developments in theory and evidence. *Ecol. Lett.* **2004**, *8*, 91–101. [CrossRef]
42. Tonnang, H.E.; Bisseleua, D.H.B.; Biber-Freudenberger, L.; Salifu, D.; Subramanian, S.; Ngowi, V.B.; Guimapi, R.Y.; Anani, B.; Kakmeni, F.M.; Affognon, H.; et al. Advances in crop insect modelling methods—Towards a whole system approach. *Ecol. Model.* **2017**, *354*, 88–103. [CrossRef]
43. Anguelov, R.; Dumont, Y.; Lubuma, J. Mathematical modeling of sterile insect technology for control of anopheles mosquito. *Comput. Math. Appl.* **2012**, *64*, 374–389. [CrossRef]
44. Dufourd, C.; Dumont, Y. Impact of environmental factors on mosquito dispersal in the prospect of sterile insect technique control. *Comput. Math. Appl.* **2013**, *66*, 1695–1715. [CrossRef]
45. Sharov, A.A. Modelling forest insect dynamics. *Caring For. Res. Chang. World. Congr. Rep.* **1996**, *2*, 293–303.
46. McNitt, J.; Chungbaek, Y.Y.; Mortveit, H.; Marathe, M.; Campos, M.R.; Desneux, N.; Brévault, T.; Muniappan, R.; Adiga, A. Assessing the multi-pathway threat from an invasive agricultural pest: *Tuta absoluta* in Asia. *Proc. R. Soc. B Boil. Sci.* **2019**, *286*, 20191159. [CrossRef] [PubMed]
47. Pinet, F.; Bimonte, S.; Miralles, A.; Le Ber, F. Special issue on “Agro-environmental decision support systems”. *Ecol. Inform.* **2015**, *30*, 327. [CrossRef]
48. Tadele, S.; Emanu, G.; Shiberu, T.; Getu, E. Biology of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) under different temperature and relative humidity. *J. Hortic. For.* **2017**, *9*, 66–73. [CrossRef]
49. Duarte, L.; Martínez, M.D.; Helena, V.; Bueno, P. Biology and population parameters of *Tuta absoluta* (Meyrick) under laboratory conditions. *Rev. Protección Veg.* **2015**, *30*, 19–29. Available online: https://www.researchgate.net/profile/Leticia_Duarte/publication/317518366_Biologia_y_parametros_poblacionales_de_Tuta_absoluta_Meyrick_bajo_condiciones_de_laboratorio/links/5a2af6bc45851552ae7a8512/Biologia-y-parametros-poblacionales-de-Tuta-absoluta-Meyr (accessed on 5 January 2019).
50. Pebesma, E.; Bivand, R.; Rowlingson, B.; Gomez-Rubio, V.; Hijmans, R.; Sumner, M.; MacQueen, D.; Lemon, J.; O'Brien, J. *Sp: Classes and Methods for Spatial Data*, 2017. Available online: <https://cran.r-project.org/web/packages/sp/index.html> (accessed on 9 October 2017).
51. Bivand, R.; Keitt, T.; Rowlingson, B.; Pebesma, E.; Sumner, M.; Hijmans, R.; Rouault, E. *Rgdal: Bindings for the “Geospatial” Data Abstraction Library*, 2017. Available online: <https://cran.r-project.org/web/packages/rgdal/index.html> (accessed on 9 October 2017).
52. Hijmans, R.J.; van Etten, J. *Raster: Geographic Data Analysis and Modeling, R Packag.* Version 2.3-12. 2014. Available online: <http://CRAN.R-Project.Org/Package=raster> (accessed on 20 December 2019).
53. Joseph, V.R.; Kang, L. Regression-Based Inverse Distance Weighting with Applications to Computer Experiments. *Technometrics* **2011**, *53*, 254–265. [CrossRef]
54. Balzter, H.; Braun, P.W.; Köhler, W. Cellular automata models for vegetation dynamics. *Ecol. Model.* **1998**, *107*, 113–125. [CrossRef]

55. Xu, X.; Du, Z.; Zhang, H. Integrating the system dynamic and cellular automata models to predict land use and land cover change. *Int. J. Appl. Earth Obs. Geoinf.* **2016**, *52*, 568–579. [CrossRef]
56. Zhao, W. Simulation of agricultural landscape dynamics using cellular automata. In Proceedings of the 2011 19th International Conference on Geoinformatics, IEEE, Shanghai, China, 24–26 June 2011; pp. 1–4. [CrossRef]
57. Moore, E.F. Machine models of self-reproduction. *Am. Math. Soc. Proc. Symp. Appl. Math.* **1970**, *14*, 17–33. Available online: https://books.google.com/books?hl=fr&lr=&id=kCyU6y9XmvQC&oi=fnd&pg=PA17&dq=Edward+F.+Moore++Machines+models+of+self-reproduction&ots=LfjZ4_9Gxk&sig=y8sz0EWluem8YZtljbbwrpAS32Y (accessed on 8 January 2019).
58. Torres, J.B.; Faria, C.A.; Evangelista, W.S.; Pratisoli, D. Within-plant distribution of the leaf miner *Tuta absoluta* (Meyrick) immatures in processing tomatoes, with notes on plant phenology. *Int. J. Pest Manag.* **2001**, *47*, 173–178. [CrossRef]
59. Hulme, P.E. Trade, transport and trouble: Managing invasive species pathways in an era of globalization. *J. Appl. Ecol.* **2009**, *46*, 10–18. [CrossRef]
60. Lounibos, L.P. Invasions by Insect Vectors of Human Disease. *Annu. Rev. Entomol.* **2002**, *47*, 233–266. [CrossRef] [PubMed]
61. Meurisse, N.; Rassati, D.; Hurley, B.P.; Brockerhoff, E.G.; Haack, R.A. Common pathways by which non-native forest insects move internationally and domestically. *J. Pest Sci.* **2018**, *92*, 13–27. [CrossRef]
62. FAO. *FAOSTAT: Production Quantities of Potatoes by Country*; FAO: Rome, Italy, 2020; Available online: <http://www.fao.org/faostat/en/#data/QC/visualize> (accessed on 27 March 2020).
63. Taher, D.; Solberg, S.Ø.; Prohens, J.; Chou, Y.-Y.; Rakha, M.; Wu, T.-H. World Vegetable Center Eggplant Collection: Origin, Composition, Seed Dissemination and Utilization in Breeding. *Front. Plant Sci.* **2017**, *8*. [CrossRef]
64. Cherif, A.; Verheggen, F. A review of *Tuta absoluta* (Lepidoptera: Gelechiidae) host plants and their impact on management strategies. *Biotechnol. Agron. Société Environ.* **2019**, *23*. Available online: <https://orbi.uliege.be/handle/2268/240504> (accessed on 26 March 2020).
65. Biondi, A.; Guedes, R.N.C.; Wan, F.-H.; Desneux, N. Ecology, Worldwide Spread, and Management of the Invasive South American Tomato Pinworm, *Tuta absoluta*: Past, Present, and Future. *Annu. Rev. Entomol.* **2018**, *63*, 239–258. [CrossRef]
66. Tonnang, H.E.Z.; Mohamed, S.F.; Khamis, F.; Ekesi, S. Identification and Risk Assessment for Worldwide Invasion and Spread of *Tuta absoluta* with a Focus on Sub-Saharan Africa: Implications for Phytosanitary Measures and Management. *PLoS ONE* **2015**, *10*, e0135283. [CrossRef]
67. A Silva, G.; Picanço, M.C.; Bacci, L.; Crespo, A.L.B.; Rosado, J.F.; Guedes, R.N.C. Control failure likelihood and spatial dependence of insecticide resistance in the tomato pinworm, *Tuta absoluta*. *Pest Manag. Sci.* **2011**, *67*, 913–920. [CrossRef]
68. Roiditakis, E.; Vasakis, E.; García-Vidal, L.; Martínez-Aguirre, M.D.R.; Rison, J.L.; Haxaire-Lutun, M.O.; Nauen, R.; Tsagkarakou, A.; Bielza, P. A four-year survey on insecticide resistance and likelihood of chemical control failure for tomato leaf miner *Tuta absoluta* in the European/Asian region. *J. Pest Sci.* **2017**, *91*, 421–435. [CrossRef]
69. Guedes, R.N.C. The tomato borer *Tuta absoluta*: Insecticide resistance and control failure. *CAB Rev. Perspect. Agric. Veter. Sci. Nutr. Nat. Resour.* **2012**, *7*. [CrossRef]
70. Never, Z.; Patrick, A.N.; Musa, C.; Ernest, M.; Zekeya, N.; Ndakidemi, P.A.; Chacha, M.; Mbega, E. Tomato Leafminer, *Tuta absoluta* (Meyrick 1917), an emerging agricultural pest in Sub-Saharan Africa: Current and prospective management strategies. *Afr. J. Agric. Res.* **2017**, *12*, 389–396. [CrossRef]
71. Radwan, E.M.; Taha, H.S. Efficacy of Certain Pesticides against Larvae of Tomato Leafminer, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Egypt. Acad. J. Boil. Sci. C, Physiol. Mol. Boil.* **2017**, *9*, 81–95. [CrossRef]
72. Megido, R.C.; Haubruge, E.; Verheggen, F.J. Pheromone-based management strategies to control the tomato leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae). A review. *Biotechnol. Agron. Soc. Environ.* **2013**, *17*, 475–482. Available online: <https://orbi.uliege.be/handle/2268/154676> (accessed on 5 January 2019).
73. Urbaneja, A.; Gonzalez-Cabrera, J.; Arnó, J.; Gabarra, R. Prospects for the biological control of *Tuta absoluta* in tomatoes of the Mediterranean basin. *Pest Manag. Sci.* **2012**, *68*, 1215–1222. [CrossRef]

74. Zappalà, L.; Biondi, A.; Alma, A.; Al-Jboory, I.J.; Arnó, J.; Bayram, A.; Chailleux, A.; El-Arnaouty, A.; Gerling, D.; Guenaoui, Y.; et al. Natural enemies of the South American moth, *Tuta absoluta*, in Europe, North Africa and Middle East, and their potential use in pest control strategies. *J. Pest Sci.* **2013**, *86*, 635–647. [[CrossRef](#)]
75. Giorgini, M.; Guerrieri, E.; Cascone, P.; Gontijo, L. Current Strategies and Future Outlook for Managing the Neotropical Tomato Pest *Tuta absoluta* (Meyrick) in the Mediterranean Basin. *Neotrop. Entomol.* **2018**, *48*, 1–17. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).