

Supplementary Materials: Titanium Nickelide in Midface Fractures Treatment

Liudmila Shamanaeva ^{1,*}, Ekaterina Diachkova ², Pavel Petruk ¹, Kirill Polyakov ¹, Igor Cherkesov ¹ and Sergei Ivanov ^{1,3}

Table S1A. Main characteristics of the research in dynamics.

Period	Paresthesia_1gr	Paresthesia_2gr	Exophthalmos_1gr	Exophthalmos_2gr	Exophthalmos_3gr	Exophthalmos_4gr	Enophthalmos_1gr	Enophthalmos_2gr	Enophthalmos_3gr	Enophthalmos_4gr	Diplopia_1gr	Diplopia_2gr	Diplopia_3gr	Diplopia_4gr
	Mean±SD (Median, Min-Max)													
	Standard error													
before	1.09±0.54 (1,0-2) 0.16	1±0.65 (1,0-2) 0.23	1.36±1.03 (1,0-3) 0.31	1.09±0.97 (1,0-3) 0.28	1.54±0.95 (1,0-3) 0.28	1.36±0.93 (1,0-3) 0.28	0.27±0.8 (0,0-2) 0.19	0.36±0.81 (0,0-2) 0.2	0.09±0.88 (0,1,0-1) 0.09	0.18±0.88 (0,0-1) 0.12	0.55±0.52 (1,0-1) 0.16	0.45±0.5 (0,0-1) 0.16	0.55±0.52 (1,0-1) 0.16	0.45±0.51 (0,0-1) 0.16
Post-operative 30d	0.55±0.65 (1,0-1) 0.16	0.55±0.67 (0,0-2) 0.2	0.27±0.95 (0,0-1) 0.14	0.27±0.89 (0,0-1) 0.14	0.09±0.9 (0,0-1) 0.09	0	0	0.54±0.81 (0,0-2) 0.2	0	0.36±0.69 (0,0-1) 0.15	0.18±0.49 (0,0-1) 0.12	0.27±0.49 (0,0-1) 0.14	0	0.45±0.51 (0,0-1) 0.16
Post-operative 365d	0.09±0.67 (0,0-1) 0.09	0.09±0.66 (0,0-1) 0.09	0	0	0	0	0	0.9±0.88 (1,0-3) 0.28	0	1.18±0.8 (1,0-3) 0.26	0	0.45±0.49 (0,0-1) 0.16	0	0.64±0.51 (1,0-1) 0.15
p	>0.05	>0.05	>0.05	>0.05	<0.01	<0.01	>0.05	<0.05	<0.01	<0.01	>0.05	<0.05	<0.01	<0.05