

Electronic Supplementary Material

A pH-responsive DNA tetrahedron/methotrexate drug delivery system used for rheumatoid arthritis treatment

Yi Jin^{1,2†}, Xingyu Ge^{3†}, Yinjin Xu¹, Siyi Wang¹, Qian Lu¹, Aidong Deng⁴, Jingjing Li^{2,*}, Zhifeng Gu^{4,*}

¹Department of Rheumatology, Affiliated Hospital of Nantong University, Medical School of Nantong University, Nantong, 226000, China; jin_yibai@126.com (Y.J.); 13862959217@163.com (Y.X.); siyiwang5@163.com (S.W.); luqian960927@163.com (Q.L.)

²Research Center of Clinical Medicine, Affiliated Hospital of Nantong University, Nantong, 226000, China

³Department of Rheumatology, Yancheng Third People's Hospital, Yancheng, 224000, China; 15896260521@163.com

⁴Department of Hand Surgery, Affiliated Hospital of Nantong University, Nantong, 226000, China; aidong_deng@163.com

*Correspondence: 13861903880@163.com (J.L.); guzfn@ntu.edu.cn (Z.G.)

†These authors contributed equally to this work.




Figure S1. Characterization of HE-TET nanoparticles. (a) The particle size of TET and HA-TET. (b) The zeta potential of TET and HA-TET. (c) The OD value of TET and HA-TET in different pH.