



Supplementary Materials:

Photocatalytic Inactivation of Plant Pathogenic Bacteria Using TiO₂ Nanoparticles Prepared Hydrothermally

László Kőrösi ^{1,*}, Botond Pertics ², György Schneider ², Balázs Bognár ³, János Kovács ⁴, Vera Meynen ⁵, Alice Scarpellini ⁶, Lea Pasquale ⁷ and Mirko Prato ⁷

¹ Research Institute for Viticulture and Oenology, University of Pécs, Pázmány P. u. 4, H-7634 Pécs, Hungary

² Department of Medical Microbiology and Immunology, Medical School, University of Pécs, Szigeti st. 12, H-7624 Pécs, Hungary; pertics.botond@pte.hu (B.P.); schneider.gyorgy@pte.hu (G.S.)

³ Institute of Organic and Medicinal Chemistry, University of Pécs, Szigeti st. 12, H-7624 Pécs, Hungary; balazs.bognar@aok.pte.hu

⁴ Environmental Analytical and Geoanalytical Research Group, Szentágotthai Research Centre, University of Pécs, Ifjúság u. 20, H-7624 Pécs, Hungary; jones@gamma.ttk.pte.hu

⁵ Laboratory of Adsorption and Catalysis, Department of Chemistry, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium; vera.meynen@uantwerpen.be

⁶ Electron Microscopy Facility, Istituto Italiano di Tecnologia, via Morego 30, 16163 Genova, Italy; alice.scarpellini@iit.it

⁷ Materials Characterization Facility, Istituto Italiano di Tecnologia, via Morego 30, 16163 Genova, Italy; Lea.Pasquale@iit.it (L.P.); Mirko.Prato@iit.it (M.P.)

* Correspondence: korosi.laszlo@pte.hu; Tel.: +36-(72)-517933




Figure S1. UV-Vis spectrum of Blacklight 368 F15W/T8/BL368 UVA lamp (Sylvania) used for the experiments.