

Isolation of Cellulose Nanocrystals from Banana Peel Using One-Pot Microwave and Mild Oxidative Hydrolysis System

Nurhidayah Azmirah Mohd Jamil ^{1,2}, Syafiqah Syazwani Jaffar ¹, Suryani Saallah ^{1,*},
Mailin Misson ¹, Shafiquzzaman Siddiquee ¹, Jumardi Roslan ³ and Wuled Lenggoro ⁴

¹ Biotechnology Research Institute, Universiti Malaysia Sabah, Jalan UMS, Kota Kinabalu 88400, Sabah, Malaysia

² Marine Aquaculture Development Centre Menggatal, Department of Fisheries Sabah, Jalan Sepanggar, Kota Kinabalu 88450, Sabah, Malaysia

³ Faculty of Food Science and Nutrition, Universiti Malaysia Sabah, Jalan UMS, Kota Kinabalu 88400, Sabah, Malaysia

⁴ Institute of Engineering, Tokyo University of Agriculture and Technology, 2-24-16 Nakacho, Koganei, Tokyo 184-8588, Japan

* Correspondence: suryani@ums.edu.my




Figure S1. Suspensions obtained after the H₂O₂-pre-treated banana peel samples were subjected to acid hydrolysis at concentration ranging from 0% (control) to 10%.