

Supplementary Materials

Radiolytic Synthesis of Pt-particle/ABS catalysts for H₂O₂ decomposition in contact lens cleaning

Yuji Ohkubo ^{1,*}, Tomonori Aoki ¹, Satoshi Seino ², Osamu Mori ², Issaku Ito ², Katsuyoshi Endo ¹ and Kazuya Yamamura ¹

¹ Graduate School of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan
[Tel and Fax: +81-6-6879-7294, and E-mail: okubo@upst.eng.osaka-u.ac.jp]

² Menicon Co., Ltd, 5-1-10 Takamori-dai, Kasugai, Aichi 487-0032, Japan

Contents

● Calibration curve for calculating the amounts of Pt	S1
● Photographs of the coloration of peroxotitanium complex	S2
● Calibration curve for calculating the H ₂ O ₂ concentration	S3

Figure S1 Calibration curve for calculating the amounts of Pt on the Pt-particle/ABS samples: relation between Pt concentration and emission intensity.




Figure S2 Photographs of the coloration of peroxotitanium complex: relation between H_2O_2 concentration and the color of peroxotitanium complex.




Figure S3 Calibration curve for calculating the H₂O₂ concentration: relation between H₂O₂ concentration and absorbance at 407 nm.

