

Supplementary

Behavioral Effect of Terahertz Waves in C57BL/6 Mice

Miao Qi ^{1,2}, Rong Liu ^{1,2}, Bing Li ^{1,2}, Shuai Wang ^{1,2}, Runze Fan ¹, Xinyi Zhao ^{1,2} and Dehui Xu ^{1*}

¹ State Key Laboratory of Electrical Insulation and Power Equipment, Centre for Plasma Biomedicine, Xi'an Jiaotong University, Xi'an, Shaanxi, 710049, China; qimiao@stu.xjtu.edu.cn (M.Q.); liurong0812@stu.xjtu.edu.cn (R.L.); lb905533242@stu.xjtu.edu.cn (B.L.); shuaiwang3059@stu.xjtu.edu.cn (S.W.); fanrunze@stu.xjtu.edu.cn (R.F.); zhaoxinyi@stu.xjtu.edu.cn (X.Z.)

² The school of Life Science and Technology, Xi'an Jiaotong University, Xi'an, Shaanxi, 710049, China.

* Correspondence: dehuixu@mail.xjtu.edu.cn

Citation: Qi, M.; Liu, R.; Li, B.; Wang, S.; Fan, R.; Zhao, X.; Xu, D. Behavioral Effect of Terahertz Waves in C57BL/6 Mice. *Biosensors* **2022**, *12*, 79. <https://doi.org/10.3390/bios12020079>

Received: 05 January 2022

Accepted: 26 January 2022

Published: 28 January 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure S1. Analysis of the open field test and forced swim test after THz treatment. (A) moved pathways (B) and distance in center in the open field test; (C) immobility time (D) and swimming time in forced swim test.