

Estrogen and Progesterone Receptors are Dysregulated at the BPH/5 Mouse Preeclamptic-Like Maternal–Fetal Interface

Viviane C. L. Gomes ¹, Bryce M. Gilbert ¹, Carolina Bernal ¹, Kassandra R. Crissman ², and Jenny L. Sones ^{3*}

¹ Department of Small Animal Clinical Sciences, Michigan State University College of Veterinary Medicine, East Lansing, MI 48824, United States.

² Department of Veterinary Clinical Sciences, Louisiana State University School of Veterinary Medicine, Baton Rouge, LA 70803, United States.

³ Colorado State University Equine Reproduction Laboratory, Fort Collins, CO 80521, United States.

* Correspondence: jenny.sones@colostate.edu

Supplementary Material

Table S1. Forward and reverse *Mus musculus*-specific primer sequences used in qRT-PCR

Gene	Primer Sequence
<i>18S</i>	F: GTAACCCGTTGAACCCCATTT R: CCATCCAATCGGTAGTAGCG
<i>Esr1</i>	F: AAT TCT GAC AAT CGA CGC CAG R: GTG CTT CAA CAT TCT CCC TCC TC
<i>Esr2</i>	F: CGG TCT GTC TGA ATG TGG TCA CTG R: TTG CCT TGG TCA TGG TAT CGC TTC
<i>Pr-A/B/C</i>	F: GAC ACT GGC TGT GGA ATT TCC R: CCA GGA TCT TGG GCA ACT G
<i>Pr-A/B</i>	F: CTG GAG ACC GAG GGC TCT R: CCA GTG CTC GAG GTT TGC TC
<i>Pr-B</i>	F: GTG GAG GGC GCT TTC TCT G R: TCT GCC TCC CTC CCT ATG AGT