

Article

# Thermal Resilience of Feeding Kinematics May Contribute to the Spread of Invasive Fishes in Light of Climate Change

Ralph Turingan \* and Tyler Sloan

Department of Biological Sciences, Florida Institute of Technology, 150 West University Boulevard, Melbourne, FL 32901, USA; turingan@fit.edu

\* Correspondence: Turingan@fit.edu; Tel.: +1-321-674-8037

Academic Editor: Chris O'Callaghan

Received: 30 September 2016; Accepted: 17 November 2016; Published: 25 November 2016

**Abstract:** As a consequence of global warming, tropical invasive species are expected to expand their range pole-ward, extending their negative impacts to previously undisturbed, high-latitude ecosystems. Investigating the physiological responses of invasive species to environmental temperature is important because the coupled effects of climate change and species invasion on ecosystems could be more alarming than the effects of each phenomenon independently. Especially in poikilotherms, the rate of motion in muscle-driven biomechanical systems is expected to double for every 10 °C increase in temperature. In this study, we address the question, “How does temperature affect the speed of jaw-movement during prey-capture in invasive fishes?” Kinematic analysis of invasive-fish prey-capture behavior revealed that (1) movement velocities of key components of the feeding mechanism did not double as water temperature increased from 20 °C to 30 °C; and (2) thermal sensitivity (Q10 values) for gape, hyoid, lower-jaw rotation, and cranial rotation velocities at 20 °C and 30 °C ranged from 0.56 to 1.44 in all three species. With the exception of lower-jaw rotation, Q10 values were significantly less than the expected Q10 = 2.0, indicating that feeding kinematics remains consistent despite the change in environmental temperature. It is conceivable that the ability to maintain peak performance at different temperatures helps facilitate the spread of invasive fishes globally.

**Keywords:** global warming; prey-capture; suction feeding; organismal performance; thermal tolerance

## 1. Introduction

Projected variations in global temperature as a consequence of climate change have underscored renewed interest in addressing questions about temperature-dependent organismal performance [1–5]. The re-emergence of emphasis on ecological physiology of organisms is rooted in the notion that physiological traits are strong determinants of species response to climate change [6–13]. In light of the climate change phenomenon, what mechanisms underlie organismal response to environmental temperature variations? In an attempt to establish the foundation for continuing investigations addressing this central question, this study was designed to explore the effects of temperature on whole-organism performance, specifically prey-capture kinematics, in three orders of invasive-teleost fishes. To our knowledge, these are the first three invasive species used to investigate the effects of environmental temperature on feeding kinematics in light of climate change. The use of invasive fishes to examine temperature-dependent performance increases the relevance of this study because of the belief that the coupled effects of climate change and species invasion on ecosystems are more alarming than the effects of each taken independently [14]. It is predicted that, as a consequence of global warming, tropical invasive species expand their range poleward, thus, extending their well-known negative biological impacts to previously undisturbed, high-latitude ecosystems [14–16].

The effects of environmental temperature on physiological processes are widespread in both animals and plants, but in fishes its effects on skeletal-muscle performance and property have been the focus of most basic and applied physiological studies. It is expected that the direct effects of environmental temperature on body temperature mediate physiological processes, most especially metabolic rate. In fishes, as with any poikilothermic vertebrate, physiological performance peaks at a narrow range of body temperature, and environmental temperature remains a major constraint in their range of distribution [4,5,17–20]. Metabolic rate, in general, and contraction rate of skeletal muscle fibers, in particular, increase with environmental temperature until a threshold level is achieved, after which these rates decline with further increase in environmental temperature [18,21–26]. The ability of an individual to perform a certain task (e.g., prey capture) driven by a temperature-dependent process (e.g., rate of contraction of buccal-expansion muscles) is constrained by the reduction in biological rates as a consequence of decrease in environmental temperature. Furthermore, the expression of temperature-induced changes in fish-muscle physiology varies according to different temporal scales (i.e., daily, seasonal, or life-history, developmental time scales) and levels of organization (i.e., from molecular to organismal to ecosystem) [27]. For example, intracellular and extracellular ionic concentration and acid balance are destabilized by instant changes in environmental temperature [28]. Seasonal temperature change may induce modifications of the properties and composition of the contractile elements of the muscle fiber [27]. This time scale may allow fish to acclimate to the new ambient temperature and drive plastic response in muscle-fiber phenotypes and trigger behavioral and other mechanisms that buffer temperature effects and maintain homeostasis [28–30]. At the whole-organism level, responses to environmental temperature change may be taxon- and ontogenetic- specific. Indeed, environmental temperature has a profound influence on the fish's ability to successfully accomplish relevant tasks such as swimming, feeding, mating, and escaping from predators.

Interestingly, previous studies investigating temperature effects on whole-organism performance in fishes have revealed mixed results. The kinematics of routine-swimming in teleost fishes have been largely consistent with physiological expectations. For example, swimming velocity doubles in response to a 10 °C increase in environmental temperature, that is, Q<sub>10</sub> values are at least 2.0 [31–34]. The physiological quotient (Q<sub>10</sub> value) indicates the magnitude of change in biological rates (e.g., swimming speed) for every 10 °C change in temperature [31–34]. An attempt to arrive at a consensus on the effects of temperature on feeding kinematics in teleost fishes remains unsuccessful perhaps because of the differences in the experimental design employed by the relatively few studies investigating this subject matter. In investigating how decrease or increase in environmental temperature, between 18 °C and 24 °C, affect prey-capture performance, Wintzer and Motta [19] concluded that it took bluegill (*Lepomis macrochirus*) longer to achieve maximum gape and lower-jaw rotation as water temperature decreased. DeVries and Wainwright [20] found that a 15 °C decrease in temperature caused only the time to reach maximum gape, among all timing parameters that underlie suction feeding performance in largemouth bass (*Micropterus salmoides*), to slightly increase. In Sloan and Turingan [35], and Turingan and Sloan [36], Repeated Measures Multivariate Analysis of Covariance revealed that environmental temperature, raised from 20 °C to 30 °C at a rate of 1 °C daily, had no effect on the magnitude and timing of prey-capture kinematics of nonnative teleost fishes in Florida, USA. Considering that the ability to successfully capture prey determines individual survivorship, it is imperative to elucidate how environmental temperature mediates prey-capture performance in organisms.

Three invasive Florida fishes—Pike killifish, *Belonesox belizanus* (Cyprinodontiformes); lionfish, *Pterois volitans* (Scorpaeniformes); and Mayan cichlid, *Cichlasoma urophthalmus* (Perciformes)—were used to determine how variable or consistent whole-organism response is, particularly in invasive species, to environmental temperature. Consistent with the prediction of climate-driven range expansion of invasive-species poleward, these invasive fishes have continued to extend their distribution northward from where they were introduced in south Florida [37–40]. The average

annual temperature within the current distribution of these invasive fishes ranges between 20 °C and 30 °C [41]. This is used as the basis for the selection of experimental temperatures in this study.

Identified as one of the most abundant nonnative fishes in Florida, *Belonesox belizanus* is native to waters (temperature is 25–37 °C) in Mexico and Central America, and was introduced into a ditch in south Florida in the late 1950s [37–43]. It is a specialist predator, feeding on small fishes using a feeding mechanism that is well designed for piscivory. This piscivore is capable of achieving a large gape with its elongated premaxillae and mandibles lined with large teeth [44–46]. The ability of pike killifish to independently rotate its premaxilla posterodorsally, facilitated by the premaxillomandibular ligament and a twisting maxilla, further enhances gape formation [45].

*Pterois volitans* is native to the Indo-Pacific Ocean (temperature is 22–28 °C) [47]. After its initial introduction in south Florida in the early 1990s, it has rapidly expanded its invasive population southward to the Caribbean and northward along the Atlantic coast of the USA [48–51]. The invasive lionfish has been identified as the likely worst threat to marine biodiversity in the Mid-Atlantic, Gulf of Mexico, and Caribbean regions [48–51]. This invasive predator feeds on a diverse group of fishes in the region, including 21 families and 41 species of teleost fishes, the majority of which have commercial, recreational, and ecological importance [50,52]. The predatory success of the lionfish is perhaps enhanced by its ability to modulate its suction-feeding repertoire, including a characteristic rapid-strike on more mobile, elusive fish and crustacean prey [53].

The native distribution of *Cichlasoma urophthalmus* ranges from eastern Mexico to Nicaragua (temperature is 22–39 °C) [54]. Following its introduction into south Florida in the 1980s, it has spread into north Florida and the Florida Bay regions [55–57]. Perhaps among the traits that enable the invasive Mayan cichlid to spread northward in Florida is its tolerance to extreme variations in salinity [55–58] and temperature [56,58]. In addition, the invasive Mayan cichlid has a generalist diet, feeding on detritus, plants, invertebrates and fish [59–62]. Its feeding apparatus includes an oral-jaw mechanism for prey capture and a well-developed pharyngeal-jaw apparatus for prey-processing [63].

This study was designed to test the hypothesis that the velocity of prey-capture kinematics, particularly buccal expansion and compression behaviors powered by skeletal-muscle in teleost fishes doubles when ambient-water temperature is raised by 10 °C.


## 2. Materials and Methods

Four *B. belizanus*, collected from the Florida Everglades National Park, four *C. urophthalmus*, collected from Merritt Island, Florida, and four *P. volitans*, collected from Port St. Lucie, Florida were acclimated to 20 °C water temperature and trained for high-speed video in the fish ecophysiology laboratory at Florida Institute of Technology for two weeks before the experiment was initiated. Each fish was housed in 38 L filming tanks filled with water that matched their Florida habitats: 0 ppt for *B. belizanus*; 24 ppt for *C. urophthalmus*, and 35 ppt for *P. volitans*. The twelve fishes were subjected to a repeated measures experimental design, in which, each fish was filmed successively in each of the three experimental temperatures 20 °C, 25 °C and 30 °C (Figure 1). Temperature in each filming tank was raised from 20 °C to the higher filming temperatures at a rate of 1 °C daily using a water heater. Once the experimental temperature was achieved, feeding sessions were recorded every other day from each fish. Each fish was filmed using a RedLake High-Speed Motionscope 2000S camera with a shutter speed of 1/1000 s at 250 frames per second while feeding on live mosquitofish (*Gambusia holbrooki*) prey at 20 °C, 25 °C and 30 °C. Prey was maintained at 20 °C ambient-room temperature. Prey girth was about 80% of peak gape of each fish; previous analyses of feeding kinematics in these fishes indicated that this relative prey-size elicited maximum prey-capture performance in fish [35,36,46]. The effects of temperature on the prey was not investigated in this study. Experimental fish was not fed 1–2 days before each recording session to ensure fish was motivated to eat and exhibit maximum performance during feeding trials [19,20].

Each fish was filmed until at least 10 feeding bouts were recorded in which the fish was perpendicular to the camera and exhibited maximum prey-capture performance. The best four films

were analyzed per fish at each of the three experimental temperatures using MaxTRAQ (Version 2.2.4.1 Innovision Systems, Inc., Columbiaville, MI, USA). Each film was played back frame-by-frame to measure maximum gape (mm), hyoid depression (mm), lower-jaw rotation (degree), and cranial rotation (degree), as well as the time (ms) to reach each of these maximum kinematic-displacement variables. Time to reach each of these maximum kinematic-displacement variables was calculated using the frame prior to mouth opening as time = 0. Reference points (i.e., kinematic hotspots) used to measure these variables are illustrated in Figure 2. Average velocity was calculated as the value of the maximum kinematic-displacement variable divided by the corresponding time to reach this maximum displacement: Gape Velocity = Maximum Gape  $\div$  Time to Reach Maximum Gape; Hyoid Velocity = Maximum Hyoid Depression  $\div$  Time to Reach Maximum Hyoid Depression; Lower-Jaw Rotation Velocity = Maximum Lower-Jaw Rotation  $\div$  Time to Reach Maximum Lower-Jaw Rotation; Cranial Rotation Velocity = Maximum Cranial Rotation  $\div$  Time to Reach Maximum Cranial Rotation. Physiological quotient ( $Q_{10}$ ) was calculated for each of the kinematic-velocity variables as  $Q_{10} = (\text{Kinematic Velocity at } 30\text{ }^{\circ}\text{C} / \text{Kinematic Velocity } 20\text{ }^{\circ}\text{C})$  (modified from Schmidt-Nielsen [64]).

## Experimental Design



**Figure 1.** Diagram depicting the experimental design investigating the effects of environmental temperature on the kinematic velocity to reach maximum gape, hyoid depression, cranial rotation, and lower-jaw rotation. Four individuals of each invasive species *Belonesox belizanus*, *Pterois volitans*, and *Cichlasoma urophthalmus* were filmed while feeding on live-fish prey at 20 °C, 25 °C, and 30 °C using high-speed video. The best four films of each individual feeding at each temperature were digitized to measure the four kinematic velocities stated above and to calculate  $Q_{10}$  values. Kinematic velocities and  $Q_{10}$  values were subjected to the appropriate statistical tests to determine the effects of temperature on prey-capture performance.

Each of the four kinematic-velocity variables was subjected to a Model I least-squares regression against temperature to define the model  $y = a + bx$ ; where  $y$  = kinematic velocity,  $a$  = intercept,  $b$  = slope, and  $x$  = environmental temperature (=20 °C, 25 °C, and 30 °C). To test the hypothesis that the  $Q_{10}$  of each of the kinematic-velocity variables was different from 2.0, a series of Paired t-Tests were conducted to compare the empirical  $Q_{10}$  values with the theoretical  $Q_{10}$  value of 2.0. All statistical tests were conducted using R.

After the experiment, each fish was sacrificed using an overdose of MS-222 solution, fixed in 10% formalin solution, and then stored in 75% ethanol solution. All specimens have been stored appropriately in the fish ecophysiology and evolution laboratory at Florida Institute of Technology for use in current and future teaching and research. All procedures for housing, maintaining and sacrificing experimental fishes strictly followed the guidelines and procedures of the Institutional Animal Care and Use Committee (IACUC) of the Florida Institute of Technology (IACUC Approval # 101202).




**Figure 2.** Diagram of the pike killifish, *Belonesox belizanus* (**top**), lionfish, *Pterois volitans* (**middle**), and Mayan cichlid, *Cichlasoma urophthalmus* (**bottom**) showing the homologous hotspots used to measure peak gape (= maximum distance measured from the anteriormost tip of the premaxilla (A) to the anteriormost tip of the dentary (C)), peak hyoid depression (= maximum distance between the center of the eye (E) to the anteriormost tip of the hyoid bar (D)), peak lower-jaw rotation (= maximum posteroventral rotation of the lower-jaw, measured as the angle formed by line segments AB (= jaw-joint) to BC), and peak cranial rotation (= maximum posterodorsal rotation of the neurocranium, measured by the angle formed by line segments AG (= dorsal tip of the pectoral-fin base) to GF (= anterior tip of the dorsal-fin base)).

### 3. Results


The three Florida invasive-fish species fed voraciously upon introduction of the prey during feeding-recording sessions at each of the three environmental temperatures, 20 °C, 25 °C, and 30 °C (Figure 3). Examination of the films revealed that the feeding behavior of each fish was consistent

with previous studies [35,36,46,53]. The pike killifish stalked its prey and within a very short distance between the predator and the prey, the fish lunged toward the prey, opened its mouth widely and snapped at the prey. The lionfish used its pectoral fins to herd the prey closer to its mouth before suction feeding to capture prey. The Mayan cichlid behaved more aggressively; as soon as the prey was introduced into the filming tank, the cichlid rapidly swam toward the prey and suction-fed on it instantaneously.



**Figure 3.** Select frames of representative films of two lionfish showing the sequence of kinematic events during prey capture in lionfish, *Pterois volitans*, at 20 °C, and 30 °C. Note that all species of invasive fishes successfully captured prey in both temperatures.

Linear regression models indicated that the average velocity of kinematic events during feeding in all three invasive species remained consistent across environmental temperatures, with the exception of the average velocity of hyoid depression in the Mayan cichlid and average velocity of lower-jaw rotation in the lionfish. Species-specific variation in elevation (=y-intercept, a) is apparent, but, the slopes of the regression, b, were not statistically different from zero, indicating that kinematic velocities were unaffected by environmental temperature (Table 1 and Figure 4).




**Figure 4.** Scatterplot showing the relationship between each of the four kinematic-velocity variables and feeding temperature 20 °C, 25 °C, and 30 °C. Results of the regression analysis that quantified the effect of temperature on each of the kinematic events are presented in Table 1.

**Table 1.** Results of the regression analysis examining the effects of temperature on the kinematic velocities of maximum gape, hyoid depression, cranial rotation and lower-jaw rotation.

Kinematics	Species	a	b	r <sup>2</sup>	p
Gape	P	0.245	0.0343	0.0734	0.163
	L	1.797	−0.0357	0.0859	0.130
	M	1.821	−0.0376	0.0893	0.122
Hyoid Depression	P	0.393	−1.14 × 10 <sup>−3</sup>	4.53 × 10 <sup>−4</sup>	0.914
	L	0.738	−0.0162	0.0853	0.131
	M	0.980	−0.0267	0.227	0.010
Cranial Rotation	P	11.385	−0.243	0.0419	0.296
	L	8.808	−0.107	9.38 × 10 <sup>−3</sup>	0.624
	M	−1.817	0.368	0.101	0.099
Lower-Jaw Rotation	P	13.365	−0.179	0.0286	0.390
	L	−3.568	0.610	0.233	0.009
	M	8.089	0.0664	3.61 × 10 <sup>−3</sup>	0.761

Coefficients defining the regression equation,  $y = a + bx$  are shown.  $y$  = velocity to reach maximum Gape, Hyoid Depression, Cranial Rotation, and Lower-Jaw Rotation.  $a$  = intercept;  $b$  = slope.  $R^2$  = Coefficient of Determination.  $p$  = probability associated with the null hypothesis that the slope of each regression is significantly different from zero. P = Pike killifish, *Belonesox belizanus*; L = Lionfish, *Pterois volitans*; M = Mayan cichlid, *Cichlasoma urophthalmus*.

The mean Q10 values of each of the kinematic velocities were significantly less than the expected Q10 value of 2.0, with the exception of the average velocity of lower-jaw rotation in the lionfish and Mayan cichlid (Table 2 and Figure 5).



**Figure 5.** Mean Q10 values for kinematic-velocity in each of the three invasive-fish species. Note that except for the mean Q10 values for lower-jaw rotation in the Mayan cichlid and lionfish, all Q10 values of kinematic velocity were significantly less than the expected Q10 value of 2.0 for fish feeding at 20 °C and 30 °C. Error bars indicate standard error of the mean.

**Table 2.** Results of the Paired *t*-Tests comparing the difference between mean Q10s of each of the kinematic velocity variables and the expected value of 2.0.

Kinematics	Species	Mean Q10	<i>t</i> -Statistic	Df	<i>p</i>
Gape	P	0.973	−7.6174	3	0.005
	L	0.922	−6.4448	3	0.008
	M	1.046	−4.3859	3	0.022
Hyoid	P	1.047	−30.0273	3	$8.113 \times 10^{-5}$
	L	0.559	−67.1036	3	$7.239 \times 10^{-6}$
	M	0.876	−9.8617	3	0.002
Cranial	P	1.049	−6.8275	3	0.006
	L	1.054	−9.7247	3	0.002
	M	1.033	−19.0769	3	$3.145 \times 10^{-4}$
Lower-Jaw	P	1.004	−6.9719	3	0.006
	L	1.294	−2.8106	3	0.063
	M	1.444	−1.6617	3	0.195

Relevant statistics are shown for P = Pike killifish, *Belonesox belizanus*; L = Lionfish, *Pterois volitans*; M = Mayan cichlid, *Cichlasoma urophthalmus*. Note that except for the Q10 of the velocity of lower-jaw rotation in Mayan cichlid and lionfish, all Q10 values of kinematic velocities are statistically less than the expected value of 2.0.

#### 4. Discussion

Empirical evidence of how organismal performance is affected by environmental change advances our understanding of the consequences of climate change and invasion of nonnative species. Investigations into the combined effects of both phenomena on native-community structure and dynamics, as well as range expansion of invasive species, are especially important considering that their combined effects are perhaps more devastating than each taken independently ([64], [www.invasivespecies.gov](http://www.invasivespecies.gov)). It has been predicted that as a consequence of the pole-ward warming

of the earth, tropical-invasive species will continue to expand their range toward higher-latitude ecosystems at alarming rates [14,65]. Well known characteristics of invasive species that enable them to impart damage in the stability of native ecosystems include: (1) they have high propensity to introduce and spread diseases [66–68]; (2) they alter community and food web structure through competition and predation [38,69–73]; (3) they hybridize with native species [74–76]; and (4) they outcompete native species and ultimately displace and even drive native species to extinction [1,2,67,69]. The latter results in the reduction of biodiversity and may even lead to biological homogenization. Although empirical evidence demonstrating the direct and indirect interactions between invasive and native species, as well as the cause-effect mechanism underlying post-invasion changes in community structure of invaded ecosystems, are elusive, it is hypothesized that climate change exacerbates the negative impacts of invasive species to ecology and society [77,78]. In our attempt to contribute to the advancement of our understanding of the impacts of the invasive species and climate change coupling to ecosystem dynamics, our discussion of the results of this study centers on the question, “How do invasive fishes deal with variations in temperature within their invasive range?”

As poikilotherms, the feeding performance of invasive fishes are expected to conform with the known effects of temperature on the physiology and ecology of heterothermic, aquatic animals [79]. For example, first, the velocity of fin propulsion during swimming and mouth-opening during feeding, behaviors fueled by skeletal-muscle contraction and relaxation, are expected to double when ambient temperature is increased by 10 °C. This is because at the physiological level of analysis, there is a two-fold increase in the rate of muscular contraction and relaxation for every 10 °C increase in temperature (i.e.,  $Q_{10} = 2.0$ ) [18,21–24,31]. Second, at the ecological level of analysis, predictable seasonal cooling and warming of lakes and rivers have contributed to the evolution of acclimatization in teleost fishes [80–82]. Third, the food habit of some temperate fishes, such as largemouth bass, *Micropterus salmoides*, and pumpkinseed sunfish, *Lepomis gibbosus*, change seasonally, consistent with the seasonal cooling and warming of lakes or rivers in temperate ecosystems [80–82].

Results of this study, as well as those of Sloan and Turingan [35] and Turingan and Sloan [36] underscore the thermal independence of prey-capture performance in invasive-teleost fishes. Suction-feeding, which is the most dominant and generalized mode of prey-capture in teleost fishes, relies primarily on the high-speed movement of cranial elements such as the jaws, hyoid, suspensorium, and cranium during mouth opening and closing (see Figure 3; [46,83–89]). A successful strike and capture of prey relies heavily on the almost simultaneous and rapid expansion of the buccal cavity to generate subambient-pressure in the buccal chamber and mouth opening [83–85]. These kinematic events are accomplished primarily by the posterodorsal rotation of the cranium, depression of the hyoid apparatus, lateral extension of the suspensorial and opercular bones, and the posteroventral rotation of the lower-jaws [83–89]. These cranial movements are driven by skeletal muscles including the epaxialis, sternohyoideus, retractor arcus palatini, levator operculi, and dilator operculi [83–85,87]. It is well known that the rate of contraction of skeletal muscle is expected to at least double for every 10 °C increase in environmental and body temperature in ectotherms such as teleost fishes [20,31,33,34,90–92]. However, temperature has no significant effects on the velocity of movement of the key elements of the prey-capture mechanism in the three contrasting models of invasive species reported here. Although, on average, prey-capture kinematics significantly differ among the three invasive species, the absence of a temperature-induced change in kinematic velocity is evident in all three invasive species [35,36].

Thermal independence of fast-start behaviors and kinematics has been found in other vertebrate animals. Navas et al. [93] concluded that the  $Q_{10}$  for “jump take-off velocity and mean swimming velocity” in the frog *Rana temporaria* was lower than  $Q_{10} = 2.0$ . “Running velocity during burst activity” in several species of lizards were less affected by temperature, as evident in the low  $Q_{10}$  values for this behavior [94–96]. “Ballistic mouth opening and tongue projection dynamics” in toads (*Bufo terrestris*) were thermally independent [91]. Lack of temperature-induced variation in the dynamics of “ballistic mouth opening” was also evident in the frog *Rana pipiens* [96].

It is noteworthy that our results do not agree with the conclusions of the only two papers reporting the effects of temperature on the feeding kinematics of native teleost fishes. Prey-capture kinematics in North American native centrarchid fishes, bluegill *Lepomis microchirus*, and largemouth bass *Micropterus salmoides*, responded to environmental-temperature change in a manner that is consistent with physiological predictions [19,20]. Among other feeding-kinematic variables that were affected by temperature, it took longer for fishes to reach peak gape in colder than in warmer temperatures [19,20]. The prevalence of thermally independent prey-capture kinematics in invasive teleost fishes underscores the need to address how they compensate for the effects of temperature on the contractile properties and contraction velocity of skeletal muscles. A direct comparison between any of the invasive fishes in this study and an ecologically relevant (e.g., as a competitor or prey) native species in Florida is imperative, given the need to address the direct impacts of invasive on native species.

Translation of the physiological effects of temperature on muscle contractile properties to whole-animal performance may be mitigated by the central nervous system [27,97–99]. Central nervous-system governed compensatory mechanisms may allow whole animals to perform at optimum levels despite variation in environmental and body temperatures [100–102]. Such compensatory mechanisms likely include (1) plasticity in the recruitment of muscle fiber types [102–105]; (2) involvement of elastic strain energy storage and recovery in muscular and tendinous tissue [106,107]; (3) occurrence of temperature-induced change in acid-base balance in muscle fiber [107,108]; (4) plasticity of thermal sensitivity of myofibrillar ATPase activity [107,108]. For example, Rome et al. [104] concluded that at lower temperatures, carp (*Cyprinus carpio*) recruited more fast-contracting (fast anaerobic) muscle fibers when environmental temperature was lower than ambient. Navas et al. [93] found that at 10 °C, optimal whole animal performance was accomplished by only 34% of muscle-power output. In the sartorius muscle of the toad *Bufo bufo*, Renaud and Stevens [107] demonstrated that short-term change in intracellular pH associated with decrease in water temperature from 25 °C to 5 °C was enough “to increase maximum force and, hence, power during isotonic shortening of muscle fiber, providing a short-term mechanism for compensation to low temperature”.

Global-climate change in general and global-temperature change in particular have important consequences for the performance of invasive species because of (1) the temperature-induced effects on physiological and mechanical processes [79,108]; (2) the likelihood that these physiological effects extend to whole-organism performance (e.g., [19,20]); and (3) the resilience of invasive species and the resistance of whole-organism performance to temperature change ([35,36]; this study). These plausible avenues where the interplay between climate-change and invasive-species phenomena may be demonstrated need further investigation and confirmation.

## 5. Conclusions

The velocity of jaw movements during prey capture in the invasive fishes, *Belonesox belizanus*, *Pterois volitans*, and *Cichlasoma urophthalmus*, were statistically unaffected by water temperature. Within the range of temperature used in this study, all invasive fishes successfully captured their prey using a stereotypical suction-feeding kinematic pattern that is unaltered by temperature. It is plausible that this seemingly temperature-resilient behavior will facilitate the successful expansion of the invasive range of these tropical-fish species as a consequence of global warming.

**Acknowledgments:** We thank Brian Bement for illustrating Figure 2. Lisa Young, Chelsea Harms, Benjamin Compton, Kayla Chapman, Brian Bement, and Matthew Sonnefeld helped with the recording of fish-feeding. This study was partially funded by Florida Sea Grant and Sigma Xi.

**Author Contributions:** Ralph Turingan conceived of the idea and wrote the manuscript. Tyler Sloan conducted all statistical analyses. Turingan and Sloan designed and conducted the experiments.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Davis, M.B.; Shaw, R.G. Range shifts and adaptive responses to Quaternary climate change. *Science* **2001**, *292*, 673–679. [[CrossRef](#)] [[PubMed](#)]
2. Baker, C.; Starger, C.J.; McClanahan, T.R.; Glynn, P.W. Coral reefs: Corals' adaptive response to climate change. *Nature* **2004**. [[CrossRef](#)] [[PubMed](#)]
3. Portner, H.O.; Farrell, A.P. Physiology and climate change. *Nature* **2008**, *322*, 690–692.
4. Portner, H.O. Integrating climate-related stressor effects on marine organisms: Unifying principles linking molecule to ecosystem-level changes. *Mar. Ecol. Prog. Ser.* **2012**, *470*, 273–290. [[CrossRef](#)]
5. Huey, R.B.; Kingsolver, J.G. Evolution of resistance to high temperature in ectotherms. *Am. Nat.* **1993**, *142*, S21–S46. [[CrossRef](#)]
6. Chown, S.L.; Gaston, K.J.; Robinson, D. Macrophysiology: Large-scale patterns in physiological traits and their ecological implications. *Func. Ecol.* **2004**, *18*, 159–167. [[CrossRef](#)]
7. Helmuth, B.; Kingsolver, J.G.; Carrington, E. Biophysics, physiological ecology, and climate change: Does mechanism matter? *Ann. Rev. Physiol.* **2005**, *67*, 177–201. [[CrossRef](#)] [[PubMed](#)]
8. Deutsch, C.A.; Tewksbury, J.J.; Huey, R.B.; Sheldon, K.S.; Ghalambor, C.K.; Haak, D.C.; Martin, P.R. Impacts of climate warming on terrestrial ectotherms across latitude. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 6668–6672. [[CrossRef](#)] [[PubMed](#)]
9. Huey, R.B.; Deutsch, C.A.; Tewksbury, J.J.; Vitt, L.J.; Hertz, P.E.; Perez, J.A.; Garland, T. Why tropical forest lizards are vulnerable to climate warming. *Proc. R. Soc. B* **2009**, *276*, 1939–1948. [[CrossRef](#)] [[PubMed](#)]
10. Kearney, M.R.; Porter, W.P. Mechanistic niche modeling: Combining physiological and spatial data to predict species ranges. *Ecol. Lett.* **2009**, *12*, 334–350. [[CrossRef](#)] [[PubMed](#)]
11. Dillion, M.E.; Wang, G.; Huey, R.B. Global metabolic impacts of recent climate warming. *Nature* **2010**, *467*, 704–706. [[CrossRef](#)] [[PubMed](#)]
12. Doak, D.F.; Morris, W.F. Demographic compensation and tipping points in climate-induced range shifts. *Nature* **2010**, *467*, 959–962. [[CrossRef](#)] [[PubMed](#)]
13. Smith, A.L.; Hewitt, N.; Klenk, N.; Bazely, D.R.; Yan, N.; Wood, S.; Henriques, I.; MacLellan, J.I.; Lipsig-Mumme, C. Effects of climate change on the distribution of invasive alien species in Canada: A knowledge synthesis of range change projections in a warming world. *Environ. Rev.* **2012**, *20*, 1–16. [[CrossRef](#)]
14. Perrings, C.; Dehnen-Shmutz, K.; Touza, J.; Williamson, M. How to manage biological invasions under globalization. *Trends Ecol. Evol.* **2005**, *20*, 212–215. [[CrossRef](#)] [[PubMed](#)]
15. Hellmann, J.J.; Byers, J.E.; Bierwagen, B.G.; Dukes, J.S. Five potential consequences of climate change for invasive species. *Conserv. Biol.* **2008**, *22*, 534–543. [[CrossRef](#)] [[PubMed](#)]
16. Shafland, P.L.; Pestrak, J.M. Lower lethal temperatures for fourteen nonnative fishes in Florida. *Environ. Biol. Fishes* **1982**, *7*, 149–156. [[CrossRef](#)]
17. Rome, L.C.; Sosnicki, A.A. The influence of temperature on mechanics of red muscle in carp. *J. Physiol.* **1990**, *427*, 151–169. [[CrossRef](#)] [[PubMed](#)]
18. Wintzer, A.P.; Motta, P.J. The effects of temperature on prey-capture kinematics of the bluegill (*Lepomis macrochirus*): Implications for feeding studies. *Can. J. Zool.* **2004**, *82*, 794–799. [[CrossRef](#)]
19. DeVries, M.S.; Wainwright, P.C. The effects of acute temperature change on prey capture kinematics in largemouth bass, *Micropterus salmoides*. *Copeia* **2006**, *3*, 437–444. [[CrossRef](#)]
20. Cossins, A.R.; Bowler, K. *Temperature Biology of Animals*; Chapman and Hall (Methuen): New York, NY, USA, 1987.
21. Clarke, A.; Johnston, N.M. Scaling of metabolic rate with body mass and temperature in teleost fish. *J. Anim. Ecol.* **1999**, *68*, 893–905. [[CrossRef](#)]
22. Gillooly, J.F.; Brown, J.H.; West, G.B.; Savage, V.M. Effects of size and temperature on metabolic rate. *Science* **2001**, *293*, 2248–2251. [[CrossRef](#)] [[PubMed](#)]
23. Josephson, R.K. Contraction dynamics and power output of skeletal muscle. *Annu. Rev. Physiol.* **1993**, *55*, 527–546. [[CrossRef](#)] [[PubMed](#)]
24. Watabe, S. Temperature plasticity of contractile proteins in fish muscle. *J. Exp. Biol.* **2002**, *205*, 2231–2236. [[PubMed](#)]

25. Malek, R.L.; Sajadi, H.; Abraham, J.; Grundy, M.A.; Gerhard, G.S. The effects of temperature reduction on gene expression and oxidative stress in skeletal muscle from adult zebrafish. *Comp. Biochem. Physiol. Part C: Toxicol. Pharmacol.* **2004**, *138*, 363–373. [[CrossRef](#)] [[PubMed](#)]
26. Johnston, I.A.; Temple, G.K. Thermal plasticity of skeletal muscle phenotype in ectothermic vertebrates and its significance for locomotory behavior. *J. Exp. Biol.* **2002**, *205*, 2305–2322. [[PubMed](#)]
27. Crawshaw, L.I.; Ackerman, R.A.; White, F.N.; Heath, M.E. Metabolic and acid-base changes during selection of warmer water by cold-acclimated fish. *Am. J. Physiol.* **1982**, *242*, R157–R161. [[PubMed](#)]
28. Lemons, D.E.; Crawshaw, L.I. Behavioral and metabolic adjustments to low temperatures in largemouth bass (*Micropterus salmoides*). *Physiol. Zool.* **1985**, *58*, 175–180. [[CrossRef](#)]
29. Clark, D.S.; Green, J.M. Seasonal variation in temperature preference of juvenile Atlantic cod (*Gadus morhua*), with evidence supporting an energetic basis for their diel vertical migration. *Can. J. Zool.* **1991**, *69*, 1302–1307. [[CrossRef](#)]
30. Rome, L.C.; Swank, D.M.; Coughlin, D.J. The influence of temperature on power production during swimming. II. Mechanics of red muscle fibers in vivo. *J. Exp. Biol.* **2000**, *203*, 333–345. [[PubMed](#)]
31. Herbing, I. Effects of temperature on larval fish swimming performance: The importance of physics to physiology. *J. Fish Biol.* **2002**, *61*, 865–876. [[CrossRef](#)]
32. Lee, C.G.; Farrell, A.P.; Lotto, A.; MacNutt, M.J.; Hinch, S.G.; Healey, M.C. The effect of temperature on swimming performance and oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and coho (*O. kisutch*) salmon stocks. *J. Exp. Biol.* **2003**, *206*, 3239–3251. [[CrossRef](#)] [[PubMed](#)]
33. Green, B.S.; Fisher, R. Temperature influences swimming speed, growth and larval duration in coral reef fish larvae. *J. Exp. Mar. Biol. Ecol.* **2004**, *299*, 115–132. [[CrossRef](#)]
34. Sloan, T.J.; Turingan, R.G. Invariant feeding kinematics of two trophically distinct invasive Florida fishes, *Belonesox belizanus* and *Cichlasoma urophthalmus* across environmental temperature regimes. *Int. J. Biol.* **2012**, *4*, 117–126. [[CrossRef](#)]
35. Turingan, R.G.; Sloan, T.J. Modeling the relationship between environmental temperature and feeding performance in Florida (USA) nonnative fishes, with implications for invasive-species response to climate change. *Annu. Rev. Res. Biol.* **2014**, *4*, 121–132. [[CrossRef](#)]
36. Belshe, J.F. *Observations of an Introduced Tropical Fish (Belonesox belizanus) in Southern Florida*; University of Miami: Coral Gables, FL, USA, 1961.
37. Miley, W.W. Ecological Impact of the Pike Killifish, *Belonesox belizanus*, Kner, (Poeciliidae) in Southern Florida. Master Thesis, Florida Atlantic University, Boca Raton, FL, USA, 1978.
38. Anderson, R. Geographic variation and aspects of the life history of *Belonesox belizanus* Kner (Pisces: Poeciliidae) from Central America. Master Thesis, University of Central Florida, Orlando, FL, USA, 1980.
39. Kerfoot, J.R.; Lorenz, J.J.; Turingan, R.G. Environmental correlates of the abundance and distribution of *Belonesox belizanus* in a novel environment. *Environ. Biol. Fish* **2011**, *92*, 125–139. [[CrossRef](#)]
40. Florida Wildlife Commission. Available online: <http://myfwc.com/> (accessed on 15 December 2012).
41. National Oceanic and Atmospheric Administration. National Oceanographic Data Center. Available online: <http://www.nodc.noaa.gov/dsdt/cwtg/all.html> (accessed on 15 December 2012).
42. Hubbs, C. Fishes of the Yucatan Peninsula. *Carnegie Inst. Wash. Publ.* **1936**, *457*, 157–287.
43. Rosen, D.E.; Bailey, R.M. The poeciliid fishes (Cyprinodontiformes): Their structure, zoogeography, and systematics. *Bull. Am. Mus. Nat. Hist.* **1963**, *126*, 1–146.
44. Greven, H.; Brenner, M. Further notes on dentition and prey capture of the Pike killifish *Belonesox belizanus* (Poeciliidae). *Bull. Fish Biol.* **2008**, *10*, 97–103.
45. Ferry-Graham, L.A.; Hernandez, L.P.; Gibb, A.; Pace, C. Unusual kinematics and jaw morphology associated with piscivory in the poeciliid, *Belonesox belizanus*. *Zoology* **2010**, *113*, 140–147. [[CrossRef](#)] [[PubMed](#)]
46. Harms, C.A.; Turingan, R.G. Dietary flexibility despite behavioral stereotypy contributes to successful invasion of the pike killifish, *Belonesox belizanus*, in Florida, USA. *Aquat. Invasions* **2012**, *7*, 547–553. [[CrossRef](#)]
47. FishBase Species Database. Available online: [www.FISHBASE.org](http://www.FISHBASE.org) (accessed on 15 December 2012).
48. Whitfield, P.E.; Hare, J.A.; David, A.W.; Harter, S.L.; Munoz, R.C.; Addison, C.M. Abundance estimates of the Indo-Pacific lionfish *Pterois volitans/miles* complex in the western North Atlantic. *Biol. Invasions* **2007**, *9*, 53–64. [[CrossRef](#)]
49. Hamner, R.M.; Freshwater, D.W.; Whitfield, P.E. Mitochondrial cytochrome *b* analysis reveals two invasive lionfish species with strong founder effects in the western Atlantic. *J. Fish Biol.* **2007**, *71*, 214–222. [[CrossRef](#)]

50. Morris, J.A.; Akins, J.L. Feeding ecology of invasive lionfish (*Pterois volitans*) in the Bahamian archipelago. *Environ. Biol. Fishes* **2009**, *86*, 389–398. [[CrossRef](#)]
51. Morris, J.A. The biology and ecology of Indo-Pacific lionfish. Dissertation, North Carolina State University, Raleigh, NC, USA, 2009.
52. Albins, M.A.; Hixon, M.A. Invasive Indo-Pacific lionfish *Pterois volitans* reduce recruitment of Atlantic coral-reef fishes. *Mar. Ecol. Prog. Ser.* **2008**, *367*, 233–238. [[CrossRef](#)]
53. Pfeiffenberger, J.A. Modulation and Scaling of Prey Capture Kinematics Through Ontogeny in Invasive Indo-Pacific Lionfish, *Pterois volitans/miles complex*. Ph.D. Thesis, Florida Institute of Technology, Melbourne, FL, USA, 2012.
54. Miller, R.R. Geographical distribution of Central American freshwater fishes. *Copeia* **1966**, *4*, 773–802. [[CrossRef](#)]
55. Martinez-Palacios, C.A.; Ross, L.G.; Rosado-Vallado, M. The effects of salinity on the survival and growth of juvenile *Cichlasoma urophthalmus*. *Aquaculture* **1990**, *91*, 65–75. [[CrossRef](#)]
56. Stauffer, J.R.; Boltz, S.E. Effect of salinity on the temperature preference and tolerance of age-0 Mayan cichlids. *Trans. Am. Fish. Soc.* **1994**, *123*, 101–107. [[CrossRef](#)]
57. Schofield, P.J.; Loftus, W.F.; Fontaine, J.A. Salinity effects on behavioural response to hypoxia in the non-native Mayan cichlid *Cichlasoma urophthalmus* from Florida Everglades wetlands. *J. Fish Biol.* **2009**, *7*, 149–156.
58. Martinez-Palacios, C.A.; Ross, L.G. The feeding ecology of Central American cichlid *Cichlasoma urophthalmus* (Gunther). *J. Fish Biol.* **1988**, *33*, 665–670. [[CrossRef](#)]
59. Bergmann, G.T.; Motta, P.J. Diet and morphology through ontogeny of the nonindigenous Mayan cichlid “*Cichlasoma (Nandopsis)*” *urophthalmus* (Gunther 1862) in southern Florida. *Environ. Biol. Fishes* **2005**, *72*, 205–211. [[CrossRef](#)]
60. Chavez-Lopez, R.; Peterson, M.S.; Brown-Peterson, N.; Morales-Gomez, A.A.; Franco-Lopez, J. Ecology of the Mayan cichlid, *Cichlasoma urophthalmus*, in the Alvarado Lagoonal system, Veracruz, Mexico. *Gulf Caribb. Res.* **2005**, *17*, 123–131. [[CrossRef](#)]
61. Hellig, C.J.; Kerschbaumer, M.; Sefc, K.M.; Koblmüller, S. Allometric shape change of the lower pharyngeal jaw correlates with a dietary shift to piscivory in a cichlid fish. *Naturwissenschaften* **2010**, *97*, 663–672. [[CrossRef](#)] [[PubMed](#)]
62. Hulsey, C.D. Function of a key morphological innovation: Fusion of the cichlid pharyngeal jaw. *Proc. R. Soc. B* **2006**, *273*, 669–675. [[CrossRef](#)] [[PubMed](#)]
63. Schmidt-Nielsen, K. *Animal Physiology: Adaptation and Environment*; Cambridge University Press: Cambridge, UK, 1997.
64. Invasive Species and Climate Change. Available online: [www.invasivespecies.gov](http://www.invasivespecies.gov) (accessed on 15 December 2012).
65. Rahel, F.J.; Olden, J.D. Assessing the effects of climate change on aquatic invasive species. *Conserv. Biol.* **2008**, *22*, 521–533. [[CrossRef](#)] [[PubMed](#)]
66. Crowder, L.B. Character displacement and habitat shift in a native cisco in Southeastern Lake Michigan: Evidence for competition? *Copeia* **1984**, *4*, 878–883. [[CrossRef](#)]
67. Douglas, M.E.; Marsh, P.C.; Minckley, W.L. Indigenous fishes of western North America and the hypothesis of competitive displacement: *Medafulgida* (Cyprinidae) as a case study. *Copeia* **1994**, *1*, 9–19. [[CrossRef](#)]
68. Gozlan, R.E.; St-Hilaire, S.; Feist, S.W.; Martin, P.; Kent, M.L. An emergent infectious disease threatens European fish biodiversity. *Nature* **2005**. [[CrossRef](#)] [[PubMed](#)]
69. Ogutu-Ohwayo, R. The decline of the native fishes of lakes Victoria and Kyoga (East Africa) and the impact of introduced species, especially the Nile perch, *Lates niloticus*, and the Nile tilapia, *Oreochromis niloticus*. *Environ. Biol. Fish* **1990**, *27*, 81–96. [[CrossRef](#)]
70. Mooney, H.A.; Cleland, E.E. The evolutionary impact of invasive species. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 5446–5451. [[CrossRef](#)] [[PubMed](#)]
71. Perry, W.L.; Lodge, D.M.; Feder, J.L. Importance of hybridization between indigenous and nonindigenous freshwater species: An overlooked threat to North American biodiversity. *Syst. Biol.* **2002**, *51*, 255–275. [[CrossRef](#)] [[PubMed](#)]
72. Singh, A.K.; Pathak, A.K.; Lakra, W.S. Invasion of an exotic fish—Common carp, *Cyprinus carpio* L. in the Ganga River, India and its impacts. *Acta Ichthyol. Piscat.* **2010**, *40*, 11–19. [[CrossRef](#)]

73. Sato, M.; Kawaguchi, Y.; Nakajima, J.; Mukai, T.; Shimatani, Y.; Onikura, N. A review of the research on introduced freshwater fishes: New perspectives, the need for research, and management implications. *Landsc. Ecol. Eng.* **2010**, *6*, 99–108. [[CrossRef](#)]
74. McKinney, M.L.; Lockwood, J.L. Biotic homogenization: A few winners replacing many losers in the next mass extinction. *Trends Ecol. Evol.* **1999**, *14*, 450–453. [[CrossRef](#)]
75. Bachelier, N.M.; Neal, J.W.; Noble, R.L. Diet overlap between native bigmouth sleepers (*Gobiomorus dormitory*) and introduced predatory fishes in a Puerto Rico reservoir. *Ecol. Freshwater Fish* **2004**, *13*, 111–118. [[CrossRef](#)]
76. Rahel, F.J. Biogeographic barriers, connectivity and homogenization of freshwater faunas: It's a small world after all. *Freshwater Biol.* **2007**, *52*, 696–710. [[CrossRef](#)]
77. Weinstein, M.R.; Litt, M.; Kertesz, D.A.; Wyper, P.; Rose, D.; Coulter, M.; McGreer, A.; Facklam, R.; Ostach, C.; Willey, B.M.; et al. Invasive infections due to a fish pathogen, *Streptococcus iniae*. *S. iniae* study group. *N. Engl. J. Med.* **1997**, *337*, 589–594. [[CrossRef](#)] [[PubMed](#)]
78. Britton, J.R.; Davies, G.D.; Brazier, M.; Pinder, A.C. A case study on the population ecology of a topmouth gudgeon (*Pseudorasbora parva*) population in the UK and the implications for native fish communities. *Aquat. Conserv.* **2006**, *17*, 749–759. [[CrossRef](#)]
79. Hochachka, P.W.; Somero, G.N. *Biochemical Adaptation: Mechanisms and Processes in Physiological Evolution*; Oxford University Press: Oxford, UK, 2002.
80. Adams, S.M.; McLean, R.B.; Parrotta, J.A. Energy partitioning in largemouth bass under conditions of seasonality of seasonally fluctuating prey availability. *Trans. Am. Fish. Soc.* **1982**, *111*, 549–558. [[CrossRef](#)]
81. Cochran, P.A.; Adelman, I.R. Seasonal aspects of daily ration and diet of largemouth bass, *Micropterus salmoides*, with an evaluation of gastric evacuation rates. *Environ. Biol. Fishes* **1982**, *7*, 265–275. [[CrossRef](#)]
82. Garcia-Berthou, E.; Moreno-Amich, R. Food of introduced pumpkinseed sunfish: Ontogenetic diet shift and seasonal variation. *J. Fish Biol.* **2000**, *57*, 29–40. [[CrossRef](#)]
83. Liem, K.F. Modulatory multiplicity in the functional repertoire of the feeding mechanism in cichlids. I. Piscivores. *J. Morphol.* **1978**, *158*, 323–360. [[CrossRef](#)]
84. Liem, K.F. Adaptive significance of intra- and interspecific differences in the feeding repertoires of cichlid fishes. *Am. Zool.* **1980**, *20*, 295–314. [[CrossRef](#)]
85. Lauder, G.V. Patterns of evolution in the feeding mechanism of actinopterygian fishes. *Am. Zool.* **1982**, *22*, 275–285. [[CrossRef](#)]
86. Wainwright, P.C.; Lauder, G.V. Feeding biology of sunfishes: Patterns of variation in the feeding mechanism. *Zool. J. Linn. Soc.* **1986**, *88*, 217–228. [[CrossRef](#)]
87. Turingan, R.G.; Wainwright, P.C. Morphological and functional bases of durophagy in the queen triggerfish, *Balistes vetula* (Pisces, tetraodontiformes). *J. Morphol.* **1993**, *215*, 101–118. [[CrossRef](#)]
88. Ferry-Graham, L.A.; Lauder, G.V. Aquatic prey capture in ray-finned fishes: A century of progress and new directions. *J. Morphol.* **2001**, *248*, 99–119. [[CrossRef](#)] [[PubMed](#)]
89. Westneat, M.W. Evolution of levers and the linkages in the feeding mechanism of fishes. *Integr. Comp. Biol.* **2004**, *44*, 378–389. [[CrossRef](#)] [[PubMed](#)]
90. Deban, S.M.; Lappin, A.K. Thermal effects on the dynamics and motor control of ballistic prey capture in toads: Maintaining high performance at low temperature. *J. Exp. Biol.* **2011**, *214*, 1333–1346. [[CrossRef](#)] [[PubMed](#)]
91. Deban, S.M.; Richardson, J.C. Cold-blooded snipers: Thermal independence of ballistic tongue projection in the salamander *Hydromantes platycephalus*. *J. Exp. Zool.* **2011**, *315*, 618–630. [[CrossRef](#)] [[PubMed](#)]
92. Navas, C.A.; James, R.S.; Wakeling, J.M.; Kemp, K.M.; Johnston, I.A. An integrative study of the temperature dependence of whole animal and muscle performance during jumping and swimming in the frog *Rana temporaria*. *J. Comp. Physiol. B* **1999**, *169*, 588–596. [[CrossRef](#)] [[PubMed](#)]
93. Christian, K.A.; Tracy, C.R. The effect of thermal environment on the ability of hatchling Galapagos land iguanas to avoid predation during dispersal. *Oecologia* **1981**, *49*, 218–223. [[CrossRef](#)]
94. Bennett, A.F. Thermal dependence of muscle function. *Am. J. Physiol.* **1984**, *247*, R217–R229. [[PubMed](#)]
95. Marsh, R.L.; Bennet, A.F. Thermal dependence of contractile properties of skeletal muscle from the lizard, *Sceleporus occidentalis* with comments on the methods for fitting and comparing force-velocity curves. *J. Exp. Biol.* **1986**, *126*, 63–77. [[PubMed](#)]
96. Sandusky, E.P.; Deban, S.M. Temperature effects on the biomechanics of prey capture in the frog *Rana pipiens*. *J. Exp. Zool.* **2012**, *317A*, 595–607. [[CrossRef](#)] [[PubMed](#)]

97. Dickinson, M.H.; Farley, C.T.; Full, R.J.; Koehl, M.A.R.; Kram, R.; Lehman, S. How animals move: An integrative view. *Science* **2000**, *288*, 100–106. [[CrossRef](#)] [[PubMed](#)]
98. Grimby, L.; Hannerz, J.; Hedman, B. The fatigue and voluntary discharge properties of single motor units in man. *J. Physiol.* **1981**, *316*, 545–554. [[CrossRef](#)] [[PubMed](#)]
99. Aagaard, P.; Simonson, E.B.; Andersen, J.L.; Magnusson, P.; Dyhre-Poulsen, P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. *J. Appl. Physiol.* **2002**, *93*, 1318–1326. [[CrossRef](#)] [[PubMed](#)]
100. Johnston, I.A.; Davison, W.; Goldspink, G. Energy metabolism of carp swimming muscles. *J. Comp. Physiol.* **1997**, *114*, 203–216. [[CrossRef](#)]
101. Rome, L.C.; Loughna, P.T.; Goldspink, G. Muscle fibre recruitment as a function of swim speed and muscle temperature in carp. *Am. J. Physiol.* **1984**, *247*, R272–R279. [[PubMed](#)]
102. Rome, L.C. Influence of temperature on muscle recruitment and muscle function in vivo. *Am. J. Physiol.* **1990**, *259*, R210–R222. [[PubMed](#)]
103. Peplowski, M.M.; Marsh, R.L. Work and power output in the hindlimb muscles of Cuban tree frogs *Osteopilu septentrionalis* during jumping. *J. Exp. Biol.* **1997**, *200*, 2861–2870. [[PubMed](#)]
104. Bojsen-Moller, J.; Magnusson, S.P.; Rasmussen, L.R.; Kjaer, M.; Aagaard, P. Muscle performance during maximal isometric and dynamic contractions is influenced by the stiffness of the tendinous structures. *J. Appl. Physiol.* **2005**, *99*, 986–994. [[CrossRef](#)] [[PubMed](#)]
105. Konow, N.; Azizi, E.; Roberts, T.J. Muscle power attenuation by tendon during energy dissipation. *Proc. R. Soc. B* **2012**, *279*, 1108–1113. [[CrossRef](#)] [[PubMed](#)]
106. Boutilier, R.G.; Glass, M.L.; Heisler, N. Blood gases, and extracellular/intracellular acid-base status as a function of temperature in the anuran amphibian *Xenopus laevis* and *Bufo marinus*. *J. Exp. Biol.* **1987**, *130*, 13–25.
107. Renaud, J.M.; Stevens, E.D. The extent of short-term and long-term compensation to temperature shown by frog and toad Sartorius muscle. *J. Exp. Biol.* **1984**, *108*, 57–75.
108. Vogel, S. *Life in Moving Fluids, the Physical Biology of Flow*, 2nd ed.; Princeton University Press: Princeton, NJ, USA, 1994; p. 467.



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).