

Relationship of thyroid function with renal hemodynamics and cholesterol metabolism in proteinuric kidney disease: A pilot study

Yoshitaka Iwazu,^{1,2,3} Kazuhiko Kotani,^{1,4} Taro Sugase,⁵ Daisuke Nagata,³

Toshiyuki Yamada¹

¹ Department of Clinical Laboratory Medicine, Jichi Medical University, Shimotsuke, Tochigi 329-0498, Japan

² Division of Anti-ageing Medicine, Center for Molecular Medicine, Jichi Medical University, Shimotsuke, Tochigi 329-0498, Japan

³ Department of Nephrology, Jichi Medical University, Shimotsuke, Tochigi 329-0498, Japan

⁴ Division of Community and Family Medicine, Center for Community Medicine, Jichi Medical University, Shimotsuke, Tochigi 329-0498, Japan

⁵ Seikai Medical Clinic Nasu, Otawara, Tochigi, 324-0063, Japan

Corresponding author

Yoshitaka Iwazu, MD, PhD

Department of Clinical Laboratory Medicine, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke 329-0498, Tochigi, JAPAN

TEL: +81-285-58-7346, FAX: +81-285-44-4869

E-mail: iwazu@jichi.ac.jp




Figure S1. Factors contributing to hypothyroidism in chronic kidney disease, and interrelationship between proteinuric kidney disease, hypothyroidism, and lipids (thyroid kidney cholesterol axis). LDL: low-density lipoprotein.

Table S1. Correlations of filtration fraction and total cholesterol levels with free triiodothyronine (FT3) and thyroxine (FT4).

Variable (units)	Filtration fraction	Total cholesterol
	β (P value)	β (P value)
FT3 (pg/mL)	0.51 (0.02)	-0.06 (0.76)
FT4 (ng/dL)	0.14 (0.48)	-0.45 (0.02)