




Figure S3.

cv. Lomonosovskij	cv. Sudarynja	cv. Gusar
		
<p>PMCs with (a) regular M I and AI; (b) 4 non oriented chromosomes at M1; (c) 2 lagging chromosomes at MI; (d) lagging chromosomes at TI; (e) normal TI; (f) regular MII; (g) 2 non oriented chromosomes at MII; (h) 2 lagging chromosomes at AII; (i) regular TII; (j)) micronucleus formation at TII; (k) tetrads of microspores.</p>	<p>PMCs with (a) regular M I and AI; (b) PMC with 1 non oriented chromosome at M1; (c) 1 precocious chromosome at AI; (d) the normal T1; (e) lagging chromosomes at TI; (f) regular MII; (g) non oriented chromosomes at MII; (h) lagging chromosome at AII; (i) 2 non oriented chromosomes at AII; (j) micronucleus formation at TII; (k) normal TII; (l) tetrads of microspores; (m) tetrad with 1 additional microspores derived from micronuclei.</p>	<p>(a) PMC with regular M I; (b) the group of PMCs with regular MI, AI, TI and abnormal MI with 1 non oriented chromosome; (c) regular TI; (d) PMCs with regular TI and TI with 1 lagging chromosome; (e) TI with 1 non oriented chromosome; (f) regular MII; (g) regular AII; (h-i) regular TII; (j) tetrads of microspores.</p>