

Can a Rotating Black Hole Be Overspun in Seven Dimensions?

Sanjar Shaymatov ^{1,2,3,4} , Bobomurat Ahmedov ^{1,5*}  and Eldor Karimbaev ¹

¹ Institute of Fundamental and Applied Research, National Research University TIIAME, Kori Niyoziy 39, Tashkent 100000, Uzbekistan; sanjar@astrin.uz (S.S.); ekarimboev@gmail.com (E.K.)

² School of Mathematics and Natural Sciences, New Uzbekistan University, Mustaqillik Ave. 54, Tashkent 100007, Uzbekistan

³ Power Engineering Faculty, Tashkent State Technical University, Tashkent 100095, Uzbekistan

⁴ Institute of Engineering Physics, Samarkand State University, University Avenue 15, Samarkand 140104, Uzbekistan

⁵ Physics Faculty, National University of Uzbekistan, Tashkent 100174, Uzbekistan

* Correspondence: ahmedov@astrin.uz

Abstract: Five-dimensional rotating black holes with two rotations could be overspun except for a single rotation, whereas a black hole in six dimensions always obeys the weak cosmic censorship conjecture (WCCC) in the weak form even for linear particle accretion. In this paper, we investigate the overspinning of a seven-dimensional rotating black hole with three rotation parameters. It is shown that a black hole in the seven dimensions cannot be similarly overspun, thereby obeying the WCCC even under linear particle accretion. It turns out that a black hole always respects the weak cosmic censorship conjecture in seven dimensions.

Keywords: Myers–Perry black hole; overspinning; weak cosmic censorship conjecture

PACS: 04.50.+h; 04.20.Dw



Citation: Shaymatov, S.; Ahmedov, B.; Karimbaev, E. Can a Rotating Black Hole Be Overspun in Seven Dimensions? *Universe* **2023**, *9*, 190. <https://doi.org/10.3390/universe9040190>

Academic Editor: Lorenzo Iorio

Received: 8 March 2023

Revised: 4 April 2023

Accepted: 10 April 2023

Published: 17 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The discovery of gravitational wave (GW) as two stellar black hole mergers [1,2] through LIGO-VIRGO detection has opened a new stage in black hole astrophysics. GW was expected to be a very powerful tool in revealing hidden properties of black holes. Very recently, the first image of the supermassive black hole of the M87 galaxy was obtained with the collaboration of the Event Horizon Telescope (EHT) [3,4]. In addition, the first image of the black hole candidate and its conceptual aspects [5,6] provides a new way to realize its presence in the universe. There exist, however, unexplored problems associated with black holes. The WCCC remains one of the unanswered questions in general relativity (GR) [7]. According to the WCCC, a black hole is always covered by an event horizon concealing singularity from observers outside. The WCCC has been tested by various tools and processes that allow the transition from a black hole to a naked singularity; thus, it has so far remained an active research area. For the first time, a gedanken experiment was proposed to test the WCCC and whether a black hole turns into a naked singularity [8]. Later, the issue of WCCC violation was approached from a different perspective, i.e., a naked singularity would be formed as a result of gravitational collapse (see, e.g., [9–15]). In GR, the gravitational collapse would play one of the most important roles in the formation of a naked singularity. However, there is no proof of the occurrence of naked singularity yet.

It was shown that a near extremal black hole cannot be turned into an extremal one. This happens because particles with suitable parameters cannot reach the extremal black hole horizon because the parameter space that allows particles to approach the horizon pinches off [8,16]. Later, this question was formulated from a different perspective; that is, could a near extremal black hole be overextremalized to create a naked singularity by impinging particles? This was first addressed by Hubeny [17] who showed that it

was achievable by destroying the horizon. Later, it was also extended to a rotating black hole [18]. It was shown that the rotating black hole can be overspun by plunging particles with suitable parameters. Here, it is worth noting that this experiment for overcharging/overspinning was initiated for a linear order particle accretion by ignoring all higher order effects. Following this thought experiment, an extensive analysis was conducted (see, e.g., [19–23] addressing overcharging/overspinning of the black holes in various gravity models). For this thought experiment, when the self-force and back-reaction effects are taken into account, it is not possible for impinging particles to reach the horizon, i.e., over-extremality cannot be approached, thereby respecting the WCCC [24–28].

So, all extensive analyses so far have been conducted for linear order accretion. Recently, Sorce and Wald [29,30] proposed a new gedanken experiment that includes second-order particle accretion. With this, they opened a new stage of investigation for testing the WCCC. This new gedanken experiment supports the WCCC, thus a black hole cannot be overspun/overcharged, which adds a peculiarity to explore the overspinning/overcharging of black holes. This was then extended to various cases. It turns out that a black hole that could be over-extremalized at the linear order cannot be overspun/overcharged when non-linear effects are involved (see, e.g., [31–38]). This experiment was also tested in Einstein–Born–Infeld and static charged Gauss–Bonnet black holes [39,40]. There was an investigation that suggested that a test magnetic field would serve as a cosmic censor [41]. The same is also true for its backreaction effect [42], i.e., the magnetic field beyond its threshold value would have a similar effect in contrast to the non-linear order effects. Similarly, the cosmological weak magnetic field could potentially be important for testing the WCCC [43].

Recent analysis shows that a five-dimensional rotating black hole has a remarkable feature that it could be overspun when it has two rotations, yet there is no overspinning in a single rotation case even under linear order effects [37]. If one switches off the rotation parameters of the black hole, it could then be overcharged [44]. This led to an interesting question—could a black hole be over-extremalized when it has both charge and spin? There is, however, no available exact solution for an analogue of the Kerr–Newman black hole in five dimensions. The only way to address this question is to consider the minimally gauged supergravity-charged rotating black hole [45], which is regarded as the very closest one to the Kerr–Newman black hole in five dimensions. What emerges here is that the ultimate case depends on which parameter is dominant. It is demonstrated that a black hole with a single rotation cannot be over-extremalized if and only if angular momentum dominates over the black hole charge. Meanwhile, the opposite result is true when the charge is dominant [36].

As discussed above, for the validity of Einstein’s gravity, Penrose proposed that black hole singularity would always be hidden behind a horizon. This is what we refer to as the WCCC. There is, however, no proof of the CCC yet, and thus it remains an open question. With this in view, it is increasingly important to test whether an existing horizon of a rotating black hole could be destroyed by overspinning, thus violating the weak WCCC. Thus, we shall focus here on the overspinning of a near extremal black hole by throwing in test particles with suitable parameters. Recently, it was shown that a black hole in six dimensions cannot be overspun by first-order particle accretion, as well as the same result holds well for a single rotation case [38]. However, a black hole with only one rotation in higher dimension > 5 cannot be overspun as one of no extremality. In this paper, we would like to verify whether the same result holds well in seven dimensions. With this motivation, we analyse seven-dimensional rotating black holes with three rotations. We show that the black hole in $d = 7$ cannot be overspun even under linear order particle accretion, similar to what is observed for six dimensions.

We describe the paper as follows: in Section 2, we present a higher dimension rotating black hole in a general form which is followed by analysis leading to the discussion of the overspinning of a black hole for particular cases in Section 3. We discuss our concluding results in Section 4.

2. Odd and Even Dimensional Rotating Myers–Perry Black Hole Spacetimes

The line element of the rotating Myers–Perry general black hole metric [46] for Einstein gravity in $d = 2n + 1$ and $d = 2n + 2$ dimensions is given by

$$\begin{aligned}
 ds^2 &= -dt^2 + (r^2 + a_i^2) \left(d\mu_i^2 + \mu_i^2 d\phi_i^2 \right) \\
 &+ \frac{\mu r^2}{\Pi F} \left(dt + a_i \mu_i^2 d\phi_i \right) + \frac{\Pi F}{\Pi - \mu r^2} dr^2,
 \end{aligned}
 \tag{1}$$

and

$$\begin{aligned}
 ds^2 &= -dt^2 + r^2 d\alpha^2 + (r^2 + a_i^2) \left(d\mu_i^2 + \mu_i^2 d\phi_i^2 \right) \\
 &+ \frac{\mu r}{\Pi F} \left(dt + a_i \mu_i^2 d\phi_i \right) + \frac{\Pi F}{\Pi - \mu r} dr^2,
 \end{aligned}
 \tag{2}$$

respectively, where

$$\begin{aligned}
 F &= 1 - \frac{a_i^2 \mu_i^2}{r^2 + a_i^2}, \\
 \Pi &= (r^2 + a_1^2) \dots (r^2 + a_n^2).
 \end{aligned}
 \tag{3}$$

Here, a_i and μ , respectively, refer to rotation parameters and the mass parameter. We note here that μ_i and α are related by $\mu_i^2 = 1$ and $\mu_i^2 + \alpha^2 = 1$ for $d = 2n + 1$ and $d = 2n + 2$, respectively, where n is the maximum number of rotation parameter a black hole can have. For odd and even dimensions, we further define Δ as follows:

$$\Delta = \Pi - \mu r^2 \quad \text{and} \quad \Delta = \Pi - \mu r.
 \tag{4}$$

A black hole horizon can be defined by $\Delta = 0$ for odd and even dimensions, respectively. Let us then write the horizon equation for odd $d = 2n + 1$ and even $d = 2n + 2$ dimensions as

$$\begin{aligned}
 r^{2n} &+ f_1(a_i^2)r^{2(n-1)} + f_2(a_i^2)r^{2n-4} + \dots - \mu r^2 \\
 &+ a_1^2 a_2^2 \dots a_n^2 = 0,
 \end{aligned}
 \tag{5}$$

and

$$\begin{aligned}
 r^{2n} &+ f_1(a_i^2)r^{2(n-1)} + f_2(a_i^2)r^{2n-4} + \dots - \mu r \\
 &+ a_1^2 a_2^2 \dots a_n^2 = 0,
 \end{aligned}
 \tag{6}$$

where f_i are functions given as a function of rotation parameters a_i . The sufficient condition for overspinning is that Equations (5) and (6) must be required to have two positive roots to define extremality, i.e., $r_- = r_+$. It is then possible to test whether an existing horizon of a rotating black hole could be destroyed by overspinning. Hence, for an existing extremality for $2n + 1$ we assume that the general solution can be found for the horizon equation as

$$(r - \gamma)^2 (r + \beta_1) (r + \beta_2) (r^2 + \alpha_1^2) \dots (r^2 + \alpha_n^2) = 0,
 \tag{7}$$

where γ has two double roots, β_1 and β_2 are two negative roots, while α_i are complex roots. From Equation (7), it is certain that the term γr^{2n-1} , which never exists in the horizon Equation (5), appears. As a result, $\gamma = 0$ suggests that two positive roots cannot exist, i.e., no extremality condition appears; thus the question of overspinning never arises. Similarly, for $d = 2n + 2$, the same general solution can also be found as

$$(r - \gamma)^2 (r + \beta_1) (r + \beta_2) (r^2 + \alpha_1^2) \dots = 0.
 \tag{8}$$

It is straightforward to show the term γr^{2n-1} in Equation (6). Thus, $\gamma = 0$ always holds – no positive double root. As a consequence of the general solution for $d = 2n + 2$, there appears no extremality condition; hence for the black hole having at least three rotations, overspinning never happens. This is the case for $n = 3$ and any $n > 3$ in $d = 2n + 1$ and $d = 2n + 2$. What happens if one of the rotations vanishes for $d = 2n + 1$ and $d = 2n + 2$? There remains only one positive root for Equations (7) and (8); hence, no question of overspinning for a number of rotations $< n$ arises. To that, we shall further focus on the case $n = 3$.

Let us then consider 3 rotation cases $n = 3$ to check whether the above statement holds well or not. So, in the case of $n = 3$ for $d = 7$, the line element of the rotating black hole metric (1) in the Boyer = –Lindquist coordinates $(t, r, \theta, \varphi, \psi, \chi)$ yields

$$\begin{aligned}
 ds^2 = & -dt^2 + \frac{\mu}{\Sigma} \left(dt - a_1 \sin^2 \theta d\varphi - a_2 \cos^2 \theta \sin^2 \chi d\psi \right. \\
 & \left. - a_3 \cos^2 \theta \cos^2 \chi d\psi \right)^2 + \frac{r^4 \Xi}{\Pi - \mu r^2} dr^2 + \Sigma d\theta^2 \\
 & + (r^2 + a_1^2) \sin^2 \theta d\varphi^2 + (r^2 + a_2^2) \cos^2 \theta \sin^2 \chi d\psi^2 \\
 & + (r^2 + a_3^2) \cos^2 \theta \cos^2 \chi d\psi^2 \\
 & + \left(r^2 + a_2^2 \cos^2 \chi + a_3^2 \sin^2 \chi \right) \cos^2 \theta d\chi^2, \tag{9}
 \end{aligned}$$

where

$$\Sigma = r^2 + a_1^2 \cos^2 \theta + a_2^2 \sin^2 \theta \sin^2 \chi + a_3^2 \sin^2 \theta \cos^2 \chi. \tag{10}$$

Here, a_i and μ are given by

$$\mu = \frac{16M}{5\pi^2}, \quad a_i = \frac{5J_i}{2M}, \tag{11}$$

and the angular coordinates range over, $\theta \in [0, \pi/2]$ and $\varphi, \psi, \chi \in [0, 2\pi]$.

3. Overspinning of Seven-Dimensional Rotating Black under a Linear Particle Accretion

3.1. Black Hole with Maximum Three Rotations A_1, A_2 and A_3 in Seven Dimensions

Here, we start to consider a black hole with three rotations in $d = 7$. It is shown that no extremality condition exists. This is a remarkable aspect of rotating black holes with three rotations. For given $n = 3$, we recall Equation (5), which solves to give six roots, i.e., four complex, one positive and one negative root. As discussed previously, an extremality requires two positive roots. Since only one positive root exists, it does not satisfy the extremality condition; therefore, no overspinning occurs for a black hole with three rotations. To be more accurate, we analyse this positive root, which is given by

$$\begin{aligned}
 r_+ = & \frac{1}{6} \left[\frac{2^{4/3} B}{\left(\sqrt{A^2 - 4B^3} - A \right)^{1/3}} \right. \\
 & \left. + 2^{2/3} \left(\sqrt{A^2 - 4B^3} - A \right)^{1/3} - 2 \left(a_1^2 + a_2^2 + a_3^2 \right) \right]^{1/2}, \tag{12}
 \end{aligned}$$

with

$$\begin{aligned}
 A = & 2a_1^6 - 3a_1^4 \left(a_2^2 + a_3^2 \right) - 3a_1^2 \left(a_2^4 - 4a_2^2 a_3^2 + a_3^4 - 3\mu \right) \\
 & + \left(a_2^2 + a_3^2 \right) \left(2a_2^4 - 5a_2^2 a_3^2 + 2a_3^4 + 9\mu \right), \tag{13}
 \end{aligned}$$

$$B = a_1^4 - a_1^2 \left(a_2^2 + a_3^2 \right) + a_2^4 - a_2^2 a_3^2 + a_3^4 + 3\mu. \tag{14}$$

For a black hole with equal rotations, i.e., $a_1 = a_2 = a_3 = a$, Equation (12) yields

$$r_+ = \left[\frac{\left(3\sqrt{3}\mu\sqrt{-(4\mu - 27a^4)} - 27a^2\mu\right)^{1/3}}{3\sqrt[3]{2}} + \frac{\sqrt[3]{2}\mu}{\left(3\sqrt{3}\mu\sqrt{-(4\mu - 27a^4)} - 27a^2\mu\right)^{1/3}} - a^2 \right]^{1/2}. \tag{15}$$

For an existing black hole horizon, $4\mu > 27a^4$ must be satisfied always. The above equation becomes a complex quantity, i.e., $r_+^2 < 0$, thus resulting in no extremality condition existing. Hence, there never arises a question of overspinning a black hole in the case of three rotations. We then further consider the possible cases to test whether an extremality condition exists.

- $A^2 - 4B^3 = 0$ which defines an extremal black hole:

For that, Equation (12) takes the following form as

$$r_+^2 = \frac{1}{6} \left[-4B^{1/2} - 2(a_1^2 + a_2^2 + a_3^2) \right]. \tag{16}$$

This clearly shows that the above equation is always negative, and thus no extremality condition, i.e., $r_+^2 < 0$ that causes no black hole horizon, exists.

- Near extremal black hole:

We assume that a near extremality condition is well defined by the following condition

$$4B^3 = A^2(1 - \epsilon^2) \tag{17}$$

with small $\epsilon \ll 0$. Recalling Equation (12), the horizon r_+ for a near extremal black hole is given by

$$r_+^2 = \frac{1}{6} \left[\frac{2^{4/3}A^{-1/3}B}{(\epsilon - 1)^{1/3}} + 2^{2/3}(\epsilon - 1)^{1/3}A^{1/3} - 2(a_1^2 + a_2^2 + a_3^2) \right]. \tag{18}$$

Since $\epsilon - 1 < 0$, we obtain $r_+^2 < 0$ that is always satisfied for a near extremal black hole, thereby resulting in no extremality condition existing. In this respect, overspinning simply loses its applicability. With this in view, here we intend to note that a black hole with three rotation parameters in $d = 7$ cannot be overspun as no extremality condition exists; therefore, the WCCC is always respected. Next, we explore black holes with two and single rotation cases.

3.2. Black Hole with Two Rotations A_1 and A_2 In Seven Dimensions

Here, we consider a black hole with two rotations in $d = 7$ to understand more deeply whether it favours the cosmic censorship conjecture. By recalling Equation (5), we first write the black hole horizon as follows:

$$r_+ = \left(\mu^{1/2} - a_2^2 \right)^{1/2}. \tag{19}$$

If one considers the mass parameter μ in terms of black hole mass M , it then yields

$$r_+ = \left[\left(\frac{16M}{5\pi^2} \right)^{1/2} - a^2 \right]^{1/2}. \tag{20}$$

From the above equation, the condition $\mu^{1/2} < a^2$ refers to an object without a horizon. However, we begin a nearly extremal black hole, according to which rotation parameters are regarded as $a_1 = a_2 = \sqrt[4]{\frac{16M}{5\pi^2}} (1 - \epsilon^2)$ with $\epsilon \ll 1$.

Here, we assume that an impinging particle has equal rotations $\delta J_\varphi = \delta J_\phi$ so that it would add an equal amount to the black hole rotations [8,18,37].

Equations (19) and (20) define the minimum threshold value as

$$\sqrt[4]{\frac{16}{5\pi^2}} (M + \delta E)^{1/4} < \frac{5}{2} \left(\frac{J + \delta J}{M + \delta E} \right), \tag{21}$$

for which the minimum threshold value for either φ or ϕ rotation takes the following form

$$\begin{aligned} \delta J_{min} = & \sqrt[4]{\frac{2^8}{5^5\pi^2}} \left(M^{5/4}\epsilon^2 + \frac{5}{4} M^{1/4}\delta E \right. \\ & \left. + \frac{5}{32} M^{-3/4}\delta E^2 \right). \end{aligned} \tag{22}$$

Since an impinging particle adds an equal amount to both rotations, the total amount due to both δJ_φ and δJ_ϕ is written as

$$\begin{aligned} \delta J_{min}^{total} = & \sqrt[4]{\frac{2^8}{5^5\pi^2}} \left(2M^{5/4}\epsilon^2 + \frac{5}{2} M^{1/4}\delta E \right. \\ & \left. + \frac{5}{16} M^{-3/4}\delta E^2 \right). \end{aligned} \tag{23}$$

This is the lower threshold value of angular momentum required for the impinging particle to fall into the black hole.

For a particle to fall into the black hole with minimum threshold, it must first reach the horizon. For that, we need to define the upper bound of the angular momentum. For the impinging particle to reach the horizon, it should have enough energy, which is in general given by

$$\delta E \geq \Omega_+^{(\varphi)} \delta J_\varphi + \Omega_+^{(\phi)} \delta J_\phi, \tag{24}$$

with angular velocities $\Omega_+^{(\varphi,\phi)}$ evaluated at the horizon. Hence, the upper threshold value of angular momentum can be written as

$$\delta J_{max}^{total} = \frac{r_+^2 + a^2}{a} \delta E. \tag{25}$$

This is what is called the upper threshold value of the angular momentum for impinging particles falling into the black hole having equal rotations $a_1 = a_2 = a$. Hence, we have

$$\delta J_{max} = \frac{5}{2} \sqrt[4]{\frac{2^8}{5^5\pi^2}} (1 + \epsilon^2) M^{1/4} \delta E. \tag{26}$$

For an impinging particle to cross the horizon, the condition $\delta J_{max} > \delta J_{min}$ must always be satisfied. If not, no parameter space, which cannot allow particles to cross the horizon and

fall into the black hole, appears. To that, we analyse the parameter space, i.e., ΔJ , and it is given by

$$\Delta J = \left(M^{1/4} \epsilon^2 \delta E - \frac{4}{5} M^{5/4} \epsilon^2 - \frac{1}{8} M^{-3/4} \delta E^2 \right) \times \left(\frac{16}{5\pi^2} \right)^{1/4}. \tag{27}$$

This clearly shows that the second and the third terms are of second order in ϵ , while the first term is of third order in ϵ . Thus, the second and the third terms dominate over the first term, resulting in indicating $\Delta J < 0$ always. With this, one can deduce that no parameter space allowing particles to overspin the black hole appears. Thus, the WCCC is always respected for a black hole with two rotations in seven dimensions.

3.3. Black Hole Having Only Single Rotation A_1 in Seven Dimensions

For single rotation, Equation (5) gives the following form for event horizon

$$r_+ = \left(\frac{\sqrt{a_1^4 + 4\mu} - a_1^2}{2} \right)^{1/2} \tag{28}$$

with the presence of no extremality condition, thereby without overspinning. This states that a black hole with a single rotation in seven dimensions cannot be overspun, similarly to what is observed in the work [38]. One can then conclude that no extremality condition for a black hole with single rotation in all higher dimensions exists except for five-dimensional rotating black holes. However, the black hole with a single rotation cannot be overspun even if it has an extremality condition [37].

We have explicitly shown that a black hole with three rotations cannot be overspun for linear particle accretion, resulting in supporting the WCC. Thus, no overspinning occurs. To better understand its dynamics, we further consider the effective gravitational potential for seven dimensions. Following Equation (4), one can obtain the effective gravitational potential for black holes having n rotations as

$$\Phi(r) \approx \frac{\Delta}{r^2} - 1 = \frac{(r^2 + a^2) \dots (r^2 + a_n^2)}{r^{2n}} - \frac{\mu}{r^{D-3}} - 1. \tag{29}$$

The above equation for seven dimensions can be explicitly written as

$$\Phi(r) = -\frac{\mu}{r^4} + \frac{a_1^2 + a_2^2 + a_3^2}{r^2} + \frac{a_1^2 a_2^2 + a_2^2 a_3^2 + a_1^2 a_3^2}{r^4} + \frac{a_1^2 a_2^2 a_3^2}{r^6}, \tag{30}$$

where a_i refers to black hole rotation parameters. In Figure 1 we demonstrate the radial profile of the effective gravitational potential $\Phi(r)$ and its first derivative. As can be seen from Figure 1, at large distances r/M , the resultant acceleration becomes repulsive for all seven-dimensional black holes, thus resulting in allowing particles not to reach the black hole horizon. In addition, this plot clearly shows that the resultant acceleration behaves attractively very near the black hole horizon due to the fact $r_h < 1$ is always satisfied. It is vital to note that the first attractive term that stems from mass in Equation (30) dominates the repulsive second term $1/r^2$. Interestingly, one can also observe the same for $\partial\Phi(r)/\partial r$. It is vital to note here that we have shown that six-dimensional black holes having a maximum of two rotations cannot be overspun for linear particle accretion (see for example [38]). Interestingly, it turns out that as shown above dynamics would be similar for both higher six and seven dimensions, thereby leading to a similar result that black holes cannot be overspun. Therefore, the same would be the case for non-linear particle accretions that always endorse weak cosmic censorship conjecture.




Figure 1. The radial profile of the gravitational effective potential $\Phi(r)$ and its first derivative $\partial\Phi(r)/\partial r$ for $D = 7$ are plotted for various combinations of black hole rotating parameters. Note that in both panels, vertical line refers to the horizon for $a_1 = a_2 = a_3 = 0.6204$ corresponding to a nearly extremal black hole.

4. Conclusions

In this paper, we studied the validity of the WCCC for a black hole with three rotations in dimension $d = 7$ for a linear order particle accretion. It is well-known that a black hole with two rotations could be overspun except for a single rotation, i.e., a black hole with only a single rotation in all higher dimensions $d > 4$ cannot be overspun, thereby favouring the WCCC [37]. Hence, the natural question then arises: what happens with the black hole in dimension $d = 6$? It turns out that black holes cannot be overspun all through [38]. With this motivation, we intended to test whether the same result holds well for the black hole in dimension $d = 7$.

We have shown that no extremal condition exists for single rotation in seven dimensions, as was expected; thus, it leads to no question for its overspinning. For two and three rotations, a black hole cannot similarly be overspun for a linear order particle accretion, thereby obeying the WCCC in dimension $d = 7$. To be more accurate, we have analysed the dynamics of overspinning by adapting the effective gravitational potential for seven dimensions $d = 7$. We have shown that the resultant gravity becomes repulsive all through at larger r so that particles cannot reach the black hole horizon. One can then infer that the dynamics would be similar for higher six and seven dimensions, yielding a similar result for which a black hole cannot be overspun; thus, no violation of the WCCC occurs. With this, we verified a remarkable result that black holes in these $d \geq 6$ higher dimensions cannot be overspun even under linear particle accretion [47]. This is a remarkable aspect of black holes in higher dimensions.

Author Contributions: Conceptualization, S.S.; methodology, S.S. and B.A.; software, S.S.; validation, S.S. and E.K.; formal analysis, S.S.; investigation, S.S. and B.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Acknowledgments: S.S. and B.A. wish to acknowledge the support from Research F-FA-2021-432 of the Uzbekistan Agency for Innovative Development.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Abbott, B.P.; Abbott, R.; Abbott, T.D.; Abernathy, M.R.; Acernese, F.; Ackley, K.; Adams, C.; Adams, T.; Addesso, P.; Adhikari, R.X.; et al. Observation of Gravitational Waves from a Binary Black Hole Merger. *Phys. Rev. Lett.* **2016**, *116*, 061102.
2. Abbott, B.P.; Abbott, R.; Abbott, T.D.; Abernathy, M.R.; Acernese, F.; Ackley, K.; Adams, C.; Adams, T.; Addesso, P.; Adhikari, R.X.; et al. Properties of the Binary Black Hole Merger GW150914. *Phys. Rev. Lett.* **2016**, *116*, 241102.
3. Event Horizon Telescope Collaboration. First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole. *Astrophys. J.* **2019**, *875*, L1.

4. Event Horizon Telescope Collaboration. First M87 Event Horizon Telescope Results. VI. The Shadow and Mass of the Central Black Hole. *Astrophys. J.* **2019**, *875*, L6.
5. Eckart, A.; Hüttemann, A.; Kiefer, C.; Britzen, S.; Zajaček, M.; Lämmerzahl, C.; Stöckler, M.; Valencia-S, M.; Karas, V.; García-Marín, M. The Milky Way's Supermassive Black Hole: How Good a Case Is It? *Found. Phys.* **2017**, *47*, 553–624.
6. Zajaček, M.; Tursunov, A.; Eckart, A.; Britzen, S. On the charge of the Galactic centre black hole. *Mon. Not. R. Astron. Soc.* **2018**, *480*, 4408–4423.
7. Penrose, R. Gravitational Collapse: The Role of General Relativity. *Riv. Nuovo Cimento* **1969**, *1*, 252.
8. Wald, R. Gedanken experiments to destroy a black hole. *Ann. Phys.* **1974**, *82*, 548–556. [[CrossRef](#)]
9. Christodoulou, D. Gravitational collapse. *Ann. N. Y. Acad. Sci.* **1986**, *470*, 147–155. [[CrossRef](#)]
10. Joshi, P.S. *Global Aspects in Gravitation and Cosmology*; Clarendon Press: Oxford, UK, 1993.
11. Joshi, P.S. Gravitational collapse: The story so far. *Pramana* **2000**, *55*, 529.
12. Goswami, R.; Joshi, P.S.; Singh, P. Quantum Evaporation of a Naked Singularity. *Phys. Rev. Lett.* **2006**, *96*, 031302.
13. Harada, T.; Iguchi, H.; Nakao, K. Physical Processes in Naked Singularity Formation. *Prog. Theor. Phys.* **2002**, *107*, 449–524.
14. Vieira, R.S.S.; Schee, J.; Kluźniak, W.; Stuchlík, Z.; Abramowicz, M. Circular geodesics of naked singularities in the Kehagias-Sfetsos metric of Hořava's gravity. *Phys. Rev. D* **2014**, *90*, 024035.
15. Giacomazzo, B.; Rezzolla, L.; Stergioulas, N. Collapse of differentially rotating neutron stars and cosmic censorship. *Phys. Rev. D* **2011**, *84*, 024022.
16. Dadhich, N.; Narayan, K. On the third law of black hole dynamics. *Phys. Lett. A* **1997**, *231*, 335–338. [[CrossRef](#)]
17. Hubeny, V.E. Overcharging a black hole and cosmic censorship. *Phys. Rev. D* **1999**, *59*, 064013. [[CrossRef](#)]
18. Jacobson, T.; Sotiriou, T.P. Overspinning a Black Hole with a Test Body. *Phys. Rev. Lett.* **2009**, *103*, 141101. [[CrossRef](#)]
19. Saa, A.; Santarelli, R. Destroying a near-extremal Kerr-Newman black hole. *Phys. Rev. D* **2011**, *84*, 027501. [[CrossRef](#)]
20. Bouhmadi-López, M.; Cardoso, V.; Nerozzi, A.; Rocha, J.V. Black holes die hard: Can one spin up a black hole past extremality? *Phys. Rev. D* **2010**, *81*, 084051.
21. Rocha, J.V.; Santarelli, R. Flowing along the edge: Spinning up black holes in AdS spacetimes with test particles. *Phys. Rev. D* **2014**, *89*, 064065.
22. Yang, S.J.; Chen, J.; Wan, J.J.; Wei, S.W.; Liu, Y.X. Weak cosmic censorship conjecture for a Kerr-Taub-NUT black hole with a test scalar field and particle. *Phys. Rev. D* **2020**, *101*, 064048.
23. Gwak, B. Weak cosmic censorship in Kerr-Sen black hole under charged scalar field. *J. Cosmol. Astropart. Phys.* **2020**, *2020*, 058.
24. Barausse, E.; Cardoso, V.; Khanna, G. Test Bodies and Naked Singularities: Is the Self-Force the Cosmic Censor? *Phys. Rev. Lett.* **2010**, *105*, 261102.
25. Zimmerman, P.; Vega, I.; Poisson, E.; Haas, R. Self-force as a cosmic censor. *Phys. Rev. D* **2013**, *87*, 041501. [[CrossRef](#)]
26. Rocha, J.V.; Cardoso, V. Gravitational perturbation of the BTZ black hole induced by test particles and weak cosmic censorship in AdS spacetime. *Phys. Rev. D* **2011**, *83*, 104037.
27. Isoyama, S.; Sago, N.; Tanaka, T. Cosmic censorship in overcharging a Reissner-Nordström black hole via charged particle absorption. *Phys. Rev. D* **2011**, *84*, 124024.
28. Colleoni, M.; Barack, L. Overspinning a Kerr black hole: The effect of the self-force. *Phys. Rev. D* **2015**, *91*, 104024.
29. Sorce, J.; Wald, R.M. Gedanken experiments to destroy a black hole. II. Kerr-Newman black holes cannot be overcharged or overspun. *Phys. Rev. D* **2017**, *96*, 104014.
30. Wald, R.M. Kerr-Newman black holes cannot be over-charged or over-spun. *Int. J. Mod. Phys. D* **2018**, *27*, 1843003. [[CrossRef](#)]
31. An, J.; Shan, J.; Zhang, H.; Zhao, S. Five-dimensional Myers-Perry black holes cannot be overspun in gedanken experiments. *Phys. Rev. D* **2018**, *97*, 104007.
32. Gwak, B. Weak cosmic censorship conjecture in Kerr-(anti)-de Sitter black hole with scalar field. *J. High Energy Phys.* **2018**, *9*, 81.
33. Ge, B.; Mo, Y.; Zhao, S.; Zheng, J. Higher-dimensional charged black holes cannot be over-charged by gedanken experiments. *Phys. Lett. B* **2018**, *783*, 440–445.
34. Ning, B.; Chen, B.; Lin, F.L. Gedanken experiments to destroy a BTZ black hole. *Phys. Rev. D* **2019**, *100*, 044043.
35. Jiang, J. Static charged Gauss-Bonnet black holes cannot be overcharged by the new version of gedanken experiments. *Phys. Lett. B* **2020**, *804*, 135365. [[CrossRef](#)]
36. Shaymatov, S.; Dadhich, N.; Ahmedov, B.; Jamil, M. Five-dimensional charged rotating minimally gauged supergravity black hole cannot be over-spun and/or over-charged in non-linear accretion. *Eur. Phys. J. C* **2020**, *80*, 481.
37. Shaymatov, S.; Dadhich, N.; Ahmedov, B. The higher dimensional Myers-Perry black hole with single rotation always obeys the Cosmic Censorship Conjecture. *Eur. Phys. J. C* **2019**, *79*, 585.
38. Shaymatov, S.; Dadhich, N.; Ahmedov, B. Six-dimensional Myers-Perry rotating black hole cannot be overspun. *Phys. Rev. D* **2020**, *101*, 044028.
39. He, Y.L.; Jiang, J. Weak cosmic censorship conjecture in Einstein-Born-Infeld black holes. *Phys. Rev. D* **2019**, *100*, 124060.
40. Jiang, J.; Zhang, M. Weak cosmic censorship conjecture in Einstein-Maxwell gravity with scalar hair. *Eur. Phys. J. C* **2020**, *80*, 196. [[CrossRef](#)]
41. Shaymatov, S.; Patil, M.; Ahmedov, B.; Joshi, P.S. Destroying a near-extremal Kerr black hole with a charged particle: Can a test magnetic field serve as a cosmic censor? *Phys. Rev. D* **2015**, *91*, 064025.

42. Shaymatov, S.; Ahmedov, B. Overcharging process around a magnetized black hole: Can the backreaction effect of magnetic field restore cosmic censorship conjecture? *Gen. Relativ. Gravit.* **2023**, *55*, 36.
43. Pandey, K.L.; Sethi, S.K.; Ratra, B. Cosmological magnetic braking and the formation of high-redshift, super-massive black holes. *Mon. Not. R. Astron. Soc.* **2019**, *486*, 1629–1640.
44. Revelar, K.S.; Vega, I. Overcharging higher-dimensional black holes with point particles. *Phys. Rev. D* **2017**, *96*, 064010.
45. Chong, Z.W.; Cvetič, M.; Lü, H.; Pope, C.N. General Nonextremal Rotating Black Holes in Minimal Five-Dimensional Gauged Supergravity. *Phys. Rev. Lett.* **2005**, *95*, 161301.
46. Myers, R.C.; Perry, M.J. Black holes in higher dimensional space-times. *Ann. Phys.* **1986**, *172*, 304–347. [[CrossRef](#)]
47. Shaymatov, S.; Dadhich, N. On overspinning of black holes in higher dimensions. *Phys. Dark Universe* **2021**, *31*, 100758.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.