

Supplementary Information


Design of inhibitors of the intrinsically disordered protein NUPR1: balance between drug affinity and target function

Bruno Rizzuti, Wenjun Lan, Patricia Santofimia-Castaño, Zhengwei Zhou, Adrián Velázquez-Campoy, Olga Abián, Ling Peng, José L. Neira, Yi Xia and Juan L. Iovanna

Table of Contents

- Scheme S1: Synthesis of ZZW-115-derived compounds.
- Fig. S1: capacity of ZZW-115-derived compounds to perform nuclear translocation
- Fig. S2: capacity of ZZW-115-derived compounds to sensitize 5-FU induced DNA damage
- ^1H , ^{13}C -NMR spectra of the compounds
- HPLC spectra of the compounds

Scheme S1. Synthesis of ZZW-115-derived compounds.



Name	R	Yield (%)	Name	R	Yield (%)
ZZW-129		68	ZZW-143		69
ZZW-130		79	ZZW-144		63
ZZW-131		74	ZZW-145		76
ZZW-132		71	ZZW-148		70
ZZW-142		76			




Figure S1: Colocalization scatter plots of NUPR1; Pearson's R value (PRV), and Mander's coefficient (MC) were calculated by using the ImageJ Coloc2 plugin. The bar indicates 20 μ m.















Figure S2: Intensity profiles shown for the effect of the ZZW-derived compounds for the ability to sensitize 5-FU induced damage. The bar indicates 20 μ m.






































HPLC analysis of ZZW-130 (Method 1)




HPLC analysis of ZZW-130 (Method 2)





HPLC analysis of ZZW-132 (Method 1)



HPLC analysis of ZZW-132 (Method 2)

