

Article

A Generalized Fejér–Hadamard Inequality for Harmonically Convex Functions via Generalized Fractional Integral Operator and Related Results

Shin Min Kang ^{1,2}, Ghulam Abbas ^{3,4}, Ghulam Farid ⁵ and Waqas Nazeer ^{6,*}

¹ Department of Mathematics and Research Institute of Natural Science, Gyeongsang National University, Jinju 52828, Korea; smkang@gnu.ac.kr

² Center for General Education, China Medical University, Taichung 40402, Taiwan

³ Department of Mathematics, Government College Bhalwal, Sargodha 40100, Pakistan; prof.abbas6581@gmail.com

⁴ Department of Mathematics, University of Sargodha, Sargodha 40100, Pakistan

⁵ Department of Mathematics, COMSATS University Islamabad, Attock Campus 43600, Pakistan; faridphdsms@hotmail.com

⁶ Division of Science and Technology, University of Education, Lahore 54000, Pakistan

* Correspondence: nazeer.waqas@ue.edu.pk; Tel.: +923214707379

Received: 5 June 2018; Accepted: 9 July 2018; Published: 11 July 2018



Abstract: In this paper, we obtain a version of the Fejér–Hadamard inequality for harmonically convex functions via generalized fractional integral operator. In addition, we establish an integral identity and some Fejér–Hadamard type integral inequalities for harmonically convex functions via a generalized fractional integral operator. Being generalizations, our results reproduce some known results.

Keywords: harmonically convex functions; Hermite–Hadamard inequality; generalized fractional integral operator; Mittag–Leffler function

MSC: 26A51, 26A33, 33E12

1. Introduction

Inequalities for convex functions, for example, the celebrated one is the Hermite–Hadamard inequality, providing a new horizon in the field of mathematical analysis. Many authors have been working on it continuously and several Hermite–Hadamard like integral inequalities have been established for many kinds of functions related to convex functions. Recently, a lot of integral inequalities of the Hermite–Hadamard type for harmonically convex functions via fractional integrals have been published (see [1–5] and references therein). The Hermite–Hadamard inequality for convex functions is stated in the following theorem.

Theorem 1. Let I be an interval of real numbers and $f : I \rightarrow \mathbb{R}$ be a convex function on I . Then, for all $a, b \in I$, the following inequality holds:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}.$$

Fejér gave a weighted version of the Hermite–Hadamard inequality stated as follows.

Theorem 2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function and $g : [a, b] \rightarrow \mathbb{R}$ is non-negative, integrable and symmetric to $\frac{a+b}{2}$. Then, the following inequality holds

$$f\left(\frac{a+b}{2}\right) \int_a^b g(x)dx \leq \int_a^b f(x)g(x)dx \leq \frac{f(a)+f(b)}{2} \int_a^b g(x)dx.$$

It is well known as the Fejér–Hadamard inequality. In the following, we give the definition of harmonically convex functions.

Definition 3. Reference [3] Let I be an interval of non-zero real numbers. Then, a function $f : I \rightarrow \mathbb{R}$ is said to be a harmonically convex function if the inequality

$$f\left(\frac{ab}{ta + (1-t)b}\right) \leq tf(b) + (1-t)f(a) \quad (1)$$

holds for $a, b \in I$ and $t \in [0, 1]$. If the inequality in Equation (1) is reversed, then f is said to be harmonically concave.

It is interesting to see that a function $f : (a, b) \rightarrow \mathbb{R}$, ($a > 0$) is harmonically convex iff the function $f \circ h : (\frac{1}{a}, \frac{1}{b}) \rightarrow \mathbb{R}$ is convex, where $h : (0, \infty) \rightarrow (0, \infty)$ is the hyperbola, i.e., $h(t) = 1/t$.

Definition 4. Reference [2] A function $h : [a, b] \subset \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ is said to be harmonically symmetric about $\frac{2ab}{a+b}$ if

$$h(x) = h\left(\frac{1}{\frac{1}{a} + \frac{1}{b} - \frac{1}{x}}\right)$$

holds for $x \in [a, b]$.

The following definition of the Riemann–Liouville fractional integral is the asset of fractional calculus.

Definition 5. Reference [6] Let $f \in L[a, b]$. Then, the two sided Riemann–Liouville fractional integral of f of order $\nu > 0$ is defined as:

$$I_{a+}^{\nu} f(x) = \frac{1}{\Gamma(\nu)} \int_a^x (x-t)^{\nu-1} f(t)dt, \quad x > a$$

and

$$I_{b-}^{\nu} f(x) = \frac{1}{\Gamma(\nu)} \int_x^b (t-x)^{\nu-1} f(t)dt, \quad x < b.$$

In the following, we give the definition of a generalized fractional integral operator which will help us to give a generalized Fejér–Hadamard inequality for harmonically convex functions and related results.

Definition 6. Reference [7] Let $\mu, \nu, k, l, \gamma, \delta$ be positive real numbers and $\omega \in \mathbb{R}$. Then, the generalized fractional integral operators containing a generalized Mittag–Leffler function for a real valued continuous function f are defined as follows:

$$\left(\epsilon_{\mu, \nu, l, \omega, a+}^{\gamma, \delta, k} f\right)(x) = \int_a^x (x-t)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k}(\omega(x-t)^{\mu}) f(t)dt,$$

and

$$\left(\epsilon_{\mu, \nu, l, \omega, b-}^{\gamma, \delta, k} f\right)(x) = \int_x^b (t-x)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k}(\omega(t-x)^{\mu}) f(t)dt,$$

where the function $E_{\mu, \nu, l}^{\gamma, \delta, k}$ is a generalized Mittag–Leffler function defined as:

$$E_{\mu, \nu, l}^{\gamma, \delta, k}(t) = \sum_{n=0}^{\infty} \frac{(\gamma)_{kn} t^n}{\Gamma(\mu n + \nu)(\delta)_{ln}}, \quad (2)$$

and $(\gamma)_{kn} = \frac{\Gamma(\gamma+kn)}{\Gamma(\gamma)}$.

For $\delta = l = 1$, the integral operator $\epsilon_{\mu,\nu,l,\omega,a^+}^{\gamma,\delta,k}$ reduces to an integral operator $\epsilon_{\mu,\nu,1,\omega,a^+}^{\gamma,1,k}$ containing generalized Mittag–Leffler function $E_{\mu,\nu,1}^{\gamma,1,k}$ introduced by Srivastava and Tomovski in [8]. Along with $\delta = l = 1$, in addition, if $k = 1$, then it reduces to an integral operator defined by Prabhaker in [9] containing Mittag–Leffler function $E_{\mu,\nu}^{\gamma}$. For $\omega = 0$, the integral operator $\epsilon_{\mu,\nu,l,\omega,a^+}^{\gamma,\delta,k}$ reduces to the Riemann–Liouville fractional integral operator.

In [7,8], properties of the generalized fractional integral operator $\epsilon_{\mu,\nu,l,\omega,a^+}^{\gamma,\delta,k}$ and the generalized Mittag–Leffler function $E_{\mu,\nu,l}^{\gamma,\delta,k}(t)$ are studied in brief. In [7], it is proved that $E_{\mu,\nu,l}^{\gamma,\delta,k}(t)$ is absolutely convergent for $k < l + \mu$ and $t \in \mathbb{R}$.

Since

$$|E_{\mu,\nu,l}^{\gamma,\delta,k}(t)| \leq \sum_{n=0}^{\infty} \left| \frac{(\gamma)_{kn} t^n}{\Gamma(\mu n + \nu)(\delta)_{ln}} \right|,$$

if we say that $\sum_{n=0}^{\infty} \left| \frac{(\gamma)_{kn} t^n}{\Gamma(\mu n + \nu)(\delta)_{ln}} \right| = S$, then

$$|E_{\mu,\nu,l}^{\gamma,\delta,k}(t)| \leq S.$$

We use this property of generalized Mittag–Leffler function subsequently in our results.

In addition, we use the following definitions of special functions known as beta and Euler type form of the hypergeometric functions (see [10]),

$$\beta(\mu, \nu) = \frac{\Gamma(\mu)\Gamma(\nu)}{\Gamma(\mu + \nu)} = \int_0^1 t^{\mu-1}(1-t)^{\nu-1} dt, \quad \mu, \nu > 0,$$

$${}_2F_1(a, b; c; w) = \frac{1}{\beta(b, c-b)} \int_0^1 t^{b-1}(1-t)^{c-b-1}(1-wt)^{-a} dt,$$

where $0 < b < c$ and $|w| < 1$.

In this paper, we give a generalized version of the Fejér–Hadamard inequality for harmonically convex functions via a generalized fractional integral operator. We also obtain bounds of the absolute differences of this generalized Fejér–Hadamard inequality for harmonically convex functions. Being generalizations, we reproduce the results proved in [1–3,5].

2. Main Results

To obtain our main results, we need the following lemmas.

Lemma 7. Reference [11] For $0 \leq a < b$ and $0 < \mu \leq 1$, we have

$$|a^\mu - b^\mu| \leq (b-a)^\mu.$$

Lemma 8. Let $f : [a, b] \subset \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$, $a < b$, be integrable and a harmonically symmetric function with respect to $\frac{2ab}{a+b}$. Then, for generalized fractional integrals, we have

$$\begin{aligned} \left(\epsilon_{\alpha,\beta,l,\omega,\frac{1}{b}^+}^{\gamma,\delta,k} f \circ g \right) \left(\frac{1}{a} \right) &= \left(\epsilon_{\alpha,\beta,l,\omega,\frac{1}{a}^-}^{\gamma,\delta,k} f \circ g \right) \left(\frac{1}{b} \right) \\ &= \frac{\left(\epsilon_{\alpha,\beta,l,\omega,\frac{1}{b}^+}^{\gamma,\delta,k} f \circ g \right) \left(\frac{1}{a} \right) + \left(\epsilon_{\alpha,\beta,l,\omega,\frac{1}{a}^-}^{\gamma,\delta,k} f \circ g \right) \left(\frac{1}{b} \right)}{2}, \end{aligned}$$

where $g(t) = \frac{1}{t}$ for all $t \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. Since f is harmonically symmetric about $\frac{2ab}{a+b}$, we have $f(\frac{1}{x}) = f(\frac{1}{\frac{1}{a} + \frac{1}{b} - x})$. By definition of generalized fractional integral operator

$$\left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{a} \right) = \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\frac{1}{a} - t \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(\frac{1}{a} - t \right)^{\mu} \right) f \left(\frac{1}{t} \right) dt, \quad (3)$$

replacing t by $\frac{1}{a} + \frac{1}{b} - x$ in Equation (3), we have

$$\begin{aligned} \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{a} \right) &= \int_{\frac{1}{b}}^{\frac{1}{a}} \left(x - \frac{1}{b} \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(x - \frac{1}{b} \right)^{\mu} \right) f \left(\frac{1}{\frac{1}{a} + \frac{1}{b} - x} \right) dx \\ &= \int_{\frac{1}{b}}^{\frac{1}{a}} \left(x - \frac{1}{b} \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(x - \frac{1}{b} \right)^{\mu} \right) f(x) dx. \end{aligned}$$

This implies

$$\left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{a} \right) = \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{b} \right). \quad (4)$$

By adding $\left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{a} \right)$ in both sides of Equation (4), we have

$$2 \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{a} \right) = \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{a} \right) + \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} f \circ g \right) \left(\frac{1}{b} \right). \quad (5)$$

Equations (4) and (5) give the required result. \square

Theorem 9. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a harmonically convex function. Let $a, b \in I, a < b, f \in L[a, b]$ and also let $g : [a, b] \rightarrow \mathbb{R}$ be a non-negative, integrable and harmonically symmetric function about $\frac{2ab}{a+b}$. Then, the following inequalities for generalized fractional integrals hold:

$$\begin{aligned} f \left(\frac{2ab}{a+b} \right) &\left[\left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{b}}^{\gamma, \delta, k} g \circ h \right) \left(\frac{1}{a} \right) + \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{a}}^{\gamma, \delta, k} g \circ h \right) \left(\frac{1}{b} \right) \right] \\ &\leq \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{b}}^{\gamma, \delta, k} f g \circ h \right) \left(\frac{1}{a} \right) + \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{a}}^{\gamma, \delta, k} f g \circ h \right) \left(\frac{1}{b} \right) \\ &\leq \frac{f(a) + f(b)}{2} \left[\left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{b}}^{\gamma, \delta, k} g \circ h \right) \left(\frac{1}{a} \right) + \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{a}}^{\gamma, \delta, k} g \circ h \right) \left(\frac{1}{b} \right) \right], \end{aligned} \quad (6)$$

where $\omega' = \omega \left(\frac{ab}{b-a} \right)^{\mu}$ and $h(t) = \frac{1}{t}$ for all $t \in \left[\frac{1}{b}, \frac{1}{a} \right]$.

Proof. Since f is a harmonically convex function, therefore, for $t \in [0, 1]$, we have

$$2f \left(\frac{2ab}{a+b} \right) \leq f \left(\frac{ab}{ta + (1-t)b} \right) + f \left(\frac{ab}{tb + (1-t)a} \right). \quad (7)$$

Multiplying both sides of Equation (7) by $t^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k}(\omega t^{\mu}) g \left(\frac{ab}{tb + (1-t)a} \right)$ and then integrating with respect to t over $[0, 1]$, we have

$$\begin{aligned} &2f \left(\frac{2ab}{a+b} \right) \int_0^1 t^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k}(\omega t^{\mu}) g \left(\frac{ab}{tb + (1-t)a} \right) dt \\ &\leq \int_0^1 t^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k}(\omega t^{\mu}) f \left(\frac{ab}{ta + (1-t)b} \right) g \left(\frac{ab}{tb + (1-t)a} \right) dt \\ &\quad + \int_0^1 t^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k}(\omega t^{\mu}) f \left(\frac{ab}{tb + (1-t)a} \right) g \left(\frac{ab}{tb + (1-t)a} \right) dt. \end{aligned} \quad (8)$$

By choosing $x = \frac{tb+(1-t)a}{ab}$ that is $\frac{ab}{ta+(1-t)b} = \frac{1}{\frac{1}{a} + \frac{1}{b} - x}$ in Equation (8), we have

$$\begin{aligned} & 2f\left(\frac{2ab}{a+b}\right) \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\frac{ab}{b-a}\right)^{\nu} \left(x - \frac{1}{b}\right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(\frac{ab}{b-a}\right)^{\mu} \left(x - \frac{1}{b}\right)^{\mu}\right) g\left(\frac{1}{x}\right) dx \\ & \leq \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\frac{ab}{b-a}\right)^{\nu} \left(x - \frac{1}{b}\right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(\frac{ab}{b-a}\right)^{\mu} \left(x - \frac{1}{b}\right)^{\mu}\right) f\left(\frac{1}{\frac{1}{a} + \frac{1}{b} - x}\right) g\left(\frac{1}{x}\right) dx \quad (9) \\ & + \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\frac{ab}{b-a}\right)^{\nu} \left(x - \frac{1}{b}\right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(\frac{ab}{b-a}\right)^{\mu} \left(x - \frac{1}{b}\right)^{\mu}\right) f\left(\frac{1}{x}\right) g\left(\frac{1}{x}\right) dx. \end{aligned}$$

Since f is harmonically symmetric about $\frac{2ab}{a+b}$, therefore, after simplification, Equation (9) becomes

$$\begin{aligned} & 2f\left(\frac{2ab}{a+b}\right) \int_{\frac{1}{b}}^{\frac{1}{a}} \left(x - \frac{1}{b}\right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega' \left(x - \frac{1}{b}\right)^{\mu}\right) g\left(\frac{1}{x}\right) dx \\ & \leq \int_{\frac{1}{a}}^{\frac{1}{b}} \left(\frac{1}{a} - x\right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega' \left(\frac{1}{a} - x\right)^{\mu}\right) f\left(\frac{1}{x}\right) g\left(\frac{1}{x}\right) dx \quad (10) \\ & + \int_{\frac{1}{b}}^{\frac{1}{a}} \left(x - \frac{1}{b}\right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega' \left(x - \frac{1}{b}\right)^{\mu}\right) f\left(\frac{1}{x}\right) g\left(\frac{1}{x}\right) dx. \end{aligned}$$

This implies

$$\begin{aligned} & 2f\left(\frac{2ab}{a+b}\right) \left(\epsilon_{\mu,\nu,l,\omega',\frac{1}{a}}^{\gamma,\delta,k} g \circ h\right) \left(\frac{1}{b}\right) \\ & \leq \left(\epsilon_{\mu,\nu,l,\omega',\frac{1}{b}}^{\gamma,\delta,k} f g \circ h\right) \left(\frac{1}{a}\right) + \left(\epsilon_{\mu,\nu,l,\omega',\frac{1}{a}}^{\gamma,\delta,k} f g \circ h\right) \left(\frac{1}{b}\right). \end{aligned}$$

Using Lemma 8 in the above inequality, we have

$$\begin{aligned} & f\left(\frac{2ab}{a+b}\right) \left[\left(\epsilon_{\mu,\nu,l,\omega',\frac{1}{a}}^{\gamma,\delta,k} g \circ h\right) \left(\frac{1}{b}\right) + \left(\epsilon_{\mu,\nu,l,\omega',\frac{1}{b}}^{\gamma,\delta,k} g \circ h\right) \left(\frac{1}{a}\right) \right] \quad (11) \\ & \leq \left(\epsilon_{\mu,\nu,l,\omega',\frac{1}{b}}^{\gamma,\delta,k} f g \circ h\right) \left(\frac{1}{a}\right) + \left(\epsilon_{\mu,\nu,l,\omega',\frac{1}{a}}^{\gamma,\delta,k} f g \circ h\right) \left(\frac{1}{b}\right). \end{aligned}$$

To prove the second half of the inequality, again from harmonically convexity of f on $[a, b]$ and for $t \in [0, 1]$, we have

$$f\left(\frac{ab}{tb+(1-t)a}\right) + f\left(\frac{ab}{ta+(1-t)b}\right) \leq f(a) + f(b). \quad (12)$$

Multiplying both sides of Equation (12) by $t^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k}(\omega t^{\mu}) g\left(\frac{ab}{tb+(1-t)a}\right)$, then integrating with respect to t over $[0, 1]$, we have

$$\begin{aligned} & \int_0^1 t^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k}(\omega t^{\mu}) f\left(\frac{ab}{tb+(1-t)a}\right) g\left(\frac{ab}{tb+(1-t)a}\right) dt \\ & + \int_0^1 t^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k}(\omega t^{\mu}) f\left(\frac{ab}{ta+(1-t)b}\right) g\left(\frac{ab}{tb+(1-t)a}\right) dt \quad (13) \\ & \leq [f(a) + f(b)] \int_0^1 t^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k}(\omega t^{\mu}) g\left(\frac{ab}{tb+(1-t)a}\right) dt. \end{aligned}$$

Setting $x = \frac{tb+(1-t)a}{ab}$ and by using harmonically symmetry of f with respect to $\frac{2ab}{a+b}$ in Equation (13), after simplification, we have

$$\begin{aligned} & \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{b}}^{\gamma, \delta, k} f g \circ h \right) \left(\frac{1}{a} \right) + \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{a}}^{\gamma, \delta, k} f g \circ h \right) \left(\frac{1}{b} \right) \\ & \leq [f(a) + f(b)] \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{b}}^{\gamma, \delta, k} g \circ h \right) \left(\frac{1}{a} \right). \end{aligned} \quad (14)$$

Using Lemma 8 in Equation (14), we have

$$\begin{aligned} & \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{b}}^{\gamma, \delta, k} f g \circ h \right) \left(\frac{1}{a} \right) + \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{a}}^{\gamma, \delta, k} f g \circ h \right) \left(\frac{1}{b} \right) \\ & \leq \frac{[f(a) + f(b)]}{2} \left[\left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{a}}^{\gamma, \delta, k} g \circ h \right) \left(\frac{1}{b} \right) + \left(\epsilon_{\mu, \nu, l, \omega', \frac{1}{b}}^{\gamma, \delta, k} g \circ h \right) \left(\frac{1}{a} \right) \right]. \end{aligned} \quad (15)$$

By joining Equations (11) and (15), we get Equation (6). \square

Remark 1. In Theorem 9,

- (i) if we put $\omega = 0$ along with $g(x) = 1$ and $\nu = 1$, then we get [3] [Theorem 2.4].
- (ii) if we put $\omega = 0$ along with $g(x) = 1$, then we get [5] [Theorem 4].
- (iii) if we put $\omega = 0$ along with $\nu = 1$, then we get [1] [Theorem 8].

Lemma 10. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on the interior of I and $f' \in L[a, b]$ where $a, b \in I$ and $a < b$. In addition, let $g : I \subset (0, \infty) \rightarrow \mathbb{R}$ be an integrable and harmonically symmetric function about $\frac{2ab}{a+b}$. Then, the following equality holds for generalized fractional integrals:

$$\begin{aligned} & \left(\frac{f(a) + f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \\ & - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (f g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (f g \circ h) \left(\frac{1}{b} \right) \right) \\ & = \left[\int_{\frac{1}{b}}^{\frac{1}{a}} \left(\int_{\frac{1}{b}}^t \left(\frac{1}{a} - s \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(\frac{1}{a} - s \right)^{\mu} \right) (g \circ h)(s) ds \right) (f \circ h)'(t) dt \right. \\ & \quad \left. - \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\int_t^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(s - \frac{1}{b} \right)^{\mu} \right) (g \circ h)(s) ds \right) (f \circ h)'(t) dt \right], \end{aligned}$$

where $h(t) = \frac{1}{t}$ for $t \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. To prove this lemma, we have

$$\begin{aligned} & \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\int_{\frac{1}{b}}^t \left(\frac{1}{a} - s \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(\frac{1}{a} - s \right)^{\mu} \right) (g \circ h)(s) ds \right) (f \circ h)'(t) dt \\ & = \left| \left(\int_{\frac{1}{b}}^t \left(\frac{1}{a} - s \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(\frac{1}{a} - s \right)^{\mu} \right) (g \circ h)(s) ds \right) (f \circ h)(t) \right|_{\frac{1}{b}}^{\frac{1}{a}} \\ & - \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\frac{1}{a} - t \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(\frac{1}{a} - t \right)^{\mu} \right) (g \circ h)(t) (f \circ h)(t) dt \\ & = \left(\int_{\frac{1}{b}}^{\frac{1}{a}} \left(\frac{1}{a} - s \right)^{\nu-1} E_{\mu, \nu, l}^{\gamma, \delta, k} \left(\omega \left(\frac{1}{a} - s \right)^{\mu} \right) (g \circ h)(s) ds \right) f(a) \\ & - \epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (f g \circ h) \left(\frac{1}{a} \right). \end{aligned}$$

This implies

$$\begin{aligned} & \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\int_{\frac{1}{b}}^t \left(\frac{1}{a} - s \right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(\frac{1}{a} - s \right)^\mu \right) (g \circ h)(s) ds \right) (f \circ h)'(t) dt \\ &= f(a) \epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{a} \right) - \epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{a} \right). \end{aligned} \quad (16)$$

Similarly,

$$\begin{aligned} & \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\int_t^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) (f \circ h)'(t) dt \\ &= \left| \left(\int_t^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) (f \circ h)(t) \right|_{\frac{1}{b}}^{\frac{1}{a}} \\ &+ \int_{\frac{1}{b}}^{\frac{1}{a}} \left(t - \frac{1}{b} \right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(t - \frac{1}{b} \right)^\mu \right) (g \circ h)(f \circ h)(t) dt \\ &= - \left(\int_{\frac{1}{b}}^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) f(b) \\ &+ \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right). \end{aligned}$$

This implies

$$\begin{aligned} & \int_{\frac{1}{b}}^{\frac{1}{a}} \left(\int_t^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{\nu-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) (f \circ h)'(t) dt \\ &= -f(b) \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{b} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right). \end{aligned} \quad (17)$$

Upon subtracting Equation (17) from Equation (16) and, using Lemma 8, we get the result. \square

Remark 2. In Lemma 10, if we take $g(x) = 1$ with $\omega = 0$, then it gives [5] (Lemma 3).

Theorem 11. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on the interior of I and $f' \in L[a, b]$ where $a, b \in I$ and $a < b$. If $|f'|$ is a harmonically convex function on $[a, b]$, $g : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ is a continuous and harmonically symmetric function with respect to $\frac{2ab}{a+b}$, then the following inequality for generalized fractional integrals holds

$$\begin{aligned} & \left| \left(\frac{f(a) + f(b)}{2} \right) \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \left. - \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_\infty S(b-a)^{\nu+1}}{\nu(ab)^{\nu-1}} (C_1(\nu)|f'(a)| + C_2(\nu)|f'(b)|), \end{aligned}$$

where $C_1(\nu) = \frac{b^{-2}}{(v+2)} {}_2F_1(2, 1; \nu+3; 1 - \frac{a}{b}) - \frac{b^{-2}}{(v+1)(v+2)} {}_2F_1(2, \nu+1; \nu+3; 1 - \frac{a}{b}) + \frac{2(a+b)^{-2}}{(v+1)(v+2)} {}_2F_1(2, \nu+1; \nu+3; \frac{b-a}{b+a})$ and $C_2(\nu) = \frac{b^{-2}}{(v+1)(v+2)} {}_2F_1(2, 2; \nu+3; 1 - \frac{a}{b}) - \frac{b^{-2}}{v+2} {}_2F_1(2, \nu+2; \nu+3; 1 - \frac{a}{b}) + \frac{4(a+b)^{-2}}{(v+1)} {}_2F_1(2, \nu+1; \nu+2; \frac{b-a}{b+a}) - \frac{2(a+b)^{-2}}{(v+1)(v+2)} {}_2F_1(\nu, \nu+1; \nu+3; \frac{b-a}{b+a})$ with $0 < \nu \leq 1$, $h(t) = \frac{1}{t}$ for all $t \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. By Lemma 10, we have

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \int_{\frac{1}{b}}^{\frac{1}{a}} \left| \left(\int_{\frac{1}{b}}^t \left(\frac{1}{a} - s \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(\frac{1}{a} - s \right)^\mu \right) (g \circ h)(s) ds \right) \right. \\ & \quad \left. - \left(\int_t^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) \right| |(f \circ h)'(t)| dt. \end{aligned} \quad (18)$$

Since g is harmonically symmetric with respect to $\frac{2ab}{a+b}$, therefore, $g\left(\frac{1}{t}\right) = g\left(\frac{1}{\frac{1}{a} + \frac{1}{b} - t}\right)$ for all $t \in \left[\frac{1}{b}, \frac{1}{a}\right]$, and we have

$$\begin{aligned} & \left| \left(\int_{\frac{1}{b}}^t \left(\frac{1}{a} - s \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(\frac{1}{a} - s \right)^\mu \right) (g \circ h)(s) ds \right) \right. \\ & \quad \left. - \left(\int_t^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) \right| \\ & = \left| \left(\int_{\frac{1}{a} + \frac{1}{b} - t}^{\frac{1}{a}} \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) \right. \\ & \quad \left. + \left(\int_{\frac{1}{a}}^t \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) \right| \\ & = \left| \left(\int_{\frac{1}{a} + \frac{1}{b} - t}^t \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) ds \right) \right| \\ & \leq \begin{cases} \int_t^{\frac{1}{a} + \frac{1}{b} - t} \left| \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) g(s) \right| ds, & t \in \left[\frac{1}{a}, \frac{a+b}{2ab}\right] \\ \int_{\frac{1}{a} + \frac{1}{b} - t}^t \left| \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) g(s) \right| ds, & t \in \left[\frac{a+b}{2ab}, \frac{1}{a}\right]. \end{cases} \end{aligned} \quad (19)$$

Using Equation (19) in (18), we have

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}+}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| r \\ & \leq \int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(\int_{\frac{1}{b}}^{\frac{1}{a} + \frac{1}{b} - t} \left| \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) \right| ds \right) |(f \circ h)'(t)| dt \\ & \quad + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\int_{\frac{1}{a} + \frac{1}{b} - t}^t \left| \left(s - \frac{1}{b} \right)^{v-1} E_{\mu,\nu,l}^{\gamma,\delta,k} \left(\omega \left(s - \frac{1}{b} \right)^\mu \right) (g \circ h)(s) \right| ds \right) |(f \circ h)'(t)| dt. \end{aligned} \quad (20)$$

Using $\|g\|_\infty = \sup_{t \in [a,b]} |g(t)|$ and absolute convergence of Mittag-Leffler function, the above inequality becomes

$$\begin{aligned} & \left| \left(\frac{f(a) + f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \|g\|_\infty S \left[\int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(\int_t^{\frac{1}{a} + \frac{1}{b} - t} \left(s - \frac{1}{b} \right)^{v-1} ds \right) |(f \circ h)'(t)| dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\int_{\frac{1}{a} + \frac{1}{b} - t}^t \left(\frac{1}{a} - s \right)^{v-1} ds \right) |(f \circ h)'(t)| dt \right] \\ & = \|g\|_\infty S \left[\int_{\frac{1}{b}}^{\frac{a+b}{2ab}} \left(\frac{(\frac{1}{a} - t)^v - (t - \frac{1}{b})^v}{v} \frac{1}{t^2} \right) |f' \left(\frac{1}{t} \right)| dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2ab}}^{\frac{1}{a}} \left(\frac{(t - \frac{1}{b})^v - (\frac{1}{a} - t)^v}{v} \frac{1}{t^2} \right) |f' \left(\frac{1}{t} \right)| dt \right]. \end{aligned} \quad (21)$$

Setting $t = \frac{ub + (1-u)a}{ab}$ in Equation (21), we have

$$\begin{aligned} & \left| \left(\frac{f(a) + f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_\infty S(b-a)^{v+1}}{v(ab)^{v-1}} \left[\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} |f' \left(\frac{ab}{(ub + (1-u)a)} \right)| du \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} |f' \left(\frac{ab}{(ub + (1-u)a)} \right)| du \right]. \end{aligned} \quad (22)$$

Since $|f'|$ is harmonically convex on $[a, b]$, it can be written as:

$$\left| f' \left(\frac{ab}{(ub + (1-u)a)^2} \right) \right| \leq u|f'(a)| + (1-u)|f'(b)|. \quad (23)$$

Using Equation (23) in Equation (22), we have

$$\begin{aligned} & \left| \left(\frac{f(a) + f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_\infty S(b-a)^{v+1}}{v(ab)^{v-1}} \left[\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} (u|f'(a)| + (1-u)|f'(b)|) du \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} (u|f'(a)| + (1-u)|f'(b)|) du \right], \end{aligned}$$

which is

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{\nu+1}}{\nu(ab)^{\nu-1}} \left[\left(\int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} u du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} u du \right) |f'(a)| \right. \\ & \quad \left. + \left(\int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} (1-u) du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} (1-u) du \right) |f'(b)| \right]. \end{aligned} \quad (24)$$

One can have, by using Lemma 7,

$$\begin{aligned} & \int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} u du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} u du \\ & = \int_0^1 \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} u du - \int_{\frac{1}{2}}^1 \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} u du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} u du \\ & = \int_0^1 \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} u du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} u du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} u du \\ & = \int_0^1 \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} u du + 2 \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} u du. \end{aligned} \quad (25)$$

On simplification, we get

$$\begin{aligned} & \int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} u du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} u du \\ & = \frac{b^{-2}}{(\nu+2)^2} {}_2F_1(2, 1; \nu+3; 1-\frac{a}{b}) - \frac{b^{-2}}{(\nu+1)(\nu+2)^2} {}_2F_1(2, \nu+1; \nu+3; 1-\frac{a}{b}) \\ & \quad + \frac{2(a+b)^{-2}}{(\nu+1)(\nu+2)^2} {}_2F_1(2, \nu+1; \nu+3; \frac{b-a}{b+a}) = C_1(\nu). \end{aligned} \quad (26)$$

Similarly,

$$\begin{aligned} & \int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} (1-u) du + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} (1-u) du \\ & = \frac{b^{-2}}{(\nu+1)(\nu+2)^2} {}_2F_1(2, 2; \nu+3; 1-\frac{a}{b}) - \frac{b^{-2}}{\nu+2} {}_2F_1(2, \nu+2; \nu+3; 1-\frac{a}{b}) \\ & \quad + \frac{4(a+b)^{-2}}{(\nu+1)^2} {}_2F_1(2, \nu+1; \nu+2; \frac{b-a}{b+a}) - \frac{2(a+b)^{-2}}{(\nu+1)(\nu+2)^2} {}_2F_1\left(\nu, \nu+1; \nu+3; \frac{b-a}{b+a}\right) = C_2(\nu). \end{aligned} \quad (27)$$

Using Equations (26) and (27) in Equation (24), we get the result. \square

Remark 3. In Theorem 11,

- (i) if we put $\omega = 0$, then we get [2] (Theorem 6).
- (ii) if we take $\nu = 1$ along with $\omega = 0$, then we get [2] (Corollary 1(1)).
- (iii) if we take $g(x) = 1$ along with $\omega = 0$, then we get [2] (Corollary 1(2)). (iv) if we take $\nu = 1$, $g(x) = 1$ along with $\omega = 0$, then we get [2] (Corollary 1(3)).

Theorem 12. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on the interior of I and $f' \in L[a, b]$ where $a, b \in I$ and $a < b$. If $|f'|^q, q > 1$ is a harmonically convex function on $[a, b]$, and $g : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be

a continuous and harmonically symmetric function about $\frac{2ab}{a+b}$, then the following inequality for generalized fractional integrals holds

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{\nu+1}}{\nu(ab)^{\nu-1}} \left[(C_3^{1-\frac{1}{q}}(\nu) (C_4(\nu) |f'(a)|^q + C_4(\nu) |f'(b)|^q)^{\frac{1}{q}} \right. \\ & \quad \left. + (C_6^{1-\frac{1}{q}}(\nu) (C_7(\nu) |f'(a)|^q + C_8(\nu) |f'(b)|^q)^{\frac{1}{q}} \right], \end{aligned}$$

where $C_3(\nu) = \frac{2(a+b)^{-2}}{\nu+1} {}_2F_1\left(2; \nu+1; \nu+3; \frac{b-a}{b+a}\right)$, $C_4(\nu) = \frac{(a+b)^{-2}}{(\nu+1)(\nu+2)} {}_2F_1\left(2; \nu+1; \nu+3; \frac{b-a}{b+a}\right)$, $C_5(\nu) = C_3(\nu) - C_4(\nu)$, $C_6(\nu) = \frac{b^{-2}}{\nu+1} {}_2F_1\left(2; 1; \nu+1; (1-\frac{a}{b})\right) - \frac{b^{-2}}{\nu+1} {}_2F_1\left(2; \nu+1; \nu+2; (1-\frac{a}{b})\right) + C_3(\nu)$, $C_7(\nu) = \frac{b^{-2}}{\nu+1} {}_2F_1\left(2; 1; \nu+2; (1-\frac{a}{b})\right) - \frac{b^{-2}}{\nu+1} {}_2F_1\left(2; \nu+1; \nu+2; (1-\frac{a}{b})\right) + C_4(\nu)$, $C_8(\nu) = C_6(\nu) - C_7(\nu)$ with $0 < \nu \leq 1$, $h(t) = \frac{1}{t}$ for all $t \in [\frac{1}{b}, \frac{1}{a}]$.

Proof. By inequality Equation (22) of Theorem 11, we have

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{\nu+1}}{\nu(ab)^{\nu-1}} \left[\int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} |f' \left(\frac{ab}{(ub+(1-u)a)} \right)| du \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} |f' \left(\frac{ab}{(ub+(1-u)a)} \right)| du \right]. \end{aligned} \quad (28)$$

Using power means, inequality Equation (28) becomes

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{\nu+1}}{\nu(ab)^{\nu-1}} \left[\left(\int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} du \right)^{1-\frac{1}{q}} \right. \\ & \quad \times \left(\int_0^{\frac{1}{2}} \frac{(1-u)^{\nu}-u^{\nu}}{(ub+(1-u)a)^2} |f' \left(\frac{ab}{(ub+(1-u)a)} \right)|^q du \right)^{\frac{1}{q}} \\ & \quad + \left(\int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} du \right)^{1-\frac{1}{q}} \\ & \quad \times \left. \left(\int_{\frac{1}{2}}^1 \frac{u^{\nu}-(1-u)^{\nu}}{(ub+(1-u)a)^2} |f' \left(\frac{ab}{(ub+(1-u)a)} \right)|^q du \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (29)$$

By using the harmonically convexity of $|f'|^q$ in Equation (29), we have

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{v+1}}{v(ab)^{v-1}} \left[\left(\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} du \right)^{1-\frac{1}{q}} \right. \\ & \quad \times \left(\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} (u|f'(a)|^q + (1-u)|f'(b)|^q) du \right)^{\frac{1}{q}} \\ & \quad + \left(\int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} du \right)^{1-\frac{1}{q}} \\ & \quad \times \left. \left(\int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} (u|f'(a)|^q + (1-u)|f'(b)|^q) du \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (30)$$

That is,

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu,\nu,l,\omega,\frac{1}{b}}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu,\nu,l,\omega,\frac{1}{a}-}^{\gamma,\delta,k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \|g\|_{\infty} S \frac{(b-a)^{v+1}}{v(ab)^{v-1}} \left[\left(\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} du \right)^{1-\frac{1}{q}} \right. \end{aligned} \quad (31)$$

$$\begin{aligned} & \times \left(\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} u du |f'(a)|^q + \int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} (1-u) du |f'(b)|^q \right)^{\frac{1}{q}} \\ & + \left(\int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} du \right)^{1-\frac{1}{q}} \left(\int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} u du |f'(a)|^q \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} (1-u) du |f'(b)|^q \right)^{\frac{1}{q}} \Bigg]. \end{aligned} \quad (32)$$

Now, we evaluate the integrals of Equation (31) by using Lemma 7:

$$\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} du \leq \int_0^{\frac{1}{2}} \frac{(1-2u)^v}{(ub + (1-u)a)^2} du = \frac{1}{2} \int_0^1 \frac{(1-u)^v}{(\frac{ub}{2} + (1-\frac{u}{2})a)^2} du. \quad (33)$$

Substituting $u = 1 - w$ in Equation (33), we have

$$\begin{aligned} & \int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} du \leq 2(a+b)^{-2} \int_0^1 w^v \left(1 - w \left(\frac{b-a}{b+a} \right) \right)^{-2} dw \\ & = 2 \frac{(a+b)^{-2}}{v+1} {}_2F_1 \left(2; v+1; v+2; \frac{b-a}{b+a} \right) = C_3(v). \end{aligned}$$

Similarly,

$$\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} u du \leq \int_0^{\frac{1}{2}} \frac{(1-2u)^v}{(ub + (1-u)a)^2} u du = \frac{1}{4} \int_0^1 \frac{u(1-u)^v}{(\frac{ub}{2} + (1-\frac{u}{2})a)^2} du. \quad (34)$$

Substituting $u = 1 - w$ in Equation (34), we have

$$\begin{aligned} \int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} u du &\leq (a+b)^{-2} \int_0^1 (1-w)w^v \left(1 - w\left(\frac{b-a}{b+a}\right)\right)^{-2} dw \\ &= \frac{(a+b)^{-2}}{(\nu+1)(\nu+2)} {}_2F_1\left(2; \nu+1; \nu+3; \frac{b-a}{b+a}\right) = C_4(\nu). \end{aligned}$$

$$\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v(1-u)}{(ub + (1-u)a)^2} (1-u) du \leq C_3(\nu) - C_4(\nu) = C_5(\nu). \quad (35)$$

$$\begin{aligned} \int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} du &= \int_0^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} du + \int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} du \\ &\leq \frac{b^{-2}}{\nu+1} {}_2F_1\left(2; 1; \nu+2; \left(1 - \frac{a}{b}\right)\right) - \frac{b^{-2}}{\nu+1} {}_2F_1\left(2; \nu+1; \nu+2; \left(1 - \frac{a}{b}\right)\right) + C_3(\nu) = C_6(\nu). \end{aligned} \quad (36)$$

$$\begin{aligned} \int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} u du &= \int_0^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} u du + \int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub + (1-u)a)^2} u du \\ &\leq \frac{b^{-2}}{\nu+2} {}_2F_1\left(2; 1; \nu+3; \left(1 - \frac{a}{b}\right)\right) - \frac{b^{-2}}{(\nu+1)(\nu+2)} {}_2F_1\left(2; \nu+1; \nu+3; \left(1 - \frac{a}{b}\right)\right) + C_4(\nu) = C_7(\nu) \end{aligned}$$

and

$$\int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub + (1-u)a)^2} (1-u) du \leq C_6(\nu) - C_7(\nu) = C_8(\nu). \quad (37)$$

Using Equations (34)–(37) in Equation (31), we get the result. \square

Remark 4. The following remarks can be obtained by giving specific values to parameters in Theorem 12:

- (i) If we take $\omega = 0$, then we get [2] (Theorem 7).
- (ii) If we take $\nu = 1$ along with $\omega = 0$, then we get [2] (Corollary 2(1)).
- (iii) If we take $g(x) = 1$ along with $\omega = 0$, then we get [2] (Corollary 2(2)).
- (iv) If we take $\nu = 1$, $g(x) = 1$ along with $\omega = 0$, then we get [2] (Corollary 2(3)).

Theorem 13. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on the interior of I such that $f' \in L[a, b]$, where $a, b \in I$ and $a < b$. If $|f'|^q$, $q > 1$ is a harmonically convex function on $[a, b]$, and $g : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a continuous and harmonically symmetric function about $\frac{2ab}{a+b}$. Then, the following inequality for generalized fractional integrals holds

$$\begin{aligned} &\left| \left(\frac{f(a) + f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ &\quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ &\leq \frac{\|g\|_{\infty} S(b-a)^{\nu+1}}{\nu(ab)^{\nu-1}} \left(C_9^{\frac{1}{p}}(\nu) \left(\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right)^{\frac{1}{q}} + C_{10}^{\frac{1}{p}}(\nu) \left(\frac{3|f'(a)|^q + |f'(b)|^q}{8} \right)^{\frac{1}{q}} \right), \end{aligned}$$

where $C_9(\nu) = \frac{(a+b)^{-2p}}{2^{-2p+1}(\nu p+1)} {}_2F_1(2p, \nu p+1; \nu p+2; \frac{b-a}{b+a})$ and $C_{10}(\nu) = \frac{b^{-2p}}{2(\nu p+1)} {}_2F_1(2p, 1; \nu p+2; \frac{1}{2}(1 - \frac{a}{b}))$ with $0 \leq \nu < 1$, $h(t) = \frac{1}{t}$ for all $t \in [\frac{1}{b}, \frac{1}{a}]$ and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. By inequality Equation (22) of Theorem 11, we have

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{v+1}}{v(ab)^{v-1}} \left[\int_0^{\frac{1}{2}} \frac{(1-u)^v - u^v}{(ub+(1-u)a)^2} |f' \left(\frac{ab}{(ub+(1-u)a)} \right)| du \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 \frac{u^v - (1-u)^v}{(ub+(1-u)a)^2} |f' \left(\frac{ab}{(ub+(1-u)a)} \right)| du \right]. \end{aligned} \quad (38)$$

By using Hölder inequality and harmonically convexity of $|f'|^q$, Equation (38) follows:

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{v+1}}{v(ab)^{v-1}} \left[\left(\int_0^{\frac{1}{2}} \frac{((1-u)^v - u^v)^p}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \right. \\ & \quad \times \left(\int_0^{\frac{1}{2}} (u|f'(a)|^q + (1-u)|f'(b)|^q) du \right)^{\frac{1}{q}} \\ & \quad \left. + \left(\int_{\frac{1}{2}}^1 \frac{(u^v - (1-u)^v)^p}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \left(\int_{\frac{1}{2}}^1 (u|f'(a)|^q + (1-u)|f'(b)|^q) du \right)^{\frac{1}{q}} \right]. \end{aligned}$$

After simplification, we have

$$\begin{aligned} & \left| \left(\frac{f(a)+f(b)}{2} \right) \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (g \circ h) \left(\frac{1}{b} \right) \right) \right. \\ & \quad \left. - \left(\epsilon_{\mu, \nu, l, \omega, \frac{1}{b}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{a} \right) + \epsilon_{\mu, \nu, l, \omega, \frac{1}{a}}^{\gamma, \delta, k} (fg \circ h) \left(\frac{1}{b} \right) \right) \right| \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{v+1}}{v(ab)^{v-1}} \left[\left(\int_0^{\frac{1}{2}} \frac{((1-u)^v - u^v)^p}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \right. \\ & \quad \times \left(\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right)^{\frac{1}{q}} \\ & \quad \left. + \left(\int_{\frac{1}{2}}^1 \frac{(u^v - (1-u)^v)^p}{(ub+(1-u)a)^{2p}} du \right)^{\frac{1}{p}} \left(\frac{|f'(a)|^q + 3|f'(b)|^q}{8} \right)^{\frac{1}{q}} \right]. \end{aligned} \quad (39)$$

We evaluate the integrals by using Lemma 7

$$\int_0^{\frac{1}{2}} \frac{((1-u)^v - u^v)^p}{(ub+(1-u)a)^{2p}} du \leq \int_0^{\frac{1}{2}} \frac{(1-2u)^{vp}}{(ub+(1-u)a)^{2p}} du = \frac{1}{2} \int_0^1 \frac{(1-u)^{vp}}{(\frac{ub}{2} + (1-\frac{u}{2})a)^{2p}} du. \quad (40)$$

Putting $u = 1 - w$ in Equation (40), we have

$$\begin{aligned} & \int_0^{\frac{1}{2}} \frac{((1-u)^v - u^v)^p}{(ub+(1-u)a)^{2p}} du \leq \frac{1}{2} \int_0^1 w^{vp} \left(\frac{a+b}{2} \right)^{-2p} \left(1 - w \left(\frac{b-a}{b+a} \right) \right)^{-2p} dw \\ & = \frac{(a+b)^{-2p}}{2^{-2p+1}(vp+1)} {}_2F_1 \left(2p, vp+1; vp+2; \frac{b-a}{b+a} \right) = C_9(v). \end{aligned} \quad (41)$$

Similarly,

$$\int_{\frac{1}{2}}^1 \frac{(u^v - (1-u)^v)^p}{(ub + (1-u)a)^{2p}} du \leq \int_{\frac{1}{2}}^1 \frac{(2u-1)^{vp}}{(ub + (1-u)a)^{2p}} du \quad (42)$$

and putting $u = 1 - w$ on the right-hand side of inequality Equation (42), we have

$$\begin{aligned} \int_{\frac{1}{2}}^1 \frac{(u^v - (1-u)^v)^p}{(ub + (1-u)a)^{2p}} du &\leq \int_0^{\frac{1}{2}} \frac{(1-2w)^{vp}}{((1-w)b + wa)^{2p}} dw \\ &= \frac{1}{2} \int_0^1 \frac{(1-w)^{vp}}{(\frac{wa}{2} + (1-\frac{w}{2})b)^{2p}} dw. \\ &= \frac{b^{-2p}}{2(vp+1)} {}_2F_1(2p, 1; vp+2; \frac{1}{2}(1 - \frac{a}{b})) = C_{10}(v). \end{aligned} \quad (43)$$

Using Equations (41) and (43) in Equation (39), we get the result. \square

Remark 5. On giving particular values to parameter in Theorem 13, we have the following results:

- (i) If we put $\omega = 0$, then we get [2] (Theorem 8).
- (i) If we put $v = 1$ along with $\omega = 0$, then we get [2] (Corollary 3(1)).
- (ii) If we put $g(t) = 1$ along with $\omega = 0$, then we get [2] (Corollary 3(2)).
- (iv) If we put $v = 1$, $g(t) = 1$ along with $\omega = 0$, then we get [2] (Corollary 3(3)).

3. Conclusions

We have obtained a generalized Fejér–Hadamard inequality for harmonically convex functions via a generalized fractional integral operator. This inequality includes several inclusions—for example, Fejér–Hadamard and Hermite–Hadamard inequalities for harmonically convex functions via Riemann–Liouville fractional integrals. Taking different specific values of parameters in the generalized Mittag–Leffler function, one can obtain results for some known fractional integral operators—for example, for fractional integral operators defined in [8,9]. In addition, we have established some bounds of the difference of the generalized Fejér–Hadamard inequality, in particular several bounds for particular values of parameters involved in the generalized Mittag–Leffler function.

Author Contributions: S.M.K. proposed the problem and supervise this work, G.A. and G.F. proved the results. W.N. collect literature and compare the results of this paper with existing results in literature and wrote the paper.

Funding: The research work of Ghulam Farid is supported by the Higher Education Commission of Pakistan under NRP 2016, Project No. 5421.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Chen, F.; Wu, S. Fejér and Hermite–Hadamard type inequalities for harmonically convex functions. *J. Appl. Math.* **2014**, *2014*, 386806.
- Kunt, M.; İşcan, İ.; Yazi, N.; Gozutok, U. On new inequalities of Hermite–Hadamard Fejér type inequalities for harmonically convex functions via fractional integrals. *SpringerPlus* **2016**, *5*, 635, doi:10.1186/s40064-016-2215-4.
- İşcan, İ. Hermite Hadamard type inequalities for harmonically convex functions. *Hacet. J. Math. Stat.* **2014**, *43*, 935–942.
- İşcan, İ.; Mehmet, K. Hermite–Hadamard–Fejér type inequalities for harmonically convex functions via fractional integrals. *Stud. Univ. Babeş-Bolyai Math.* **2015**, *60*, 355–366.
- İşcan, İ.; Wu, S. Hermite–Hadamard type inequalities for harmonically convex functions via fractional integrals. *Appl. Math. Comput.* **2014**, *238*, 237–244.
- Tomovski, Z.; Hiller, R.; Srivastava, H.M. Fractional and operational calculus with generalized fractional derivative operators and Mittag–Leffler function. *Integral Transforms Spec. Funct.* **2011**, *21*, 797–814.
- Salim, L.T.O.; Faraj, A.W. A Generalization of Mittaf–Leffler function and integral operator associated with integral calculus. *J. Frac. Calc. Appl.* **2012**, *3*, 1–13.

8. Srivastava, H.M.; Tomovski, Z. Fractional calculus with an integral operator containing generalized Mittag–Leffler function in the kernel. *Appl. Math. Comput.* **2009**, *211*, 198–210.
9. Prabhakar, T.R. A singular integral equation with a generalized Mittag–Leffler function in the kernel. *Yokohama Math. J.* **1971**, *19*, 7–15.
10. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of Fractional Differential Equations*; Elsevier: New York, NY, USA; London, UK, 2006.
11. Prudnikov, A.P.; Brychkov, Y.A.; Marichev, O.J. *Integral and Series of Elementary Functions*; Nauka: Moscow, Russia, 1981.



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).