

a. FEV1 (% of predicted value)

Responders			Non-responders		
T0 (68 patients)	78.8 ± 19.7		T0 (22 patients)	70.9 ± 20.4	
T1 (68 patients)	86.8 ± 19.7		T1 (22 patients)	77.8 ± 17.4	
T2 (55 patients)	87.7 ± 19.1		T2 (18 patients)	73.6 ± 16.5	
T3 (36 patients)	90.8 ± 19.5		T3 (17 patients)	83.6 ± 17.7	
T4 (25 patients)	85.9 ± 18.4		T4 (14 patients)	76.4 ± 19.7	
p			p		
T0	T1	n.s.	T0	T1	n.s.
	T2	n.s.		T2	n.s.
	T3	0.002		T3	0.001
	T4	n.s.		T4	n.s.

b. FENO (ppb)

Responders			Non-responders		
T0 (68 patients)	46.4 ± 30.9		T0 (22 patients)	42.9 ± 25.9	
T1 (68 patients)	33.9 ± 23.3		T1 (22 patients)	39.9 ± 27.2	
T2 (55 patients)	29.6 ± 23.3		T2 (18 patients)	37.3 ± 28.2	
T3 (36 patients)	23.2 ± 18.1		T3 (17 patients)	34.1 ± 27.3	
T4 (25 patients)	19.1 ± 10.5		T4 (14 patients)	21.3 ± 12.6	
p			p		
T0	T1	0.043	T0	T1	n.s.
	T2	0.035		T2	n.s.
	T3	0.008		T3	n.s.
	T4	<0.001		T4	n.s.

Table S1 – Trends of the analyzed variables over time in responders and non-responders, comparisons with baseline. Panel a: FEV1 (% of predicted value); Panel b:- FENO (ppb).