

Selective Extraction of Platinum(IV) from the Simulated Secondary Resources Using Simple Secondary Amide and Urea Extractants

Yuki Ueda ^{1,2,*}, Shintaro Morisada ³, Hidetaka Kawakita ³ and Keisuke Ohto ³

¹ Materials Sciences Research Center, Japan Atomic Energy Agency, Tokai, Ibaraki 391-1195, Japan

² Advanced Science Research Center, Japan Atomic Energy Agency, Tokai, Ibaraki 391-1195, Japan

³ Department of Chemistry and Applied Chemistry, Faculty of Science and Engineering, Saga University, 1-Honjo, Saga 840-8502, Japan; morisada@cc.saga-u.ac.jp (S.M.); kawakita@cc.saga-u.ac.jp (H.K.); ohtok@cc.saga-u.ac.jp (K.O.)

* Correspondence: ueda.yuki@jaea.go.jp; Tel.: +81-29-284-3960

CONTENT

1. ¹H NMR spectra of (a) *N*-(2-ethylhexyl)acetamide (MonoAA), (b) MonoAA exchanged with D₂O, and (c) 2-ethylhexylamine as a starting material.

2. IR spectra of *N*-(2-ethylhexyl)acetamide (MonoAA).



3. ¹H NMR spectra of (a) *N*-(2-ethylhexyl)octanamide (MonoOA), (b) MonoOA exchanged with D₂O, and (c) 2-ethylhexylamine as a starting material.

4. IR spectra of *N*-(2-ethylhexyl)octanamide (MonoOA).

5. ¹H NMR spectra of (a) *N,N*-bis(2-ethylhexyl)acetamide (BisAA), (b) BisAA exchanged with D₂O, and (c) di(2-ethylhexyl)amine as a starting material.

6. IR spectra of *N,N*-bis(2-ethylhexyl)acetamide (BisAA).


1. ^1H NMR Spectra of N-(2-ethylhexyl)acetamide (MonoAA).



¹ H NMR	δ / ppm	Assignment	Multiplicity	Integral ratio
CDCl ₃	0.90	a	m	5.60 (6H)
TMS	1.28–1.43	b	m	8.19 (9H)
	2.00	c	s	2.53 (3H)
	3.19	d	t	2.00 (2H)
	5.60	e	br	0.85 (1H)

Figure S1. ^1H NMR spectra of (a) *N*-(2-ethylhexyl)acetamide (MonoAA), (b) MonoAA exchanged with D_2O , and (c) 2-ethylhexylamine as a starting material.



2. IR Spectra of *N*-(2-ethylhexyl)acetamide (MonoAA).



FT-IR	ν / cm^{-1}	Assignment
ATR	1558	N–H (bending)
SeZn	1647	C=O (stretching)
	2927	–CH ₂ – (stretching)
	2958	–CH ₃ (stretching)
	3088	N–H (stretching)
	3290	N–H (stretching)

Figure S2. IR spectra of *N*-(2-ethylhexyl)acetamide (MonoAA).


3. ^1H NMR Spectra of *N*-(2-ethylhexyl)octanamide (MonoOA).



^1H NMR	δ / ppm	Assignment	Multiplicity	Integral ratio
CDCl ₃	0.89	a	m	8.37 (9H)
TMS	1.28–1.42	b	m	17.3 (17H)
	1.62	c	m	1.74 (2H)
	2.17	d	t	2.03 (2H)
	3.20	e	t	2.00 (2H)
	5.49	f	br	1.24 (1H)

Figure S3. ^1H NMR spectra of (a) *N*-(2-ethylhexyl)octanamide (MonoOA), (b) MonoOA exchanged with D₂O, and (c) 2-ethylhexylamine as a starting material.


4. IR Spectra of *N*-(2-ethylhexyl)acetamide (MonoAA).



FT-IR	ν / cm^{-1}	Assignment
ATR	1556	N–H (bending)
SeZn	1641	C=O (stretching)
	2926	–CH ₂ – (stretching)
	2956	–CH ₃ (stretching)
	3084	N–H (stretching)
	3290	N–H (stretching)


Figure S4. IR spectra of *N*-(2-ethylhexyl)octanamide (MonoOA).

5. ^1H NMR Spectra of *N,N*-bis(2-ethylhexyl)acetamide (BisAA).



^1H NMR	δ / ppm	Assignment	Multiplicity	Integral ratio
CDCl_3	0.84–0.92	a	m	14.3 (12H)
TMS	1.26	b	m	21.8 (16H)
	2.09	c	s	3.00 (3H)
	3.13	d	d	1.69 (4H)
	3.27	e	m	1.61 (2H)

Figure S5. ^1H NMR spectra of (a) *N,N*-bis(2-ethylhexyl)acetamide (BisAA), (b) BisAA exchanged with D_2O , and (c) di(2-ethylhexyl)amine as a starting material.

6. IR Spectra of *N,N*-bis(2-ethylhexyl)acetamide (BisAA).

FT-IR	ν / cm^{-1}	Assignment
ATR	1647	C=O (stretching)
SeZn	2927	-CH ₂ - (stretching)
	2959	-CH ₃ (stretching)

Figure S6. IR spectra of *N,N*-bis(2-ethylhexyl)acetamide (BisAA).