

Supplementary information

The Synthesis, crystal structure and magnetic properties of mono-scorpionate Eu(III) complexes

Kira E. Vostrikova*, Taisiya S. Sukhikh and Alexander N. Lavrov

Table S1. Crystal data and structure refinement for the compounds.

Identification code	(1)	(1a)	(2)	(3)
Empirical formula	C ₁₄ H ₁₈ EuN ₁₁ O ₁₀	C ₁₀ H ₁₂ EuN ₉ O ₁₀	C ₁₈ H ₂₅ EuN ₁₀ O ₉	C _{32.9} H _{37.24} Eu ₂ F ₁₈ N ₁₄ O _{20.73} S ₆
Formula weight	652.35	570.25	677.44	1798.66
Temperature/K	150(2)	150(2)	150(2)	150(2)
Space group	<i>P</i> -1	<i>I</i> 2/c	<i>P</i> 2 ₁ /n	<i>P</i> 2 ₁ /c
<i>a</i> /Å	9.4083(2)	14.9027(3)	10.8259(6)	21.596(3)
<i>b</i> /Å	11.0349(3)	15.6143(3)	14.9846(7)	17.912(2)
<i>c</i> /Å	11.7157(3)	16.1279(4)	16.4034(8)	18.668(2)
α°	99.6960(10)	90	90	90
β°	93.4410(10)	108.4710(10)	95.786(2)	115.419(4)
γ°	102.6850(10)	90	90	90
Volume/Å ³	1163.71(5)	3559.55(13)	2647.4(2)	6522.3(14)
<i>Z</i>	2	8	4	4
ρ_{calc} g/cm ³	1.862	2.128	1.700	1.832
μ /mm ⁻¹	2.769	3.601	2.434	2.224
<i>F</i> (000)	644.0	2224.0	1352.0	3530.0
Crystal size/mm ³	0.16 × 0.15 × 0.10	0.20 × 0.16 × 0.08	0.10 × 0.10 × 0.08	0.12 × 0.10 × 0.03
Radiation	MoKα ($\lambda = 0.71073$)	MoKα ($\lambda = 0.71073$)	MoKα ($\lambda = 0.71073$)	MoKα ($\lambda = 0.71073$)
2θ range for data collection/°	5.168 to 59.232	3.886 to 61.058	4.316 to 61.048	3.086 to 51.484
Index ranges	-13 ≤ <i>h</i> ≤ 13, -15 ≤ <i>k</i> ≤ 15, -16 ≤ <i>l</i> ≤ 16	-21 ≤ <i>h</i> ≤ 21, -22 ≤ <i>k</i> ≤ 22 -22 ≤ <i>l</i> ≤ 22	-13 ≤ <i>h</i> ≤ 15, -21 ≤ <i>k</i> ≤ 21 -23 ≤ <i>l</i> ≤ 22	
Reflections collected	26611	44683	36783	
Independent reflections	6515 [$R_{\text{int}}=0.0256$, $R_{\text{sigma}}=0.0220$]	5332 [$R_{\text{int}}=0.0308$, $R_{\text{sigma}}=0.0207$]	8022 [$R_{\text{int}}=0.0473$, $R_{\text{sigma}}=0.0393$]	5332 [$R_{\text{int}}=0.0308$, $R_{\text{sigma}}=0.0207$]
Data/restraints/parameters	6515/3/333	5332/2/277	8022/0/350	5332/2/277
Goodness-of-fit on <i>F</i> ²	1.062	1.091	1.072	1.091
Final R indexes [$I \geq 2\sigma(I)$]	$R_1=0.0151$, $wR_2=0.0357$	$R_1=0.0184$, $wR_2=0.0370$	$R_1=0.0276$, $wR_2=0.0525$	$R_1=0.0184$, $wR_2=0.0370$
Final R indexes [all data]	$R_1=0.0156$, $wR_2=0.0360$	$R_1=0.0281$, $wR_2=0.0414$	$R_1=0.0354$, $wR_2=0.0581$	$R_1=0.0281$, $wR_2=0.0414$
Largest diff. peak/hole/e Å ⁻³	0.42/-0.55	0.55/-0.84	0.52/-0.62	0.55/-0.84




Figure S1. The asymmetric unit in the crystal structure of [Eu(HCPZ₃)H₂O(CF₃SO₃)₃]₂ (3).




Figure S2. Square antiprism coordination environment - of Eu³⁺ sites in [Eu(HCPz₃)H₂O(CF₃SO₃)₃]₂ (3) for: (a) around Eu1 central atom; (b) around Eu2 central atom.

Table S2. Geometry analysis of the complexes by SHAPE software.

Compound	CN	Polyhedron geometry		
[Eu(HCPz ₃)(NO ₃) ₃ H ₂ O] 1	[ML ₁₀]	Sphenocorona, C _{2v}	Bicapped square antiprism, D _{4d}	
		3.534	6.356	
[Eu(HCPz ₃)(NO ₃) ₃ H ₂ O] 1a	[ML ₁₀]	Sphenocorona, C _{2v}	Bicapped square antiprism, D _{4d}	
		3.225	4.793	
[Eu(HC(Pz ^{Me} ₂) ₃)(NO ₃) ₃] 2	[ML ₉]	Spherical tricapped trigonal prism, D _{3h}	Spherical capped square antiprism, C _{4v}	Muffin, C _s
		2.05	2.910	3.027
[Dy(Rad)(NO ₃) ₃]		2.220	3.025	3.256
[Eu(HCPz ₃)H ₂ O(OTf) ₃] 3				
Eu01	[ML ₈]	Square antiprism D _{4d}	Triangular dodecahedron, D _{2d}	Biaugmented trigonal prism, C _{2v}
		0.354	2.184	1.543
Eu02	[ML ₈]	Square antiprism D _{4d}	Triangular dodecahedron, D _{2d}	Biaugmented trigonal prism, C _{2v}
		0.426	1.919	1.368




Figure S3. Dimerized species in **1** (left) and **1a** (right).




Figure S4. The temperature-dependent molar magnetic susceptibility χ_M of **1–3** at a field of 1 kOe (open red circles). Solid green circles show the χ_M data after subtraction of the contribution from Eu(II) impurities (0.01–0.017 %).