

Article

# Combining Celery Oleoresin, Limonene and Rhamnolipid as New Strategy to Control Endospore-Forming *Bacillus cereus*

Paula de Camargo Bertuso <sup>1</sup>, Débora M. Drappé Mayer <sup>1</sup> and Marcia Nitschke <sup>2,\*</sup>

<sup>1</sup> Interunits Graduate Program in Bioengineering (EESC/FMRP/IQSC), University of São Paulo, Trabalhador São-Carlense Av., 400, São Carlos, SP 13566-590, Brazil; paulabertuso@usp.br (P.d.C.B.); debora\_mayer@hotmail.com (D.M.D.M.)

<sup>2</sup> São Carlos Institute of Chemistry (IQSC), University of São Paulo, Trabalhador São-Carlense Av., 400, P.O. Box 780, São Carlos, SP 13560-970, Brazil

\* Correspondence: nitschke@iqsc.usp.br

**Abstract:** Foodborne diseases (FBD) are a great problem worldwide, leading millions of people to seek medical help and to significant economic losses for industry. Among the agents implicated in FDB is *Bacillus cereus*, a Gram-positive, toxigenic and endospore-forming bacterium. In this study, rhamnolipid (RL) biosurfactant, celery oleoresin (OR) and limonene (LN) were evaluated as bio-based alternatives for controlling the growth of vegetative cells and endospores of *B. cereus*. To address their antimicrobial activity, the compounds were tested separately and in combination. Results demonstrate that, when combined with RL, both OR and LN have lower minimal inhibitory concentration (MIC) values and increased endospore inhibition potential. A percentage of endospore inhibition from 73% to 98%, corresponding to a 2.8–3.6 log reduction in spore outgrowth, was observed. RL inhibited *B. cereus* growth and endospore germination and potentially enhanced the antimicrobial efficacy of the natural hydrophobic compounds tested.

**Keywords:** *Bacillus cereus*; rhamnolipid; *Apium graveolens*; oleoresin; limonene



**Citation:** Bertuso, P.d.C.; Mayer, D.M.D.; Nitschke, M. Combining Celery Oleoresin, Limonene and Rhamnolipid as New Strategy to Control Endospore-Forming *Bacillus cereus*. *Foods* **2021**, *10*, 455.  
<https://doi.org/10.3390/foods10020455>

Academic Editors: Xanel Vecino and Ana Belén Moldes

Received: 8 December 2020

Accepted: 16 February 2021

Published: 19 February 2021

Corrected: 2 February 2023

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Foodborne diseases (FBD) resulting from the ingestion of foodstuffs contaminated with microorganisms or chemicals represent a growing public health problem worldwide [1]. It is estimated that 600 million people fall sick and 420 thousand die every year after ingestion of contaminated food [2]. Between 2007 and 2017, 96% of FBD outbreaks registered in Brazil were caused by bacteria, including *Bacillus cereus* [3]. In 2016, a food poisoning outbreak was reported in New York from a Chinese fast food chain and 33 *B. cereus* isolates were found on bean samples [4]. In outbreaks registered in France from 2007 to 2014, as little as 400 CFU/g *B. cereus* was found in the incriminated foods and reported to be enough to cause symptoms [5]. Those studies help illustrate how difficult it is to eliminate the bacterium in industrial processing plants and show its capacity to cause disease in humans even when present at low numbers. Considering the importance of controlling such pathogens to public health, food manufactures are continuously searching for innovative methods to guarantee the safety of their products.

Consumers' preference for natural additives rather than artificial ones [6] stimulates the industry to find new bio-based and green preservatives. The use of natural biocontrol agents derived from microbes or plants such as bacteriocins, endolysins and essential oils (EOs) is gaining increasing interest, especially due to the development of microbial tolerance to disinfectants [7].

EOs are complex, volatile and hydrophobic compounds formed by plants as secondary metabolites [8,9]. Their antimicrobial activity against food pathogens has been extensively reported [10–13]. Oleoresins (OR), in contrast, are viscous mixtures of essential oils and resins that are extracted from spices through organic solvents [14,15]. ORs can be found

in liquid form when adding solvents such as propylene glycol, which also facilitates their use in food products [14]. ORs contain both volatile and non-volatile components and present a high shelf life since they are practically absent of water, thus reducing oxidative degradation, flavor loss and microbial contamination [16]. D-limonene (LN) is the major constituent of several EOs, especially from citrus species, and its presence has been associated with the antimicrobial activity of such compounds [11,12].

Although plant-derived oils and oleoresins have shown antimicrobial effects, their use is limited, due to the higher concentrations needed to be effective, their low water solubility and their strong sensorial impact [17]. An alternative to this problem is the use of emulsifiers, such as surfactants, that permit improving water solubility, reducing the amount of the oil needed and its undesirable sensorial effects.

Microbial-derived surfactants can replace synthetic ones with some advantages since they show similar surfactant and emulsifier characters, are eco-friendly and bio-based and are good candidates in the development of innovative “green” food products [18]. Rhamnolipids (RL) are glycolipid biosurfactants, produced primarily by *Pseudomonas* sp., that have shown potential as antimicrobial, anti-adhesive and anti-biofilm agents against food pathogens [19–22].

Within this context, this work investigates the antimicrobial potential of celery (*Apium graveolens*) OR, limonene and their combination with rhamnolipids against planktonic cells and endospores of *B. cereus*.

## 2. Materials and Methods

### 2.1. Oils, Oleoresins and Biosurfactant

Oleoresin (8%) extracted from *Apium graveolens* seeds was kindly donated by Beraca Sabará S.A (Santa Bárbara d’Oeste, Brazil). Limonene (97% purity—Cutrale, Araraquara, Brazil) was kindly donated by Prof. André L. M. Porto. Rhamnolipid biosurfactant (90% purity) was acquired from AGAE Technologies (Corvallis, OR, USA).

### 2.2. Mixture Stock Solutions

OR or LN was mixed with propylene glycol (1:1 or 1:0.5 for the endospores germination experiment) before being added to the culture broth containing a final concentration of 0.02% of Tween 80 or 5000 µg/mL of RL. The final concentration of OR and LN in the stock solutions was 80,000 µg/mL, and such values were based on the maximum amount of propylene glycol determined not to affect the bacterial growth, which was previously defined as 12.5%.

The mixtures were homogenized by vortexing and filtered (0.45 µm). RL was diluted in culture broth and further sterilized by filtration (0.22 µm). The final concentration of RL stock solution was 50,000 µg/mL.

### 2.3. Microorganism

The *Bacillus cereus* ATCC 33,018 strain was stored at −20 °C on TSB (Tryptic Soy Broth—Himedia) supplemented with 6 g/L of yeast extract (TSYEB) and 20% (v/v) glycerol.

### 2.4. Identification of Celery Oleoresin Components by GC-MS

Fatty acid methyl esters (FAME) of celery OR (10 mg) were prepared under stirring using concentrated sulfuric acid (1 drop) and methanol (1 mL) at 60 °C for 30 min. After cooling, 2 mL of NaCl and 1 mL of n-hexane were added, and the solution was stirred and left undisturbed for phase separation. The upper phase was separated, and chromatography analysis was performed using a Shimadzu QP2010 (Shimadzu Corporation, Kyoto, Japan) system comprising an AOC-20 auto-sampler and gas chromatograph interfaced with a mass spectrometer (GC-MS QP2010 Plus) with a J&W Scientific DB-5MS (Folsom, CA, USA) (5% phenylmethylpolysiloxane) fused silica capillary column (30 cm × 0.25 mm i.d., 0.25 µm film thickness). The operation was carried out in electron impact mode at 70 eV. Helium (99.999%) was used as carrier gas at a constant flow of 0.99 mL/min, and an injec-

tion volume of 1  $\mu\text{L}$  was employed (split ratio of 1:10). The injector temperature was 250  $^{\circ}\text{C}$ , and the ion source temperature was 250  $^{\circ}\text{C}$ . The oven temperature was programmed from 50 (isothermal for 1 min), with an increase of 5  $^{\circ}\text{C}/\text{min.}$ , to 300  $^{\circ}\text{C}$ , ending with a 10-min isothermal period at 300  $^{\circ}\text{C}$ . Mass spectra were taken at 70 eV with a scan interval of 0.3 s and fragments of 40 to 500 Da [23]. The identification of the methyl esters was performed by computerized matching of the acquired mass spectra with those stored in mass spectral libraries of the equipment data system. For terpene analysis, a mixture of hydrocarbons ( $\text{C}_9\text{H}_{20}$ – $\text{C}_{18}\text{H}_{38}$ ) was injected under the conditions described above and identification of constituents was also performed by computerized matching and by comparison of the spectra obtained with those of the databank and considering the relative retention index (RRI) [24].

#### 2.5. Determination of Minimal Inhibitory Concentration (MIC) and Minimal Bactericidal Concentration (MBC)

Stock cultures of *B. cereus* were transferred to TSEYA (Tryptic Soy Agar supplemented with 6 g/L of yeast extract) and incubated for 24 h at 37  $^{\circ}\text{C}$ . Cells were transferred to 5 mL of TSEYB and incubated for an additional 24 h at 37  $^{\circ}\text{C}$ . An aliquot of 1 mL of cell suspension was transferred to 4 mL of fresh TSEYB and incubated for 3 h at 37  $^{\circ}\text{C}$ . Antimicrobial activity was performed on 96-well microplates using the micro-broth dilution technique based on Clinical and Laboratory Standards Institute [25] guidelines. Microplate wells were filled with 100  $\mu\text{L}$  of TSEYB followed by addition of 100  $\mu\text{L}$  of the tested mixture on the first column and a 2-fold serial dilution. After serial dilutions of the antimicrobials, 20  $\mu\text{L}$  of the standardized bacterial inoculum ( $10^7$  CFU/mL) was added to each well and the microplates were incubated at 37  $^{\circ}\text{C}$  for 24 h. After visual inspection, 10  $\mu\text{L}$  from the wells where no bacterial growth was observed was spotted on agar plates and incubated at 37  $^{\circ}\text{C}$  for 24 and 48 h. MBC was defined as the lowest MIC concentration where no viable growth was detected. Subsequently, 20  $\mu\text{L}$  of 0.1% tetrazolium bromide (MTT-Sigma Aldrich, St. Louis, MO, USA) solution was added to the wells for 1 h to confirm the presence or absence of growth. The MIC was defined as the lowest concentration of antimicrobial mixture that showed no change in the MTT original color.

#### 2.6. Time–Kill Assay

The bacterial growth in the presence of antimicrobials was evaluated using a time dependence assay as described by Verma, 2007 [26]. The tests were conducted in glass tubes filled with 5 mL of culture broth containing one of the antimicrobial mixtures (RL, OR + RL or LN + RL) at MIC and 16 $\times$  MIC concentration. Inoculum was prepared as described above (2.5) and 1 mL of standardized bacterial suspension was added to the tubes following incubation at 37  $^{\circ}\text{C}$ . At different time intervals, the number of viable cells was determined by the drop method [27]. Control tubes with no antimicrobial addition were also assessed.

#### 2.7. Endospore Germination Inhibition

Stock cultures of *B. cereus* were transferred to TSEYA and incubated for 24 h at 37  $^{\circ}\text{C}$ . After that, cells were transferred to a modified nutrient agar plate to favor sporulation (nutrient agar supplemented with 0.06 g/L  $\text{MgSO}_4$  and 0.35 g/L  $\text{KH}_2\text{PO}_4$ ) and incubated for 10 days at 37  $^{\circ}\text{C}$  [28].

Resulting cultures were suspended in sterilized distilled water, and the optical density was adjusted to 0.5 at 610 nm. This OD corresponds to  $3 \times 10^7$  cells/mL. Cell suspension was then heated to 75  $^{\circ}\text{C}$  for 20 min to assure only the endospores were present [28]. To confirm sporulation, malachite green staining was performed (2.8). An amount of 1 mL of endospore suspension was then added to 5 mL of TSEYB containing specific concentrations (16 $\times$  MIC and 32 $\times$  MIC) of RL, OR + RL or LN + RL. Samples were further incubated at 37  $^{\circ}\text{C}$  without agitation. At specific time points (0, 2, 6, 10 and 24 h), a 0.1 mL aliquot from each treatment was taken and added to 0.9 mL of saline solution (NaCl 0.86%) to perform a 10-fold serial dilution. Viable cells were counted by using the drop method [27].

Non-treated cells at time 0 were used as control for germination inhibition experiments. Log reduction was calculated for each specific time comparatively to the control.

The percentage of endospore germination inhibition and the log reduction were calculated using the following equations [28]:

$$\% \text{ endospore germination inhibition} = \frac{(CFU_{\text{control}} - CFU_{\text{treated}})}{CFU_{\text{control}}} \times 100, \quad (1)$$

$$\text{Log reduction} = \log_{10} \left( \frac{CFU_{\text{control}}}{CFU_{\text{treated}}} \right) \quad (2)$$

## 2.8. Endospore Staining

A 2 mL sample of culture broth was centrifuged at 10,000 rpm for 10 min. The supernatant was discarded, and cells were washed with 2 mL distilled water and centrifuged again. Supernatant was discarded and cell pellet was resuspended in 50 µL distilled water. Cellular suspension was then transferred to a glass slide, heat fixed and covered with malachite green solution (5%). Slides were passed through a flame for 5 min, paying attention to not boil the dye. The slide was then washed and stained with a safranin solution (2.5%) for 30 s [29,30]. After being air dried, the slides were observed by bright-field microscopy using immersion lens.

## 2.9. Statistics

MIC and MBC values were expressed as the mode of at least three independent replicates. All other data are represented as the mean  $\pm$  SD of three independent replicates. Analysis of variance (ANOVA) and the Tukey test were performed using OriginPro, version 8.5 (OriginLab Corporation, Northampton, MA, USA).

## 3. Results and Discussion

A preliminary screening using food-derived essential oils and oleoresins showed that *Apium graveolens* OR was efficient in controlling the growth of food pathogens such as *Listeria monocytogenes* and *Bacillus cereus* (data not shown). Based on such previous observations, we developed this new study to investigate, in detail, the antimicrobial activity demonstrated by the celery OR. Considering that one of the main difficulties in the application of EO and OR is their low water solubility, we also evaluated their combination with rhamnolipids since the amphiphilic nature of the biosurfactant may improve solubility and favor the delivery of hydrophobic compounds to cell targets. In addition, rhamnolipids also demonstrate antimicrobial activity against several Gram-positive food pathogens [31]; thus, we hypothesize that their combination with OR and/or EO can improve the antimicrobial potential of the natural compounds. *Bacillus cereus* was selected as our model study bacterium because of its importance as a food pathogen, along with its endospore-forming ability.

### 3.1. Analysis of Celery Oleoresin

To identify the main components present in the celery OR, a GC-MS analysis was performed and the results are shown in Table 1. Terpenes (limonene), sesquiterpenes ( $\beta$ -selinene), flavor (aromatic) compounds (3-buthylphthalide, sedanolide) and fatty acid esters were found as the major compounds. Although no reports using oleoresin are available for comparison, most studies in the literature identified terpenes and sesquiterpenes as the main active components of celery seed EO [32,33]. Therefore, we also investigated the antimicrobial activity of limonene (LN) since it is considered the major component of celery EO responsible for the antimicrobial activity against several pathogens [33,34].

**Table 1.** Chemical composition of celery (*Apium graveolens*) oleoresin.

	RT (min) <sup>1</sup>	Compound	% GC-MS	Exp RRI <sup>2</sup>	Lit RRI <sup>3</sup>
1	9.965	limonene	9.48	1039	1024
3	13.415	pentyl ciclohexa-1,3-diene	0.36	1158	1156
4	18.980	1-phenyl-1-pentanone	0.57	1358	1364
5	20.705	β-kariophyllene	0.23	1423	1417
6	22.455	β-selinene	3.11	1494	1489
7	22.615	α-selinene	0.77	1501	1498
8	23.190	methyl dodecanoate	0.06	1524	–
11	24.725	kariophyllene oxide	0.27	1586	1582
12	26.295	3-butylphthalide	5.67	1655	1647
13	26.415	β-eudesmol	0.86	1661	1649
14	26.730	3Z-butylidenephthalide	0.20	1675	1671
16	27.765	sedanenolide	7.41	1722	1719
17	27.915	neocnidilide	1.75	1728	1722
18	28.040	Z-ligustilide	0.34	1734	1734
19	29.980	methyl pentadecanoate	0.27	1815	–
23	31.435	methyl 7,10,13-hexadecatrienoate	0.28	1859	–
24	31.535	methyl 7-hexadecenoate	0.37	1862	–
25	31.600	methyl 9-hexadecenoate	0.87	1865	–
27	32.045	methyl hexadecanoate	8.23	1878	–
28	32.745	hexadecanoic acid	0.08	1900	–
31	33.985	methyl heptadecanoate	0.26	1947	–
35	35.280	methyl 9,12-octadecadienoate	11.68	1996	–
36	35.465	methyl 9-octadecenoate	30.50	2003	–
37	35.875	methyl octadecanoate	3.58	2019	–
38	36.115	9-octadecenoic acid	1.03	2028	–
40	38.880	methyl 11-eicosenoate	0.23	2134	–
42	39.375	methyl eicosanoate	0.66	2152	–
44	41.030	methyl heneicosanoate	0.08	2216	–
47	42.620	methyl docosanoate	0.56	2276	–
48	44.155	methyl tricosanoate	0.18	2334	–
52	45.630	methyl tetracosanoate	0.58	2391	–
53	46.785	trans-squalene	0.14	2435	–
54	47.065	methyl pentacosanoate	0.08	2445	–
57	48.440	methyl hexadocosanoate	0.40	2498	–
60	51.070	methyl octacosanoate	0.49	2598	–
61	52.705	β-stigmasterol	0.51	2660	–
63	54.045	methyl triacontanoate	0.16	2711	–
		Identified	92.30		
		Non-identified	7.70		
		Fatty esters	60.63		
		Total	100.00		

<sup>1</sup> Retention time in minutes. <sup>2</sup> Experimental relative retention index. <sup>3</sup> Literature relative retention index.

### 3.2. Minimal Inhibitory Concentration (MIC) and Minimal Bactericidal Concentration (MBC)

Vegetative *B. cereus* cells were treated with solutions containing RL, OR, and LN, alone or in combination, to determine the minimal concentrations necessary to inhibit growth and to kill the bacterial cells after 24 and 48 h of exposure (Table 2).

**Table 2.** Antimicrobial activity of the tested compounds against *B. cereus* vegetative cells.

Compound	MIC (µg/mL)	MBC 24h (µg/mL)	MBC 48h (µg/mL)
RL	98	1562.5	1562.5
OR	40,000	>40,000	>40,000
OR + RL	2500 OR + 156.3 RL	20,000 OR + 1250 RL	>40,000 OR + 2500 RL
LN	2500	>40,000	>40,000
LN + RL	1250 LN + 78.1 RL	20,000 LN + 1250 RL	20,000 LN + 1250 RL



When considered alone, RL was able to inhibit cell growth with concentrations as low as 98 µg/mL, and cell death was obtained with 1562.5 µg/mL. OR showed the highest MIC value of the tested compounds. It was not possible to determine the MBC of either oil based on the concentrations tested. When combined with RL, on the other hand, both OR and LN showed a reduction in the MIC and exhibited MBC. The mixture containing OR + RL was able to inhibit cell growth at 2500 + 156.3 µg/mL and kill cells at 20,000 + 1250 µg/mL (24 h), while the LN + RL mixture had an MIC of 1250 + 78.1 µg/mL and an MBC of 20,000 + 1250 µg/mL. These results suggest that, when combined with RL, the antimicrobial effect of OR and LN is enhanced. By contrast, it is also possible to assume that OR may have an inhibitory effect on RL, since Table 2 shows that more RL is needed to reach the MIC when combined with OR.

### 3.3. Time-Kill Assays

A new set of experiments was conducted using the previously determined MIC and MBC values, in order to study the bacterial behavior in the presence of the antimicrobials during the time-defined intervals. Since no MBC was observed even for OR and LN alone, they were not considered for this assay.


The growth of *B. cereus* cells in the presence of RL, OR + RL and LN + RL revealed that the values initially designated as MBC are not capable of eradicating the cell population (Figure 1), although they were 16 times higher than the MIC concentration. Rhamnolipid treatment reduced the viable cell population by around 4 log and the population was maintained at this level after 24 h (Figure 1a). The combination of RL with LN and OR improved their antimicrobial efficacy, corroborating the data shown in Table 2. For LN + RL, there is no apparent difference between the MIC and the 16x MIC values (Figure 1c). On the contrary, for OR + RL, it is possible to observe that the increase in the mixture concentration caused a decrease in cell survival (Figure 1b). This fact can be explained by the increasing concentration of active compounds present in OR, such as terpenes and sesquiterpenes (Table 1).

It is also worth noting that all curves show roughly a 2 log drop in CFU/mL after only 2 h of contact. We speculate that this behavior is caused by the endospore-forming ability of *B. cereus* under adverse situations. The initial drop in cell counts means that some cells died within this period of time, while the remaining cells switched on into endospores. To confirm this hypothesis, test samples were stained, and microscopy revealed the presence of endospores (Figure 2).

The inconsistency in MBC values described in Table 2 can be explained by the count method utilized. We observed that during serial dilutions to perform the viable count, the first undiluted sample did not show growth after incubation; however, in the subsequent decimal dilutions of the same sample, several colonies were observed (Supplementary Materials Figure S1). This fact is caused by the antimicrobial dilution factor; thus, in the first sample (without dilution), the drug concentration was able to avoid cell growth, while after dilution, the remaining cells (or endospores) could grow. As the MBC was performed after 24 and 48 h of incubation by plating the original sample in a fresh plate without dilution, the remaining antimicrobial concentration was sufficient to inhibit the growth or, possibly, endospore germination. In addition, as shown in Table 2, the MBC of the OR + RL mixture was increased after 48 h, corroborating the statements described above.

Antimicrobial activity against *Bacillus cereus* was previously reported for RL [31,35], limonene [36,37] and celery EO [35]; however, there are no reports regarding the use of celery oleoresin and/or EO components associated with RL. Similar studies using the particular compounds showed that celery seed EO (100% v/v) presented an inhibition zone of 33 mm against *B. cereus* cells [32]. In another study, 250 µg/mL of limonene was able to inhibit *B. cereus* growth; however, no MBC was observed [37]. In addition, an inhibition zone of 6.3–6.7 mm was reported when 420 µg of limonene was tested against two strains of *B. cereus* [36]. Rhamnolipids showed an MIC of 19.5 µg/mL [31] and an inhibition zone ranging from 13 to 22 mm against *B. cereus* strains [35]. These examples point out that


differences regarding the strains' sensitivity, methodology, purity of the compounds and their formulations can influence the antimicrobial activity; thus, it is difficult to compare our results with the ones available in the literature.



**Figure 1.** Time–kill curves for *B. cereus* with (a) rhamnolipid (RL), (b) celery oleoresin (OR) + RL, and (c) limonene (LN) + RL. Lines show the log CFU/mL for the control (black), minimum inhibitory concentration (MIC) (red) and 16× MIC (blue). Error bars show the standard deviation of at least three independent replicates.

Considering that the mechanism involved in the antimicrobial activity of both RL and LN/EO is related to disturbance in the permeability/integrity of the bacterial cell membrane [34,38–40], we can suggest that the combination of a biosurfactant and active hydrophobic compounds may in fact increase their solubility, favoring interaction with the cells.

As observed, the presence of RL and its combination with OR and LN can inhibit the growth of vegetative cells but also induces *B. cereus* sporulation (Figure 2). It was also demonstrated that under the presence of the antimicrobials, the population was maintained relatively constant, suggesting they might also inhibit endospore germination. To test this hypothesis, further germination inhibition assays were conducted.




**Figure 2.** Images of endospore staining of (a) control, (b) OR + RL MIC 14h and (c) OR + RL MIC 24h treatments. Endospores are visible on images (b) and (c) as an uncolored region inside the cells. Some endospores are indicated by arrows. Magnification 1000 $\times$ .

### 3.4. Endospore Inhibition


The endospores of *B. cereus* were treated with RL and its respective mixtures with celery OR and LN. The percentage of germination inhibition (CFU) and log reduction in outgrowth were calculated after different times of exposure. RL showed a reduction around 93% in germination after 24 h. Additionally, the increase in RL concentration and time of exposure did not significantly increase the inhibition of endospore germination (Figure 3). A maximum of a 3.6 log reduction in endospore outgrowth was also observed after 6 h compared to the control at the same time (Figure 4a). When RL was mixed with OR, the maximum inhibition percentage (73%) and log reduction (2.8) were obtained for the highest concentration tested. As also shown in Figure 1b, the increase in concentration improved the activity of the OR + RL mixture, probably due to the increase in the active compounds present in OR (Table 1). As observed in vegetative cells, OR also seems to inhibit the effect of RL on endospores, since when both antimicrobials are combined, lower endospore germination inhibition values are displayed compared to RL alone (Figure 1).

The combination of LN + RL demonstrated the highest endospore inhibition values after 24 h of exposure, achieving a maximum of 98% inhibition. An important decrease in cell counts over time was also observed, reaching a 3.3 log reduction after 24 h, as shown in Figure 4c. As previously discussed, one of the main components of OR is LN (around 9.5%) and it may be responsible (at least in part) for OR's antimicrobial effect. This could explain why, when LN is in its purified form, it shows higher antimicrobial potential compared with OR samples.





**Figure 3.** Percentage of *B. cereus* endospore germination inhibition of RL, OR + RL, and LN + RL treatments. The concentrations used for this experiment were 16 $\times$  and 32 $\times$  MIC. Error bars represent the standard deviation of at least three independent replicates. For each time of exposure, treatments with the same letters did not differ significantly ( $p < 0.05$ ).



**Figure 4.** Log reduction in endospore germination after treatment with (a) RL, (b) OR + RL, and (c) LN + RL. Lines show the log CFU/mL for the control (black), 16 $\times$  MIC (red) and 32 $\times$  MIC (blue). Error bars show the standard deviation of at least three independent replicates.

Regardless of RL and EO, antimicrobial action against vegetative *B. cereus* cells is well documented, though reports on their activity against endospores are scarce in the literature. Huang et al., 2007 [41], evaluated the potential of a lipopeptide biosurfactant produced by *B. subtilis* to inactivate *B. cereus* endospores. The lipopeptide displayed an MIC of 156.25  $\mu\text{g/mL}$ , and the authors proposed the combination with heat to reduce endospore germination. A 2 log reduction was attained after treatment of *B. cereus* endospores for 7.6 h at 29.6  $^{\circ}\text{C}$  with 3.46 mg/mL of the lipopeptide. A spore coat disruption was also observed,

and the authors proposed that the surfactant nature of the compound could favor the binding to lipoproteins and internalization of the biosurfactant, resulting in coat damage.

The increasing in lipophilic character of organic acids and alcohols enhanced the inhibition of *B. cereus* spore germination. The accumulation of such compounds in the inner membrane was correlated with the length of their carbon chains and consequent endospore germination inhibition [42].

Further reports have demonstrated that the amphiphilic nature of surfactants, more precisely, their hydrophilic–lipophilic balance (HBL), can be correlated with endospore inactivation potential. The hydrophobic compounds citral, *p*-cymene and bornyl acetate were more effective at inhibiting *B. subtilis* endospores than hydrophilic compounds such 2,3 dihydrobenzofuran and  $\beta$ -pinene [43]. The average HBL value of the compounds causing a significant 1–2 log reduction in endospores was 9.3, which is similar to several chemical surfactants. According to the authors, the structural proteins of the spore coat can be changed due to binding of their polar and apolar groups to hydrophilic and hydrophobic moieties of the surfactants.

Green tea polyphenol samples including a crude extract, epigallocatechin-3-gallate and their respective lipophilic derivatives were reported to prevent endospore germination in different *Bacillus* species. The four types of polyphenols inhibited *B. cereus* spore germination by 94–100% with a log reduction ranging from 1.27 to 3.0. A disruption in the *B. cereus* spore coat, changing in coat morphology and spore agglutination, was observed after the polyphenol treatment [28].

Although the exact mechanisms involved in spore inhibition are unknown, it is a consensus that the lipophilic nature of chemical compounds influences their activity. Since the endospore surface is hydrophobic, molecules with this character might easily interact with several targets present in the inner/outer coats, cortex, membrane and core [44].

The HBL of RL is dependent (among other factors) on the purity and composition of homologous compounds present in the mixture, and several different values have been reported in the literature, ranging from 6.5 to 24 [45,46]. Thus, the amphiphilic character presented by the rhamnolipids may account for the results observed in our work. The combination of RL with the hydrophobic compounds LN and OR can improve their solubility and consequent interaction with endospore structures, as also proposed for vegetative cells. Our results demonstrate that RL potentially inhibits *B. cereus* vegetative cells and endospore germination and also enhances the antimicrobial action of the oil-derived compounds tested. Further studies should be conducted to understand the mechanism involved in the endospore inhibition by the compounds.

To our knowledge, this is the first report on using RL and its mixture with LN and celery OR to inhibit *B. cereus* endospores. Combinations of such compounds with physical methods (heat, radiation, pressure) or the development of micro/nanoemulsions exploring RL and OR/EO components may result in innovative strategies to control this important food pathogen.

**Supplementary Materials:** The following are available online at <https://www.mdpi.com/2304-8158/10/2/455/s1>, Figure S1: Images of the Petri dishes used for viable cell counting. On the undiluted sample, growth after incubation was not observed; however, as the dilution of the sample occurred, several colonies were observed. (a) LN + RL; (b) RL. Numbers on plates represent how many times the samples were 10-fold diluted.

**Author Contributions:** Conceptualization, M.N.; methodology, D.M.D.M. and P.d.C.B.; formal analysis, D.M.D.M. and P.d.C.B.; investigation, D.M.D.M. and P.d.C.B.; writing—original draft preparation, P.d.C.B. and M.N.; writing—review and editing, P.d.C.B. and M.N.; supervision, project administration and funding acquisition, M.N. All authors have read and agreed to the published version of the manuscript.

**Funding:** The authors would like to thank COORDENAÇÃO DE APERFEIÇOAMENTO DE PESSOAL DE NÍVEL SUPERIOR (CAPES)—Grant n° 001, for PB's and DM's scholarships and FUN-

DAÇÃO DE AMPARO À PESQUISA DO ESTADO DE SÃO PAULO (FAPESP)—Grant n° 2019/14405-9, for financial support.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author.

**Acknowledgments:** The authors would like to thank Beraca Sabará S.A. for the celery oleoresin donation. The authors also are grateful to André L. M. Porto for the limonene donation and GC-MS facility and support.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## References

1. World Health Organization. Available online: <http://www.who.int/mediacentre/news/releases/2015/foodborne-disease-estimates/en/> (accessed on 5 December 2020).
2. World Health Organization. Fact Sheets: Food Safety. Available online: <https://www.who.int/news-room/fact-sheets/detail/food-safety> (accessed on 3 December 2020).
3. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Vigilância das Doenças Transmissíveis. Surtos de Doenças Transmitidas por Alimentos no Brasil. Available online: <http://portal.arquivos.saude.gov.br/images/pdf/2017/maio/29/Apresentacao-Surtos-DTA-2017.pdf> (accessed on 3 December 2020).
4. Carroll, L.M.; Wiedmann, M.; Mukherjee, M.; Nicholas, D.C.; Mingle, L.A.; Dumas, N.B.; Cole, J.A.; Kovac, J. Characterization of Emetic and Diarrheal *Bacillus cereus* Strains From a 2016 Foodborne Outbreak Using Whole-Genome Sequencing: Addressing the Microbiological, Epidemiological, and Bioinformatic Challenges. *Front. Microbiol.* **2019**, *10*, 144. [CrossRef]
5. Glasset, B.; Herbin, S.; Guillier, L.; Cadel-Six, S.; Vignaud, M.-L.; Grout, J.; Pairaud, S.; Michel, V.; Hennekinne, J.-A.; RamaRao, N.; et al. *Bacillus cereus*-induced food-borne outbreaks in France, 2007 to 2014: Epidemiology and genetic characterisation. *Eurosurveillance* **2016**, *21*. [CrossRef]
6. Carrocho, M.; Morales, P.; Ferreira, I.C. Natural food additives: Quo vadis? *Trends Food Sci. Technol.* **2015**, *45*, 284–295. [CrossRef]
7. Gray, J.A.; Chandry, P.S.; Kaur, M.; Kocharunchitt, C.; Bowman, J.P.; Fox, E.M. Novel Biocontrol Methods for *Listeria monocytogenes* Biofilms in Food Production Facilities. *Front. Microbiol.* **2018**, *9*, 1–12. [CrossRef]
8. El Asbahani, A.; Miladi, K.; Badri, W.; Sala, M.; Addi, E.H.A.; Casabianca, H.; El Mousadik, A.; Hartmann, D.; Jilale, A.; Renaud, F.N.R.; et al. Essential oils: From extraction to encapsulation. *Int. J. Pharm.* **2015**, *483*, 220–243. [CrossRef]
9. Swamy, M.K.; Akhtar, M.S.; Sinniah, U.R. Antimicrobial Properties of Plant Essential Oils against Human Pathogens and Their Mode of Action: An Updated Review. *Evid.-Based Complement. Altern. Med.* **2016**, *2016*, 1–21. [CrossRef]
10. Hyldgaard, M.; Mygind, T.; Meyer, R.L. Essential Oils in Food Preservation: Mode of Action, Synergies, and Interactions with Food Matrix Components. *Front. Microbiol.* **2012**, *3*, 12. [CrossRef]
11. Okunowo, W.O.; Oyedele, O.; Afolabi, L.O.; Matanmi, E. Essential Oil of Grape Fruit (*Citrus paradisi*) Peels and Its Antimicrobial Activities. *Am. J. Plant Sci.* **2013**, *4*, 1–9. [CrossRef]
12. Gomes, M.S.; Cardoso, M.D.G.; Soares, M.J.; Batista, L.R.; Machado, S.M.F.; Andrade, M.A.; De Azeredo, C.M.O.; Resende, J.M.V.; Rodrigues, L.M.A. Use of Essential Oils of the Genus *Citrus* as Biocidal Agents. *Am. J. Plant Sci.* **2014**, *5*, 299–305. [CrossRef]
13. Andrade, B.F.M.T.; Barbosa, L.N.; Probst, I.D.S.; Júnior, A.F. Antimicrobial activity of essential oils. *J. Essent. Oil Res.* **2014**, *26*, 34–40. [CrossRef]
14. Sowbhagya, H.B. Chemistry, Technology, and Nutraceutical Functions of Celery (*Apium graveolens* L.): An Overview. *Crit. Rev. Food Sci. Nutr.* **2013**, *54*, 389–398. [CrossRef] [PubMed]
15. Arshad, H.; Ali, T.M.; Abbas, T.; Hasnain, A. Effect of Microencapsulation on Antimicrobial and Antioxidant Activity of Nutmeg Oleoresin Using Mixtures of Gum Arabic, OSA, and Native Sorghum Starch. *Starch-Stärke* **2018**, *70*, 1–12. [CrossRef]

16. King, K. Packaging and storage of herbs and spices. In *Handbook of Herbs and Spices*; Peter, K.V., Ed.; Woodhead Publishing Series in Food Science, Technology and Nutrition; Woodhead Publishing: Cambridge, UK, 2006; Volume 3, pp. 86–102, ISBN 9781845690175. Available online: <http://www.sciencedirect.com/science/article/pii/B9781845690175500051> (accessed on 26 November 2020).
17. Lucera, A.; Costa, C.; Conte, A.; Del Nobile, M.A. Food applications of natural antimicrobial compounds. *Front. Microbiol.* **2012**, *3*, 287. [CrossRef] [PubMed]
18. Nitschke, M.; E Silva, S.S. Recent food applications of microbial surfactants. *Crit. Rev. Food Sci. Nutr.* **2018**, *58*, 631–638. [CrossRef]
19. Magalhães, L.; Nitschke, M. Antimicrobial activity of rhamnolipids against *Listeria monocytogenes* and their synergistic interaction with nisin. *Food Control* **2013**, *29*, 138–142. [CrossRef]
20. De Araujo, L.V.; Guimarães, C.R.; Marquita, R.L.D.S.; Santiago, V.M.; De Souza, M.P.; Nitschke, M.; Freire, D.M.G. Rhamnolipid and surfactin: Anti-adhesion/antibiofilm and antimicrobial effects. *Food Control* **2016**, *63*, 171–178. [CrossRef]
21. E Silva, S.; Carvalho, J.; Aires, C.; Nitschke, M. Disruption of *Staphylococcus aureus* biofilms using rhamnolipid biosurfactants. *J. Dairy Sci.* **2017**, *100*, 7864–7873. [CrossRef]
22. Clements, T.; Ndlovu, T.; Khan, S.; Khan, W. Biosurfactants produced by *Serratia* species: Classification, biosynthesis, production and application. *Appl. Microbiol. Biotechnol.* **2019**, *103*, 589–602. [CrossRef]
23. Marangon, C.A.; Martins, V.C.; Leite, P.M.; Santos, D.A.; Nitschke, M.; Plepis, A.M.G. Chitosan/gelatin/copaiba oil emulsion formulation and its potential on controlling the growth of pathogenic bacteria. *Ind. Crop. Prod.* **2017**, *99*, 163–171. [CrossRef]
24. Adams, R.P. *Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry*, 4th ed.; Allured Publishing Corporation: Carol Stream, IL, USA, 2007; ISBN 978-1-932633-21-4.
25. CLSI. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard*, 9th ed.; CLSI Document M07-A9; CLSI: Wayne, PA, USA, 2012; ISBN 1-56238-784-7.
26. Verma, P. Methods for determining bactericidal activity and antimicrobial interactions: Synergy testing, time-kill curves, and population analysis. In *Antimicrobial Susceptibility Testing Protocols*; Schwalbe, R., Steele-Moore, L., Goodwin, A.C., Eds.; Taylor & Francis: Boca Raton, FL, USA, 2007; pp. 275–299, ISBN 9780429132346.
27. Miles, A.A.; Misra, S.S.; Irwin, J.O. The estimation of the bactericidal power of the blood. *Epidemiol. Infect.* **1938**, *38*, 732–749. [CrossRef]
28. Ali, B.; Lee, L.H.; Laskar, N.; Shaikh, N.; Tahir, H.; Hsu, S.D.; Newby, R.; Valsechi-Diaz, J.; Chu, T. Modified Green Tea Polyphenols, EGCG-S and LTP, Inhibit Endospore in Three *Bacillus* spp. *Adv. Microbiol.* **2017**, *7*, 175–187. [CrossRef]
29. Schaeffer, A.B.; Fulton, M.D. A simplified method of staining endospores. *Science* **1933**, *77*, 194. [CrossRef] [PubMed]
30. Hamouda, T.; Shih, A.Y.; Baker, J.R., Jr. A rapid staining technique for the detection of the initiation of germination of bacterial spores. *Let. Appl. Microbiol.* **2002**, *34*, 86–90. [CrossRef]
31. Ferreira, J.D.F.; Vieira, E.A.; Nitschke, M. The antibacterial activity of rhamnolipid biosurfactant is pH dependent. *Food Res. Int.* **2019**, *116*, 737–744. [CrossRef] [PubMed]
32. Hassanen, N.H.M.; Eissa, A.M.F.; Hafez, S.A.M. Antioxidant and antimicrobial activity of celery (*Apium graveolens*) and coriander (*Coriandrum sativum*) herb and seed essential oils. *Int. J. Curr. Microbiol. Appl. Sci.* **2015**, *4*, 284–296.
33. Dąbrowska, J.A.; Kunicka-Styczyńska, A.; Śmigielski, K.B. Biological, chemical, and aroma profiles of essential oil from waste celery seeds (*Apium graveolens* L.). *J. Essent. Oil Res.* **2020**, *32*, 308–315. [CrossRef]
34. Espina, L.; Gelaw, T.K.; De Lamo-Castellví, S.; Pagán, R.; García-Gonzalo, D. Mechanism of Bacterial Inactivation by (+)-Limonene and Its Potential Use in Food Preservation Combined Processes. *PLoS ONE* **2013**, *8*, e56769. [CrossRef]
35. Ndlovu, T.; Rautenbach, M.; Vosloo, J.A.; Khan, S.; Khan, W. Characterisation and antimicrobial activity of biosurfactant extracts produced by *Bacillus amyloliquefaciens* and *Pseudomonas aeruginosa* isolated from a wastewater treatment plant. *AMB Express* **2017**, *7*, 1–19. [CrossRef]
36. De Martino, L.; De Feo, V.; Fratianni, F.; Nazzaro, F. Chemistry, Antioxidant, Antibacterial and Antifungal Activities of Volatile Oils and their Components. *Nat. Prod. Commun.* **2009**, *4*, 1741–1750. [CrossRef] [PubMed]
37. Guimarães, A.C.; Meireles, L.M.; Lemos, M.F.; Guimarães, M.C.C.; Endringer, D.C.; Fronza, M.; Scherer, R. Antibacterial Activity of Terpenes and Terpenoids Present in Essential Oils. *Molecules* **2019**, *24*, 2471. [CrossRef] [PubMed]
38. Ortiz, A.; Teruel, J.A.; Espuny, M.J.; Marqués, A.; Manresa, Á.; Aranda, F.J. Effects of dirhamnolipid on the structural properties of phosphatidylcholine membranes. *Int. J. Pharm.* **2006**, *325*, 99–107. [CrossRef]
39. Sotirova, A.V.; Spasova, D.I.; Galabova, D.N.; Karpenko, E.; Shulga, A. Rhamnolipid-Biosurfactant Permeabilizing Effects on Gram-Positive and Gram-Negative Bacterial Strains. *Curr. Microbiol.* **2008**, *56*, 639–644. [CrossRef] [PubMed]
40. Li, Z.-H.; Cai, M.; Liu, Y.-S.; Sun, P.; Luo, S.-L. Antibacterial Activity and Mechanisms of Essential Oil from *Citrus medica* L. var. *sarcodactylis*. *Molecules* **2019**, *24*, 1577. [CrossRef] [PubMed]
41. Huang, X.; Lu, Z.; Bie, X.; Lü, F.; Zhao, H.; Yang, S. Optimization of inactivation of endospores of *Bacillus cereus* by antimicrobial lipopeptides from *Bacillus subtilis* fmbj strains using a response surface method. *Appl. Microbiol. Biotechnol.* **2006**, *74*, 454–461. [CrossRef] [PubMed]
42. Van Melis, C.; Almeida, C.B.; Kort, R.; Groot, M.N.; Abee, T. Germination inhibition of *Bacillus cereus* spores: Impact of the lipophilic character of inhibiting compounds. *Int. J. Food Microbiol.* **2012**, *160*, 124–130. [CrossRef]
43. Cho, W.-I.; Cheigh, C.-I.; Hwang, H.-J.; Chung, M.-S. Sporocidal Activities of Various Surfactant Components against *Bacillus subtilis* Spores. *J. Food Prot.* **2015**, *78*, 1221–1225. [CrossRef]

- 
44. Cho, W.-I.; Chung, M.-S. Bacillus spores: A review of their properties and inactivation processing technologies. *Food Sci. Biotechnol.* **2020**, *29*, 1447–1461. [[CrossRef](#)]
  45. De Rienzo, M.D.; Stevenson, P.; Marchant, R.; Banat, I. Pseudomonas aeruginosa biofilm disruption using microbial surfactants. *J. Appl. Microbiol.* **2016**, *120*, 868–876. [[CrossRef](#)] [[PubMed](#)]
  46. Xie, Y.; Li, Y.; Ye, R. Effect of Alcohols on the Phase Behavior of Microemulsions Formed by a Biosurfactant—Rhamnolipid. *J. Dispers. Sci. Technol.* **2005**, *26*, 455–461. [[CrossRef](#)]