

Article

The Effect of Guisangyou Tea on Abnormal Lipid Metabolism in Mice Induced by High-Fat Diet

Yan Zhu ^{1,†}, Xianghui Zhou ^{2,†}, Nan Ling ², Qiming Yu ^{2,*}, Huijuan Wang ¹ and Qizhen Du ^{1,*}

¹ College of Food and Health Sciences, Zhejiang A & F University, Hangzhou 311300, China; yueliangyan@163.com (Y.Z.); wanghuijuan1016@126.com (H.W.)

² College of Public Health, Guilin Medical University, Guilin 541100, China; ZHOUXH_0472@163.com (X.Z.); lingnan0118@163.com (N.L.)

* Correspondence: qm_yu19@glmc.edu.cn (Q.Y.); qizhendu@163.com (Q.D.)

† These authors contributed equally to this work.

Supplementary Materials

Table S1. The nutrient composition of normal diet and high-fat diet.

Ingredient	Normal diet (g)	High-fat diet (g)
Corn starch	397.5	227.5
Casein	200	200
Dyetrose	132	132
Lard	0	170
Sucrose	100	100
Soybean oil	70	70
α -cellulose	50	50
Minerals mixture ^a	35	35
Vitamins mixture ^a	10	10
DL-Methionine	3	3
Choline bitartrate	2.5	2.5
Total weight	1000	1000

^a Minerals and Vitamins mixture according to AIN-93 diet (Teklad, Madison, WI, USA)

Table S2. The mouse organ indexes of the control group (CON), high fat diet group (HFD) and high fat diet-WE treatment group (WE)

Group	Organ index/%			
	Liver	Kidney	Spleen	Heart
CON	3.913±0.143 ^b	1.223±0.054 ^a	0.341±0.033 ^a	0.461±0.025 ^a
HFD	4.928±0.135 ^a	1.226±0.128 ^a	0.322±0.025 ^a	0.440±0.025 ^a
WE	4.379±0.164 ^b	1.218±0.138 ^a	0.352±0.034 ^a	0.454±0.024 ^a

Significant differences between groups are denoted by different letters ($P < 0.05$).