



Occurrence of Ochratoxin A in Coffee: Threads and Solutions—A Mini-Review

Ana Lúcia Leitão 1,2

Review

- ¹ Departamento de Ciências e Tecnologia da Biomassa, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, Campus da Caparica, 2829-516 Caparica, Portugal; aldl@fct.unl.pt
- ² MEtRICs, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Campus da Caparica, 2829-516 Caparica, Portugal

Received: 29 December 2018; Accepted: 26 April 2019; Published: 8 May 2019



Abstract: Ochratoxin A (OTA) is a widespread bioactive extrolite from secondary metabolism of fungi which presence in foods like coffee is of public health concern, particularly for heavy drinkers. Coffee is one of the most consumed and appreciated non-alcoholic beverage in the world. Its production from the plantation to the coffee cup involves several steps that would determine the final concentration of OTA in the beverage. This review gives an overview of OTA contamination in roasted coffee beans in different countries and mitigation strategies for OTA reduction.

Keywords: ochratoxin A; coffee; mycotoxin; threads; treatment

1. Coffee as a Unique Beverage

Coffee is one of the most popular beverages nowadays all over the world, with an annual global production of 10.4×10^6 tons and it plays an important cultural and economic role [1]. It contains more than 1500 chemical components, being responsible for sensorial and functional properties of the coffee beverage, the second most valuable legally traded commodity [2–4]. A recent study by Samoggia et al. (2018) indicated that consumers' preferences for a specific coffee type mainly depended on sensory aspects like taste and aroma and functional motives to improve alertness and physical performance, beside habits and traditions of the consumer [5]. Coffee can be consumed as a hot or cold beverage, usually as the result of an infusion of properly processed coffee beans, being consumed per se as part or as the last component of meals or as an ingredient in snacks. It is, generally, produced from two biologically dissimilar species the Coffea arabica and Coffea canephora var. robusta, representing 60% and 40%, respectively, of world production [6]. Coffee from Coffea arabica (Arabica) and Coffea canephora var. robusta (Robusta) have several differences not only in terms of botanical, genetic, and morphological features of bean but also in their chemical composition and environmental conditions [7]. For example, while Arabica is adapted to lower temperatures and reduced humidity characteristics of high altitudes, Robusta is cultivated from sea level up to 1000 m; caffeine concentration endogenously present in the green coffee extracts of Robusta, often, is double that of Arabica, being dependent also of geographical origin and producing a drink with pronounced bitterness [6,7]. The tocopherols contents of Arabica are higher than Robusta and while 90% of tocopherols remain after roasting in the case of Arabica, a degradation of approximately 25% is observed for β -tocopherol in Robusta [8]. Recently, it was reported that the roasting procedure affected differentially coffee varieties in what concerns to antimutagenic activity, while no effect was observed in Robusta varieties, roasting enhances time-dependently the antimutagenicity of the Brazil variety (Arabica) [9].

Coffee is a unique beverage with healthy properties that make it a functional food due to its content in chlorogenic acids, caffeine, trigonelline, diterpenes and tocopherols present in raw

beans. Therefore, the coffee extracts exhibited free radical scavenging and induced detoxification enzymes [9,10]. Additionally, beneficial effects of coffee such as anti-mutagen, inductor of the activation of DNA repair, as well as in the decrease of mortality from neurological and cardiovascular diseases, and diabetes type II has been reported [10,11]. Despite the literature support the role of coffee as having an immunomodulatory action, it seems to be disease-specific [11].

Recently, a study investigated the hypothesis that habitual coffee consumption is associated with lower risk of frailty, physical function impairment, and disability in women with aged 60 years or older [12]. The conclusion was that habitual coffee consumption of two or more cup of coffee/day, in older people, was not associated with increased risk of functional impairment, and it can be positive in the case of individuals with diabetes, hypertension or obesity [12].

2. Ochratoxin A as a Coffee Contaminant

2.1. Ochratoxin A

The mycotoxin ochratoxin A (OTA) is a low molecular weight ubiquitous secondary metabolite, a weak organic acid consisting of an amino acid phenylalanine and a dihydro-isocoumarin linked by a peptide bond (Table 1).

		Reference
Name	N-[(5-chloro-3,4-dihydro-8-hydroxy-3-methyl-1-oxo-1 <i>H-</i> 2-benzopyran-7-yl)carbonyl]-L-phenylalanine	
Chemical structure		
Formula	$C_{20}H_{18}CINO_6$	[13]
Molecular weight	tolecular weight $403.8 \text{ g} \cdot \text{mol}^{-1}$	
	168–173 °C (drying for 1 hour at 60 °C)	[14]
Melting point	159 °C (recrystallized from benzene-hexane)	[15]
	169 °C (recrystallized from xylene)	[16]
$[\alpha]^{21}D$	-46.8° ($c = 2.65$ mmol/L in chloroform)	[14]
[α] ²⁰ D	-118° ($c = 1.1$ mmol/L in chloroform)	[16]
	214 nm ($\varepsilon = 37.2 \times 10^{-3} \text{ L} \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$)	
λ_{max}	282 nm ($\varepsilon = 0.89 \times 10^{-3} \mathrm{L \cdot mol^{-1} \cdot cm^{-1}}$)	[14]
	$332 \text{ nm} (\varepsilon = 6.33 \times 10^{-3} \text{ L} \cdot \text{mo}^{-1} \cdot \text{cm}^{-1})$	[1]]
	(c = 0.0281 mmol/L in methanol)	
рКа	4.2–4.4 and 7.0–7.3	[17]

Table 1. Structure, formula and physical characteristics of ochratoxin A (OTA).

OTA is a white crystalline, odorless, solid compound with poor aqueous solubility, but moderately soluble in chloroform, methanol, ethanol and xylene. It is heat-stable, being described as the most common member in group of ochratoxins [18,19]. The group of ochratoxins is principally formed by OTA, ochratoxin B (OTB) and ochratoxin C (OTC). From a chemical perspective, while OTA is chlorinated, ochratoxin B is not chlorinated, and ochratoxin C is the ethyl ester of OTA [18]. Although OTA appear to be overall the most toxic, from recent investigations a different and more detailed picture have resulted with cell models showing in lower or similar toxicity of OTB compared to OTA; while there seems no doubt that OTB is less toxic in vivo compared to OTA, similar acutely cytotoxic in vitro is observed. Meanwhile OTC seems to be equally acutely toxic in vivo and in vitro compared

to OTA [20]. Nowadays, based on the metabolism of animals and humans, the number of molecules belonging to this group has increased substantially, especially on OTA derivatives. Examples of this are hydroxylated, conjugated or metabolites where phenylalanine moiety is absent [21]. Despite its stability, OTA can be transformed mainly to metabolites, which were considered to have low or no toxicity (Figure 1) [22–25]. Today's knowledge on comparative toxicity of OTA and its metabolites shows that ochratoxin α (OT α) is approximately 100 fold less toxic than OTA; 10-hydroxyochratoxin A (10-OH OTA) is non-genotoxic, and conjugated ochratoxin forms are, as so far, not toxic; meanwhile the 4-hydroxyochratoxin A (4-OH OTA) shows similar toxicity, and lactone-opened OTA (OP-OTA) could be even more toxic than OTA [20].



Figure 1. Structures of ochratoxin A (**OTA**) and transformation bioproducts ochratoxin B (**OTB**), 10-hydroxyochratoxin A (**10-OH-OTA**), ochratoxin α (**OT** α), 14-decarboxy-ochratoxin A (**14-DC-OTA**), lactone-opened OTA (**OP-OA**), 4-hyroxyochratoxin A (**4-OH-OTA**) and OTA conjugated metabolites adapted from [23–25].

This mycotoxin was reported for the first time in 1965 [16] and nine years later it was described in coffee [26]. Although humans can be exposed to OTA by inhalation or dermal contact, dietary intake is the principal source of OTA for humans because it is found in a variety of foods, like for example; maize, wheat, rice, sorghum, barley, oats, rye, bread, flour, pasta, infant cereals, grapes, apples, peaches, pears, strawberries, oranges, figs, mangoes, tomatoes, wine, watermelons, nuts, sesame seeds, rapeseed, spice, coffee beans, cocoa, peanuts, chickpeas, soybeans, milk and milk-based baby formulae, cheese, eggs, yam, garlic, onions, potatoes, pork, fish, poultry, dried beans, jerky, smoked or salted dried fish, and raisins. More recently, its presence was also detected in bottled water [27], food coloring agents and plant food supplements [28]. Therefore, avoiding the consumption of OTA is nearly impossible, having as a consequence the presence of low but detectable levels of OTA in human blood, coffee being one important source. In the 1970s it was reported the first studies that identified the presence of OTA in human blood in the Balkans [29]. In fact, OTA can be found in human body due to its long half-life of 35 days [30]. In the review of Malir et al. (2016) it is compiled published data on OTA in

human blood samples from healthy persons, and concentrations higher than $1.0 \mu g/L$ were observed in several countries.

OTA displays a vast toxicity, including neurotoxic, teratogenic, immunotoxic, carcinogenic, hepatotoxic, embryotoxic and especially nephrotoxic in laboratory and farm animals [19,21]. Toxicokinetics data showed that the susceptibility to cancer depending on the species and sex of the tested animals [19,31]. The available kinetic data of OTA for humans is rather limited than for animals, but gender (male population showed higher values), and particularly season (in summer the highest values were obtained) and geographic location (identify with dietary habits of each particular region) appear to be the most relevant factors that influencing OTA concentrations in plasma [32].

In 1993, the International Agency for Research on Cancer has evaluated OTA as a possible human carcinogen (Group 2B) based on inadequate evidence in humans for the carcinogenicity of OTA [33]. Surprisingly, chronic exposure to low OTA doses could be even more severe effects than acute exposure to a high dose [21,34]. It is generally accepted that the kidney is the classical target organ of OTA [34]. The toxin at nanomolar concentrations in renal cells may be regarded as a modulator of cellular signaling, since it interacts with certain cellular key targets resulting in specific alterations of cell function and/or phenotype and even may stable re-program the cells. At these concentrations OTA can induce apoptosis in culture cells, while necrosis has only a minor role in cellular death [34]. Later, Schwerdt showed that exposure to very low OTA concentrations (0.3–10 nm) for prolonged time (up to 14 days) led not only to cell hypertrophy, but also affect the cellular fate in human proximal tubule cells [35]. Moreover, findings of increased secretion of fibronectin, NF-kB activity and modified collagen pattern, was proposed for OTA induced tubule-interstitial fibrosis which is frequently observed in Balkan endemic nephropathy [35]. Toxicogenomic studies of Balkan nephropathy on the human genes that influence susceptibility to OTA, lead to the hypotheses about the association between Balkan nephropathy to be produced by OTA [36]. Nowadays, it is assumed that OTA is the probable causal agent in the development of Balkan endemic nephropathy, a chronic tubulointerstitial kidney disease that may lead to kidney tumors and irreversible kidney failure [13,37].

An in vitro study conducted in human fetal gastric epithelium cells (GES-1) showed that OTA could induce a G2 arrest, increasing intracellular reactive oxygen species (ROS) production and consequently inducing DNA damage. The oxidative DNA damage elevated the levels of 8-hydroxydeoxyguanosine and DNA double-strand breaks. It was also observed that OTA exposure induced the phosphorylation of the ataxia telangiectasia-mutated (ATM) protein suggesting that OTA-induced oxidative DNA damage triggered the ATM-dependent pathways [38]. Interestingly, among normal human cells, OTA could induce a G1 arrest in human peripheral blood mononuclear cells [39].

At present, although there is insufficient data regarding in utero exposure and the fetotoxicity of OTA, it is known that the level of OTA in human umbilical cord blood is higher than maternal blood at the time of delivery. Considering that human blood samples from healthy individuals could be relatively high as described above, it is recommended that OTA exposure of pregnant women should kept as low as possible in order to reduce the possibility to develop cancer early in life, or even increase susceptibility to chronic disease or cancer later in life [40].

A recent data analysis on OTA suggested a strong evidence of its carcinogenicity in humans [21], probably due to induction of oxidative stress by OTA [13] or through oxidative DNA lesions coupled with direct DNA adducts via quinone formation [17]. This last theory concurs with the Maaroufi and Arlt studies, which detected OTA-DNA adducts in tissues of humans exposed to OTA [37,41]. On the other hand, other studies postulated that OTA induced DNA damage in association with ROS production in several cell types including human kidney (HK-2), proximal tubular cells and colonic adenocarcinoma cell line (CaCo-2), as well as peripheral blood mononuclear cells and hepatoma cell lines (HepG2) [38,39,42–44]. Indeed, it is believed that OTA genotoxicity is through oxidative metabolism, playing and important role in OTA-mediated carcinogenesis [17]. Furthermore, a model for the role of miR-122, the most abundant liver micro-RNA (miRNA), in OTA-induced hepatocyte apoptosis in vivo and in vitro was proposed [45].The expression of miR-122 in HepG2 was upregulated when treated

with OTA. miR-122 is a primary effector of OTA-induced hepatocyte apoptosis. The mechanism across CCNG1/P53 and Bcl-w/caspase-3 pathways was described [45].

The mechanism of toxicity of OTA is complex and new data are necessary to better understand all the events implicated. Nevertheless, taking into account the previous and recently available data, there are voices in the scientific community that propose upgrade its carcinogenicity from Group 2B to at least Group 2A (probably carcinogenic to humans) [20,35,46–48] or even Group 1 (carcinogenic to humans) [21].

2.2. Fungal Contamination and Coffee

Green coffee beans are prone to fungal attack. Among these microorganisms, Aspergillus species such as Aspergillus carbonarius, A. niger, A. ochraceus and A. westerdijkiae are the most widely OTA producers in tropical and semi-tropical coffee plantations [7,49]; meanwhile in temperate regions Penicillium species such as P. verruculosum, P. brevicompactum, P. crustosum, P. olsonii and P. oxalicum are usually found [50,51]. Recently, a strain of Byssochlamys spectabilis isolated from roasted coffee from Mexico markets was able to produce OTA [52]. An EU study confirmed that green coffee is a raw material susceptible to OTA contamination [53]. The mycotoxigenic potencial to contaminate the coffee beans depends on several factors such as species and strain of fungus, water activity, environmental (climatic conditions, storage and transportation) and processing (wet and dry processing) conditions [7,54–56]. In fact, OTA has been reported at different stages of the food chain, indicating that fungus colonization may occur before or after coffee harvest, depending on environmental or storage conditions, particularly in the case that coffee seeds are not dried to a safe moisture percentage or if rehydrated during any of the phases of drying, storage or transportation [55]. Factors such as the adequate cherry maturation using normal water, fertilizers and maintenance, as well as harvest only ripe cherries which were not in contact with soil to prevent microorganism contamination and the appearance of black and sour defective beans, and the reduction of defects at purchasing are important points in green coffee production chain. Finally, the transport should be done to well-dried green coffee (10–12% MC) and storage conditions should assure an atmosphere at 50–70% RH and a temperature below 20 °C to keep OTA levels generally low [55,57]. Therefore, the critical steps for OTA accumulation are considered to be the harvest and the postharvest handling of coffee cherries [7]. Additionally, OTA is stable under acidic conditions, high temperatures and pressure steam sterilization, making difficult the control of its sanitary quality in order to minimize the risk of its intake from coffee [19,58]. Taking into account the risk that OTA presents to human health, the European Union has developed regulations and fixed a maximum level for OTA in the roasted coffee bean and ground roasted coffee of 5 μ g/kg of ochratoxin A [59]. Additionally, based on the toxicological data, the European Safety Authority fixed a tolerable weekly intake (TWI) of 120 ng/kg body weight [60].

3. Can Ochratoxin A Reduce Coffee Consumption?

An EU study reported that there is a number of data sufficiently high to provide sound information on the green and processed coffee contamination by OTA [53]. The percentage of positive samples was 36% and 46% for green coffee beans and roasted coffee, respectively, among EU member states, with average of OTA contamination levels of 3.641 μ g/kg and 1.092 μ g/kg, respectively. Nevertheless, countries like Germany, Greece, Italy, Spain and Netherlands reported positive samples for roasted coffee with at least 5.000 μ g/kg of OTA [53]. The occurrence of OTA in roasted coffee bean has also been reported by other studies in concentrations ranging between 0.09 μ g/kg and 12.1 μ g/kg (Table 2). The highest level of OTA (12.1 μ g/kg) was found in a sample from Germany [61]. Although in countries like Argentina, Brazil, France and Portugal, OTA was also detected in samples exceeding the maximum level established by EU, the number of samples was reduced comparing to sampling [62–66]. Despite of the lack of consensus in literature regarding the percentage of toxin that is transferred into beverage during brewing, it is known that home-processing methods can influence the OTA concentration in coffee brews [67–71]. It was reported that a reduction of approximately 50% of the toxic potential of roast and ground coffee was obtained in expresso coffee, meanwhile in moka and auto-drip brews, a lower efficacy was achieved [69]. Malir noticed that the amount of OTA transfer into beverages was dependent on raw material contamination, time of contact, amount of ground coffee and water used as well as temperature of water and pH of the beverages. In the study conducted by this research group, values ranged from 22.3% to 66.1% of OTA transfer into coffee were found [67].

Country	n	% Positive	Average (µg/kg)	Range of OTA (µg/kg)	Reference
Argentina	24	54	1.00	0.11-5.78	[64]
Brazil	34	68	0.9	0.3–6.5	[65]
Brazil	16	31		0.09–9	[62]
Canada	71	59	0.6	0.1–2.3	[72]
Chile	24	100	0.47	tr-0.84	[3]
France	30	100		tr-11.9	[66]
Germany	490	44	1.48	<0.15-12.1	[61]
Hungary	38	58	0.5	0.17-0.91	[73]
Italy	30	27	0.27	tr-<5.0	[74]
Portugal	11	27	1.66	0.71-10.31	[63]
Spain	72	49	1.05	1.21-4.21	[75]
UK	20	85	0.6	0.2–2.1	[76]

Table 2. Occurrence of OTA in roasted and/or ground roasted coffee by country.

tr, traces (values of ochratoxin A between LOD and LOQ).

The consumption of this pleasant functional food due its taste and aroma could be taken with caution due to the presence of OTA, especially to high consumers where an intake of more than four coffees by day is a possible scenario. Moreover, if coffee beverage is prepared with approximately 7 g of ground coffee for 100 mL of water, the intake of a high consumer weighing 60 kg could corresponds to half of tolerate daily intake of ochratoxin A set up by JECFA, which made OTA intake via coffee beverage not negligible [66]. Nowadays, it is estimated that coffee accounts for approximately 12% of the total intake of OTA, being considered the third source of OTA exposure [3,77]. In a comparison study to assess human exposure to the OTA and 2'R-ochratoxin A (2'R-OTA, previously named 14R-Ochratoxin A), a product from thermal degradation, between coffee and noncoffee drinkers showed that no correlation exist between the amounts of coffee consumption and OTA or 2'R-OTA concentrations. However, 2'R-OTA was only detected in blood of coffee drinkers [78].

4. Strategies to Reduce Ochratoxin A

4.1. Coffee Roasted

Coffee roasting, one of the most important stages to obtain the brew, where non-enzymatic browning occurs depending on time and temperature that can be applied to green coffee samples, is responsible for the characteristics of the final product like the pleasant flavor as a consequence of chemical reactions. Besides, this interesting effect that make coffee an unique beverage, the roasting process is described as a step that decreases OTA levels [7,49,64,79,80]. Indeed, until 1988, it was believed that the roasting process completely degraded this mycotoxin. Tsubouchi et al. reported for the first time the occurrence of ochratoxin A in coffee brews [81]. Several studies have been published from then till now, particularly, using coffee "naturally contaminated" or "artificially contaminated" in order to demonstrate the effect of roasting on the OTA concentration (Table 3).

Method	% Reduction	Remarks	Reference
	67	Temp = $470 \degree$ C; Time = 2.5 min; batch size = 5 kg	
	63	Temp = 490 °C; Time = 2.5 min; batch size = 5 kg	
Deacting	74	Temp = 490 °C; Time = 4 min; batch size = 10 kg	[00]
Roasting	53	Temp = 400 °C; Time = 10 min; batch size = 15 kg	[82]
	84	Temp = 425 °C; Time = 10 min; batch size = 15 kg	
		$[OTA]_{initial} = 4.9 \ \mu g/kg;$ (Gothot RN 100 pilot size roaster)	
	77.9 ^a ; 64.1 ^b	Temp = 450 °C ; Time = 6 min; batch size = 3 kg	
Desetions	90.5 ^a ; 88.7 ^b	Temp = 450 °C ; Time = 7 min; batch size = 3 kg	[00]
Roasting	96.8 ^a ; 95.1 ^b	Temp = $450 \degree$ C; Time = 9 min; batch size = 3 kg	[80]
	,	^a [OTA] _{initial} = 29.30 µg/kg; ^b [OTA] _{initial} = 9.64 µg/kg; (Laboratory roaster,	
		model 500, STA)	
Deasting	21.1	Temp = 180 °C; Time = 10 min; [OTA] _{initial} = 29.36 μg/kg	[02]
Koasting	31.1	(Normal coffee roaster)	[83]
Deasting	70.1	Temp = 260 °C; Time = 5 min; batch size = 0.5 kg ; [OTA] _{initial} = $2.54 \mu \text{g/kg}$;	[60]
Roasting	79.1	(Precision Coffee Roaster model, Hearthware Home Products)	[09]
	25.1 ^a ; 15.1 ^b	Temp = 230 °C; Time = 6 min; batch size = 0.2 kg	
	59.5 ^a ; 67.9 ^b	Temp = $230 \degree$ C; Time = $9 \min$; batch size = 0.2 kg	
	87.0 ^a ; 89.6 ^b	Temp = 230 °C; Time = 12 min; batch size = 0.2 kg	
	97.2 ^a ; 95.1 ^b	Temp = 230 °C; Time = 15 min; batch size = 0.2 kg	
Roasting		(Rotating cylinder- Probat-Werne type RE1 pilot roaster)	[49]
nousing	7.4 ^a ; 9.3 ^b	Temp = 230 °C; Time = 0.9 min; batch size = 0.2 kg	[17]
	36.0 ^a ; 30.9 ^b	Temp = 230 °C; Time = 1.7 min; batch size = 0.2 kg	
	72.4 ^a ; 64.3 ^b	Temp = 230 °C; Time = 2.6 min; batch size = 0.2 kg	
	77.6 ^a ; 73.4 ^b	Temp = 230 °C; Time = 3.5 min; batch size = 0.2 kg	
		(Fluidized bed - Neuhaus Neotec pilot roaster)	
		$a[OTA]_{initial} = 5.3 \ \mu g/kg; b[OTA]_{initial} = 57.2 \ \mu g/kg$	
	5	$a_w = 0.59$; Dried green coffee beans;	
	9	$a_w = 0.65$; Incubated at RH 76 for 72 h;	
Gamma	20	$a_w = 0.71$; Incubated at RH 99 for 72 h	[84]
radiation	90	$a_w = 0.82$; Steamed for 1 h	[04]
	≈100	$a_w = 0.93$; Soaked for 20 h	
		[OTA] _{initial} = 50 ppb; radiation = 10 kGy (Gamma Chamber 5000)	
	0	P = 50 MPa; Time storage = 10 days; [OTA] _{initial} = 4.4 µg/kg	
	54.6	P = 200 MPa; Time storage = 10 days; [OTA] _{initial} = 3.9 µg/kg	
	93.2	P = 400 MPa; Time storage = 10 days; [OTA] _{initial} = 4.4 µg/kg	
High- pressure	96.2	P = 600 MPa; Time storage = 10 days; [OTA] _{initial} = 4.5 µg/kg	
	14.6	P = 50 MPa; Time storage = 20 days; [OTA] _{initial} = 4.4 µg/kg	
	43.6	$P = 200 \text{ MPa}$; Time storage = 20 days; $[OTA]_{initial} = 3.9 \mu g/kg$	
	93.2	$P = 400 \text{ MPa}$; Time storage = 20 days; $[OTA]_{initial} = 4.4 \mu g/kg$	
	96.0	P = 600 MPa; Time storage = 20 days; [OTA] _{initial} = 4.5 µg/kg	
	19.1	P = 50 MPa; Time storage = 30 days; [OTA] _{initial} = 4.4 µg/kg	[60]
	45.7	$P = 200$ MPa; Time storage = 30 days; $[OTA]_{initial} = 3.9 \mu g/kg$	
	91.9	P = 400 MPa; Time storage = 30 days; [OTA] _{initial} = 4.4 µg/kg	
	95.5	P = 600 MPa; Time storage = 30 days; [O1A] _{initial} = 4.5 µg/kg	
	26.9	P = 50 MPa; Time storage = 50 days; [OTA] _{initial} = 4.4 µg/kg	
	36.3	P = 200 MPa; Time storage = 50 days; [OTA] _{initial} = 3.9 µg/kg	
	90.8	r = 400 MPa; Time storage = 50 days; [OTA] _{initial} = 4.4 µg/kg	
	74.4	I = 000 WIFa; Time storage = 50 days; [OTAJ _{initial} = 4.5 µg/kg (BaoTou KoFa High Processive Technology Ca.)	
		(Da0100 Kera right ressure rechnology C0.)	

Table 3. Physical approaches for detoxification of OTA.

One of these interesting studies investigated the effect of roasting conditions on the reduction of OTA during roasting [82]. Roasting conditions varied from clearly roasting in commercial scale equipment to clearly outside the usually roasting practice, and no statistically significant differences between the color and the levels of the applied roasting were observed. In this study a mean value for OTA reduction of 68% on mycotoxin content was achieved. Nevertheless, the highest OTA reduction (84%) was found for the combination of long plus dark roasting, indicating that statistical significance could be obtained in darker colors and longer times of treatment [82]. The influence of roasting process on the OTA levels in coffee beans was also reported by Romani and collaborators. The reduction of OTA

content was a function of the severity of the thermal treatment and seems to be generally related to the initial OTA concentration (high percentages of OTA reduction in the highest contaminated samples) [80]. The effect of two different roasting techniques, rotating cylinder and fluidized bed, on OTA levels of green coffee beans (Coffea arabica) artificially contaminated with Aspergillus westerdijkiae have been reported [49]. A reduction on OTA levels above 96% and 75% was achieved under rotating cylinder and fluidized bed at the same roasting degree, respectively [49]. In a study conducted by Drunday and Pacin, the average of OTA levels in green coffee was almost six times greater than roasted ground coffee samples from markets located in Buenos Aires Province [64]. On contrary, Tsubouchi et al. described roasting at 200 °C during 10 and 20 min as an ineffective treatment in the reduction of OTA levels of contaminated coffee beans, since only a decrease of 0–12% was observed [71]. The discrepancy between these findings is difficult to explain but the OTA reduction during roasting may be due to: Epimerization at the C3 position [70,83], thermal degradation with possible involvement of moisture and a physical removal of OTA with chaff [79,83]. In a model system simulating degradation reactions of coffee roasting two products were identified, 14-(R)-ochratoxin A the major compound formed and 14-decarboxy-ochratoxin A at traces levels [24]. Detoxification of OTA by the formation of ochratoxin A saccharide esters and ochratoxin α amide during the coffee roasting was proposed [22,23]. When OTA is submitted to temperatures of 200 °C and above, a partial decomposition of the mycotoxin was observed and the ochratoxin α amide was formed as the thermal degradation product by a cleavage between the carbon atom C-14 and the nitrogen atom [22]; while in the case of ochratoxin A saccharide esters, a ester linkage between the carboxyl group of OTA and the primary hydroxyl group of the carbohydrates of the matrix was demonstrated [23]. Furthermore, ochratoxin α amide can be formed by photolytic degradation during the exposure of OTA to blue and white light, suggesting light treatment as a possible treatment to destroy OTA [22].

4.2. Coffee Composition: Caffeine

Coffee contains caffeine, 1,3,7-trimethylxanthine, a psychoactive molecule and one of the most important active compounds with antioxidant properties, being the major water-soluble component of coffee. Caffeine is described as a compound able to reduce type 2 diabetes mellitus disease and incidence of cancer, as well as protect against Parkinson's disease and prevent bronchopulmonary dysplasia in premature neonates [86–89]. Meanwhile, caffeine concentration varies substantially, depending on coffee plant species, method of bean roasting, as well as drinks preparation [2]. Several studies have suggested that caffeine content could be a strategy to prevent OTA in green coffee bean, based on the fact that its presence reduces OTA production [90–92]. Buchanan et al. (1982) reported the effect of caffeine on growth and OTA production by Aspergillus ochraceus NRRL 3174 in YES medium. Caffeine was less inhibitory to growth than OTA synthesis. A dose related decrease in OTA accumulation was observed. A concentration of 3 mg/mL of caffeine inhibited OTA almost completely (98%). Despite different OTA inhibition percentages, a similar trend in AMY medium supplemented with 4 mg/mL of caffeine after 14 days of incubation was observed for mycotoxigenic strains, namely A. flavus PC65, A. versicolor XVII/19, Penicillium urticase NRRL 989, P. roqueforti M251 and P. expansum NRRL 973 [91]. In contrast, Aspergillus ochraceus strains S-235-100 isolated from green coffee beans produced higher amounts of OTA when grown on YES medium supplemented with 0.5% and 1.0% of caffeine, reaching a maximum after 15 to 20 days of incubation [92]. However, in the same study, it was investigated in solid medium the effect of caffeine on OTA production by 15 strains of A. ochraceus, three strains of A. elegans and one of A. sclerotiorum, and concluded that caffeine, generally, had little or no effect on OTA production by the majority of A. ochraceus strains, but did inhibit toxin production by A. elegans and A. sclerotiorum strains. While, Nehad et al. (2005) reported that caffeine at concentrations of 1.0% and 2.0% completely prevented OTA production by A. ochraceus in YES medium [83]. Recently, in a study conducted by Akbar et al. (2016) demonstrated that OTA production by strains from Aspergillus section *Circumdati* and *Aspergillus* section *Nigri* groups isolated from coffee had differential tolerance to concentrations of caffeine. If low concentrations, less than 0.5%, of caffeine were inhibitory to OTA

production by some strains, namely *A. carbonarius, A. westerdijkiae* and *A. niger*, at least 1.5% of caffeine was needed for *A. steynii* inhibition; while 4% of caffeine was required for *A. ochraceus*. However, OTA production was stimulated with 3% of caffeine [90]. In summary, it appears that caffeine has different effects on OTA production depending on its concentration but, particularly, of mycotoxigenic species that have the ability to contaminate coffee beans.

4.3. Physical Methods

Several approaches have been developed to detoxify OTA, including the use of organic and inorganic adsorbents, irradiation, high-pressure, ultrasound, ozone, and biodegradation and/or biological adsorption. Factors such as nutritional quality, organoleptic properties, safety, availability, and cost-effectiveness are determinant in the implementation of these technologies. However, it is out of the scope of the present review to describe the different methods to detoxify OTA in general foods since it is focused in the coffee matrix, but for those readers who are interested in this aspect we recommend other reviews published elsewhere [93–95]. In fact, in coffee, the number of reports on OTA detoxification is very scarce (Figure 2).



Figure 2. Ochratoxin A in coffee, including its natural sources derived from fungal contamination of beans, their possible toxic effects in humans (black lines) and the potential treatments for minimizing its effects by reducing the fungal contamination of coffee beans (red lines).

One of these examples investigated ozone and ultrasound as alternative processes in the treatment of fermented coffee [96]. The results showed that ozone and ultrasound exhibits a potential in reducing OTA in coffee, due to filamentous fungi inactivation. Besides the biological effect, both treatments reduced the total soluble solids and increased the phenolic compounds content, increasing fruit astringency which concentration is inversely related with brew quality. Meanwhile, this was not observed in sensorial analysis since no perceptible changes in the quality of the beverage were detected. According to these researchers, ozone and ultrasounds could be applied as pre-treatments before coffee fermentation, as they minimally affect the quality of coffee naturally contaminated with OTA. Nutrient losses or changes are negligible compared to the benefits, improving food safety of the fermented coffee [96].

Recently, the detoxification of OTA using gamma radiation and high-pressure processing (HPP) were proposed, taking into account not only the feasibility and efficacy of the method but also because it overall not promote coffee organoleptic changes [84,93]. In fact, both approaches are known to keep foods' original freshness, but efficacy depends on food composition, particularly water activity. In the case of gamma radiation processing, the study performed by Kumar et al. (2012) showed that green coffee beans having aw of 0.93 when submitted to a dose of 10 kGy OTA was almost completely degraded; while aw of 0.82, 0.65 and 0.59 resulted in approximately 90%, 9% and 5% of OTA reduction, respectively, for an initial concentration of 50 ppb [84]. The influence of HPP in the inhibition of *Aspergillus fresenii* growth was study by Chen et al. (2018), indicating that HPP at pressures of 400 MPa or higher inhibited fungal growth on coffee beans; although no effect was observed on OTA its accumulation was prevented. Additionally, it was observed that 400 MPa not only maintained OTA levels, but also increase γ -aminobutyric acid content after 50 days of storage improving the nutritional value of coffee [85]. Therefore, HPP seems to be a promising technology not to destroy OTA but indirectly avoiding its increment in coffee beans.

Despite the consensus regarding to flexibility and promising results of these new technologies in the treatment of coffee, there is a great concern on definition of optimum treatment conditions (well-directed standard operating procedures in approved laboratories) and toxicity of degradation products formed, and consequent implications on human and animal health.

5. Conclusions

Coffee that is more than a mere beverage, due to its functional and sensorial properties, it could represent a threat, especially to the higher coffee consumer, attending to OTA levels. Good manufacturing practices and hygiene standards during coffee production is strongly recommended to reduce the risk of contamination processed coffee. Additionally, it is necessary to invest in knowledge in order to develop and/or improve effective and friendly technologies to detoxify coffee beans. The inclusion of gamma radiation in the postharvest processing chain of green coffee beans could be a promising approach in order to not only eliminate OTA producer fungi but also the toxicity of performed OTA. Ozone and ultrasound could be applied as pre-treatments before coffee fermentation, while HPP could be introduced in steps presenting high risk of microbiological hazards in the coffee industry without altering the manufacturing process. These promising methods could be complementarily applied toward reducing OTA contamination in the coffee without affecting the sensorial attributes that makes coffee one of the most pleasant products, promoting food safety and quality. However, further studies on emerging new technologies are needed not only in what concerns to improvement and control but also in their application in the coffee treatment, ensuring their effectiveness and safety use in the coffee processing.

Funding: This research received no external funding.

Acknowledgments: Author thanks to MEtRICs, ref. UID/EMS/04077/2019.

Conflicts of Interest: The author declares no conflicts of interest.

References

- 1. Organization, I.C. Annual Review 2016/2017; International Coffee Organization: London, UK, 2018; pp. 1–50.
- 2. Dorea, J.G.; da Costa, T.H. Is coffee a functional food? Br. J. Nutr. 2005, 93, 773–782. [CrossRef] [PubMed]
- 3. Galarce-Bustos, O.; Alvarado, M.; Vega, M.; Aranda, M. Occurence of ochratoxin A in roasted and instant coffees in Chilean. *Food Control* 2014, *46*, 102–107. [CrossRef]
- Nishitsuji, K.; Watanabe, S.; Xiao, J.; Nagatomo, R.; Ogawa, H.; Tsunematsu, T.; Umemoto, H.; Morimoto, Y.; Akatsu, H.; Inoue, K.; et al. Effect of coffee or coffee components on gut microbiome and short-chain fatty acids in a mouse model of metabolic syndrome. *Sci. Rep.* 2018, *8*, 16173. [CrossRef]

- 5. Samoggia, A.; Riedel, B. Coffee consumption and purchasing behavior review: Insights for further research. *Appetite* **2018**, *129*, 70–81. [CrossRef] [PubMed]
- Jeszka-Skowron, M.; Sentkowska, A.; Pryrzynska, K.; Peña, M.P. Chlorogenic acids, caffeine content and antioxidant properties of green coffee extracts: Influence of green coffee bean preparation. *Eur. Food Res. Technol.* 2016, 242, 1403–1409. [CrossRef]
- Napolitano, A.; Fogliano, V.; Tafuri, A.; Ritieni, A. Natural occurrence of ochratoxin A and antioxidant activities of green and roasted coffees and corresponding byproducts. J. Agric. Food Chem. 2007, 55, 10499–10504. [CrossRef]
- 8. Alves, R.C.; Casal, S.; Alves, M.R.; Oliveira, M.B. Discrimination between arabica and robusta coffee species on the basis of their tocopherol profiles. *Food Chem.* **2009**, *114*, 295–299. [CrossRef]
- 9. Priftis, A.; Mitsiou, D.; Halabalaki, M.; Ntasi, G.; Stagos, D.; Skaltsounis, L.A.; Kouretas, D. Roasting has a distinct effect on the antimutagenic activity of coffee varieties. *Mutat. Res.* **2018**, *829–830*, 33–42. [CrossRef]
- Butt, M.S.; Sultan, M.T. Coffee and its consumption: Benefits and risks. *Crit. Rev. Food Sci. Nutr.* 2011, 51, 363–373. [CrossRef]
- 11. Sharif, K.; Watad, A.; Bragazzi, N.L.; Adawi, M.; Amital, H.; Shoenfeld, Y. Coffee and autoimmunity: More than a mere hot beverage! *Autoimmun. Rev.* **2017**, *16*, 712–721. [CrossRef]
- 12. Machado-Fragua, M.D.; Struijk, E.A.; Graciani, A.; Guallar-Castillon, P.; Rodriguez-Artalejo, F.; Lopez-Garcia, E. Coffee consumption and risk of physical function impairment, frailty and disability in older adults. *Eur. J. Nutr.* **2018**. [CrossRef]
- 13. Tao, Y.; Xie, S.; Xu, F.; Liu, A.; Wang, Y.; Chen, D.; Pan, Y.; Huang, L.; Peng, D.; Wang, X.; et al. Ochratoxin A: Toxicity, oxidative stress and metabolism. *Food Chem. Toxicol.* **2018**, *112*, 320–331. [CrossRef]
- 14. Pohland, A.E.; Schuller, P.L.; Steyn, P.S.; Van Egmond, H.P. Physico-chemical data for selected mycotoxins. *Pure Appl. Chem.* **1982**, *54*, 2219–2284. [CrossRef]
- 15. Natori, S.; Sakaki, S.; Kurata, H.; Udagawa, S.; Ichinoe, M. Chemical and cytotoxicity survey on the production of ochratoxins and penicillic acid by *Aspergillus ochraceus* Wilhelm. *Chem. Pharm Bull.* (*Tokyo*) **1970**, *18*, 2259–2268. [CrossRef]
- 16. van der Merwe, K.J.; Steyn, P.S.; Fourie, L.; Scott, D.B.; Theron, J.J. Ochratoxin A, a toxic metabolite produced by *Aspergillus ochraceus* Wilh. *Nature* **1965**, *205*, 1112–1113. [CrossRef]
- 17. Pfohl-Leszkowicz, A.; Manderville, R.A. Ochratoxin A: An overview on toxicity and carcinogenicity in animals and humans. *Mol. Nutr. Food Res.* **2007**, *51*, 61–99. [CrossRef]
- 18. Bayman, P.; Baker, J.L. Ochratoxins: A global perspective. Mycopathologia 2006, 162, 215–223. [CrossRef]
- 19. el Khoury, A.; Atoui, A. Ochratoxin a: General overview and actual molecular status. *Toxins* (*Basel*) **2010**, 2, 461–493. [CrossRef]
- Heussner, A.H.; Bingle, L.E. Comparative Ochratoxin Toxicity: A Review of the Available Data. *Toxins (Basel)* 2015, 7, 4253–4282. [CrossRef]
- Malir, F.; Ostry, V.; Pfohl-Leszkowicz, A.; Malir, J.; Toman, J. Ochratoxin A: 50 Years of Research. *Toxins (Basel)* 2016, 8, 191. [CrossRef]
- 22. Bittner, A.; Cramer, B.; Harrer, H.; Humpf, H.-U. Structure elucidation and in vitro cytotoxicity of ochratoxin α amide, a new degradation product of ochratoxin A. *Mycotoxin Res.* **2015**, *31*, 83–90. [CrossRef]
- 23. Bittner, A.; Cramer, B.; Humpf, H.U. Matrix binding of ochratoxin A during roasting. *J. Agric. Food Chem.* **2013**, *61*, 12737–12743. [CrossRef]
- 24. Cramer, B.; Konigs, M.; Humpf, H.U. Identification and in vitro cytotoxicity of ochratoxin A degradation products formed during coffee roasting. *J. Agric. Food Chem.* **2008**, *56*, 5673–5681. [CrossRef]
- 25. Koszegi, T.; Poor, M. Ochratoxin A: Molecular Interactions, Mechanisms of Toxicity and Prevention at the Molecular Level. *Toxins (Basel)* **2016**, *8*, 111. [CrossRef]
- 26. Levi, C.P.; Trenk, H.L.; Mohr, H.K. Study of the occurrence of ochratoxin A in green coffee beans. J. Assoc. Off. Anal. Chem. 1974, 57, 866–870.
- 27. Mata, A.T.; Ferreira, J.P.; Oliveira, B.R.; Batoreu, M.C.; Barreto Crespo, M.T.; Pereira, V.J.; Bronze, M.R. Bottled water: Analysis of mycotoxins by LC-MS/MS. *Food Chem.* **2015**, *176*, 455–464. [CrossRef]
- 28. Solfrizzo, M.; Piemontese, L.; Gambacorta, L.; Zivoli, R.; Longobardi, F. Food coloring agents and plant food supplements derived from *Vitis vinifera*: A new source of human exposure to ochratoxin A. *J. Agric. Food Chem.* **2015**, *63*, 3609–3614. [CrossRef]

- 29. Hult, K.; Plestina, R.; Habazin-Novak, V.; Radic, B.; Ceovic, S. Ochratoxin A in Human Blood and Balkan Endemic Nephropathy. *Arch. Toxicol* **1982**, *51*, 313–321. [CrossRef]
- 30. WHO. Ochratoxin A.; WHO: Geneva, Switzerland, 2008; pp. 356-429.
- 31. Malir, F.; Ostry, V.; Pfohl-Leszkowicz, A.; Novotna, E. Ochratoxin A: Developmental and reproductive toxicity-an overview. *Birth Defects Res. B Dev. Reprod. Toxicol.* **2013**, *98*, 493–502. [CrossRef]
- 32. Coronel, M.B.; Sanchis, V.; Ramos, A.J.; Marin, S. Review. Ochratoxin A: Presence in human plasma and intake estimation. *Food Sci. Technol. Int.* **2010**, *16*, 5–18. [CrossRef]
- 33. IARC. Some Naturally Occuring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins; WHO/IARC, Ed.; IARC Press: Lyon, France, 1993; Volume 56, pp. 489–521.
- 34. Gekle, M.; Sauvant, C.; Schwerdt, G. Ochratoxin A at nanomolar concentrations: A signal modulator in renal cells. *Mol. Nutr. Food Res.* **2005**, *49*, 118–130. [CrossRef]
- 35. Schwerdt, G.; Holzinger, H.; Sauvant, C.; Konigs, M.; Humpf, H.U.; Gekle, M. Long-term effects of ochratoxin A on fibrosis and cell death in human proximal tubule or fibroblast cells in primary culture. *Toxicology* **2007**, 232, 57–67. [CrossRef]
- 36. Stefanovic, V.; Toncheva, D.; Polenakovic, M. Balkan nephropathy. Clin. Nephrol. 2015, 83, 64-69. [CrossRef]
- Maaroufi, K.; Pfohl-Leszkowicz, A.; Achour, A.; el May, M.; Grosse, Y.; Hammami, M.; Ellouz, F.; Creppy, E.E.; Bacha, H. Ochratoxin A genotoxicity, relation to renal tumors. *Arch. Inst. Pasteur Tunis* 1994, 71, 21–31.
- Cui, J.; Liu, J.; Wu, S.; Wang, Y.; Shen, H.; Xing, L.; Wang, J.; Yan, X.; Zhang, X. Oxidative DNA damage is involved in ochratoxin A-induced G2 arrest through ataxia telangiectasia-mutated (ATM) pathways in human gastric epithelium GES-1 cells in vitro. *Arch. Toxicol.* 2013, *87*, 1829–1840. [CrossRef]
- Liu, J.; Wang, Y.; Cui, J.; Xing, L.; Shen, H.; Wu, S.; Lian, H.; Wang, J.; Yan, X.; Zhang, X. Ochratoxin A induces oxidative DNA damage and G1 phase arrest in human peripheral blood mononuclear cells in vitro. *Toxicol. Lett.* 2012, 211, 164–171. [CrossRef]
- 40. Woo, C.S.; El-Nezami, H. Maternal-Fetal Cancer Risk Assessment of Ochratoxin A during Pregnancy. *Toxins (Basel)* **2016**, *8*, 87. [CrossRef]
- 41. Arlt, V.M.; Pfohl-Leszkowicz, A.; Cosyns, J.; Schmeiser, H.H. Analyses of DNA adducts formed by ochratoxin A and aristolochic acid in patients with Chinese herbs nephropathy. *Mutat. Res.* **2001**, *494*, 143–150. [CrossRef]
- 42. Arbillaga, L.; Azqueta, A.; Ezpeleta, O.; Lopez de Cerain, A. Oxidative DNA damage induced by Ochratoxin A in the HK-2 human kidney cell line: Evidence of the relationship with cytotoxicity. *Mutagenesis* **2007**, *22*, 35–42. [CrossRef]
- Guerra, M.C.; Galvano, F.; Bonsi, L.; Speroni, E.; Costa, S.; Renzulli, C.; Cervellati, R. Cyanidin-3-O-betaglucopyranoside, a natural free-radical scavenger against aflatoxin B1- and ochratoxin A-induced cell damage in a human hepatoma cell line (Hep G2) and a human colonic adenocarcinoma cell line (CaCo-2). *Br. J. Nutr.* 2005, 94, 211–220. [CrossRef]
- Schaaf, G.J.; Nijmeijer, S.M.; Maas, R.F.; Roestenberg, P.; de Groene, E.M.; Fink-Gremmels, J. The role of oxidative stress in the ochratoxin A-mediated toxicity in proximal tubular cells. *Biochim. Biophys. Acta* 2002, 1588, 149–158. [CrossRef]
- 45. Zhu, L.; Yu, T.; Qi, X.; Yang, B.; Shi, L.; Luo, H.; He, X.; Huang, K.; Xu, W. miR-122 plays an important role in ochratoxin A-induced hepatocyte apoptosis in vitro and in vivo. *Toxicol. Res.* **2016**, *5*, 160–167. [CrossRef]
- 46. Hennemeier, I.; Humpf, H.U.; Gekle, M.; Schwerdt, G. Role of microRNA-29b in the ochratoxin A-induced enhanced collagen formation in human kidney cells. *Toxicology* **2014**, 324, 116–122. [CrossRef]
- 47. Mally, A. Ochratoxin a and mitotic disruption: Mode of action analysis of renal tumor formation by ochratoxin A. *Toxicol. Sci.* **2012**, *127*, 315–330. [CrossRef]
- 48. Pfohl-Leszkowicz, A.; Manderville, R.A. An update on direct genotoxicity as a molecular mechanism of ochratoxin a carcinogenicity. *Chem. Res. Toxicol.* **2012**, *25*, 252–262. [CrossRef]
- 49. Castellanos-Onorio, O.; Gonzalez-Rios, O.; Guyot, B.; Fontana, T.A.; Guiraud, J.P.; Schorr-Galindo, S.; Durand, N.; Suárez-Quiroz, M. Effect of two different roasting techniques on the Ochratoxin A (OTA) reduction in coffee beans (*Coffea arabica*). *Food Control* **2011**, *22*, 1184–1188. [CrossRef]
- 50. Alvindia, D.G.; de Guzman, M.F. Survey of Philippine coffee beans for the presence of ochratoxigenic fungi. *Mycotoxin Res.* **2016**, 32, 61–67. [CrossRef]
- 51. Vega, F.E.; Posada, F.; Peterson, S.W.; Gianfagna, T.J.; Chaves, F. *Penicillium* species endophytic in coffee plants and ochratoxin A production. *Mycologia* **2006**, *98*, 31–42. [CrossRef]

- Casas-Junco, P.P.; Ragazzo-Sanchez, J.A.; Ascencio-Valle, F.J.; Calderon-Santoyo, M. Determination of potentially mycotoxigenic fungi in coffee (*Coffea arabica* L.) from Nayarit. *Food Sci. Biotechnol.* 2018, 27, 891–898. [CrossRef]
- EU. Assessment of Dietary Intake of Ochratoxin A by the Population of EU Member States; Reports on Tasks for Scientific Cooperation; Directorate-General Health and Consumer, Ed.; Scientific Cooperation: Brussels, Belgium, 2002; p. 153.
- 54. Batista, L.R.; Chalfoun, S.M.; Prado, G.; Schwan, R.F.; Wheals, A.E. Toxigenic fungi associated with processed (green) coffee beans (*Coffea arabica* L.). *Int. J. Food Microbiol.* **2003**, *85*, 293–300. [CrossRef]
- 55. Bucheli, P.; Taniwaki, M.H. Research on the origin, and on the impact of post-harvest handling and manufacturing on the presence of ochratoxin A in coffee. *Food Addit. Contam.* **2002**, *19*, 655–665. [CrossRef]
- 56. Paterson, R.R.M.; Lima, N. How will climate change affect mycotoxins in food? *Food Res. Int.* **2010**, 43, 1902–1914. [CrossRef]
- 57. Taniwaki, M.H.; Teixeira, A.A.; Teixeira, A.R.R.; Copetti, M.V.; Iamanaka, B.T. Ochratoxigenic fungi and ochratoxin A in defective coffee beans. *Food Res. Int.* **2014**, *61*, 161–166. [CrossRef]
- 58. Trivedi, A.B.; Doi, E.; Kitabatake, N. Detoxification of Ochratoxin A on Heating under Acidic and Alkaline Conditions. *Biosci. Biotechnol. Biochem.* **1992**, *56*, 741–745. [CrossRef]
- Commission, E. Commission regulation (EC) N° 1881/2006 of 19 December 2006, setting maximum levels for certain contaminants in foodstuffs. Off. J. Eur. Union 2006, L 364, 5–24.
- 60. EFSA. Opinion of the scientific panel on contaminants in the food chain on a request from the commission related to ochratoxin A in food. *EFSA J.* **2006**, *365*, 1–56.
- 61. Otteneder, H.; Majerus, P. Ochratoxin A (OTA) in coffee: Nation-wide evaluation of data collected by German Food Control 1995-1999. *Food Addit. Contam.* **2001**, *18*, 431–435. [CrossRef]
- Bandeira, R.D.C.C.; Uekane, T.M.; Cunha, C.P.; Geaquinto, L.R.O.; Cunha, V.S.; Caixeiro, J.M.R.; Godoy, R.L.O.; Cruz, M.H.C. Development and validation of a method for the analysis of ochratoxin A in roasted coffee by liquid chromatography/electrospray-mass spectrometry in tandem (LC/ESI-MS/MS). *Quim. Nova* 2012, 35, 66–71. [CrossRef]
- 63. Benites, A.J.; Fernandes, M.; Boleto, A.R.; Azevedo, S.; Silva, S.; Leitão, A.L. Occurrence of ochratoxin A in roasted coffee samples commercialized in Portugal. *Food Control* **2017**, *73*, 1223–1228. [CrossRef]
- 64. Drunday, V.; Pacin, A. Occurrence of Ochratoxin A in coffee beans, ground roasted coffee and soluble coffee and method validation. *Food Control* **2013**, *30*, 675–678.
- 65. Leoni, L.A.; Soares, L.M.; Oliveira, P.L. Ochratoxin A in Brazilian roasted and instant coffees. *Food Addit. Contam.* **2000**, *17*, 867–870. [CrossRef] [PubMed]
- 66. Tozlovanu, M.; Pfohl-Leszkowicz, A. Ochratoxin A in roasted coffee from French supermarkets and transfer in coffee beverages: Comparison of analysis methods. *Toxins* (*Basel*) **2010**, *2*, 1928–1942. [CrossRef]
- 67. Malir, F.; Ostry, V.; Pfohl-Leszkowicz, A.; Toman, J.; Bazin, I.; Roubal, T. Transfer of ochratoxin A into tea and coffee beverages. *Toxins (Basel)* **2014**, *6*, 3438–3453. [CrossRef]
- 68. Micco, C.; Grossi, M.; Miraglia, M.; Brera, C. A study of the contamination by ochratoxin A of green and roasted coffee beans. *Food Addit. Contam.* **1989**, *6*, 333–339. [CrossRef] [PubMed]
- 69. Perez De Obanos, A.; Gonzalez-Penas, E.; Lopez De Cerain, A. Influence of roasting and brew preparation on the ochratoxin A content in coffee infusion. *Food Addit. Contam.* **2005**, *22*, 463–471. [CrossRef]
- 70. Studer-Rohr, I.; Dietrich, D.R.; Schlatter, J.; Schlatter, C. The occurrence of ochratoxin A in coffee. *Food Chem. Toxicol.* **1995**, *33*, 341–355. [CrossRef]
- 71. Tsubouchi, H.; Yamamoto, K.; Hisada, K.; Sakabe, Y.; Udagawa, S. Effect of roasting on ochratoxin A level in green coffee beans inoculated with *Aspergillus ochraceus*. *Mycopathologia* **1987**, *97*, 111–115. [CrossRef]
- 72. Lombaert, G.A.; Pellaers, P.; Chettiar, M.; Lavalee, D.; Scott, P.M.; Lau, B.P. Survey of Canadian retail coffees for ochratoxin A. *Food Addit. Contam.* **2002**, *19*, 869–877. [CrossRef]
- 73. Fazekas, B.; Tar, A.K.; Zomborszky-Kovacs, M. Ochratoxin a contamination of cereal grains and coffee in Hungary in the year 2001. *Acta Vet. Hung.* **2002**, *50*, 177–188. [CrossRef]
- 74. Prelle, A.; Spadaro, D.; Denca, A.; Garibaldi, A.; Gullino, M.L. Comparison of clean-up methods for ochratoxin A on wine, beer, roasted coffee and chili commercialized in Italy. *Toxins (Basel)* **2013**, *5*, 1827–1844. [CrossRef]
- 75. Coronel, M.B.; Marin, S.; Cano, G.; Ramos, A.J.; Sanchis, V. Ochratoxin A in Spanish retail ground-roasted coffee: Occurence and assessment of the exposure in Catalonia. *Food Control* **2011**, *23*, 414–419. [CrossRef]

- 76. Patel, S.; Hazel, C.M.; Winterton, A.G.; Gleadle, A.E. Survey of ochratoxin A in UK retail coffees. *Food Addit. Contam.* **1997**, *14*, 217–222. [CrossRef] [PubMed]
- 77. Uekane, T.M.; Bandeira, R.D.C.C.; Silva, M.C. Comparação de métodos de análise para ocratoxina A no café: Uma revisão. *Prespectivas da Ciência e Tecnologia* **2010**, *2*, 44–54.
- Cramer, B.; Osteresch, B.; Munoz, K.A.; Hillmann, H.; Sibrowski, W.; Humpf, H.U. Biomonitoring using dried blood spots: Detection of ochratoxin A and its degradation product 2'R-ochratoxin A in blood from coffee drinkers. *Mol. Nutr. Food Res.* 2015, 59, 1837–1843. [CrossRef] [PubMed]
- 79. Blanc, M.; Pittet, A.; Munoz-Box, R.; Viani, R. Behavior of Ochratoxin A during Green Coffee Roasting and Soluble Coffee Manufacture. *J. Agric. Food Chem.* **1998**, *46*, 673–675. [CrossRef]
- 80. Romani, S.; Pinnavaia, G.G.; Dalla Rosa, M. Influence of roasting levels on ochratoxin a content in coffee. *J. Agric. Food Chem.* **2003**, *51*, 5168–5171. [CrossRef] [PubMed]
- 81. Tsubouchi, H.; Terada, H.; Yamamoto, K.; Sakabe, Y. Ochratoxin A found in commercial roast coffee. *J. Agric. Food Chem.* **1988**, *36*, 540–542. [CrossRef]
- 82. van der Stegen, G.H.; Essens, P.J.; van der Lijn, J. Effect of roasting conditions on reduction of ochratoxin a in coffee. *J. Agric. Food Chem.* **2001**, *49*, 4713–4715. [CrossRef]
- Nehad, E.A.; Farag, M.M.; Kawther, M.S.; Abdel-Samed, A.K.; Naguib, K. Stability of ochratoxin A (OTA) during processing and decaffeination in commercial roasted coffee beans. *Food Addit. Contam.* 2005, 22, 761–767. [CrossRef]
- 84. Kumar, S.; Kunwar, A.; Gautam, S.; Sharma, A. Inactivation of A. ochraceus spores and detoxification of ochratoxin A in coffee beans by gamma irradiation. *J. Food Sci.* **2012**, *77*, T44–T51. [CrossRef]
- Chen, B.Y.; Huang, H.W.; Cheng, M.C.; Wang, C.Y. Influence of high-pressure processing on the generation of gamma-aminobutyric acid and microbiological safety in coffee beans. J. Sci. Food Agric. 2018, 98, 5625–5631. [CrossRef]
- 86. Bhupathiraju, S.N.; Pan, A.; Malik, V.S.; Manson, J.E.; Willett, W.C.; van Dam, R.M.; Hu, F.B. Caffeinated and caffeine-free beverages and risk of type 2 diabetes. *Am. J. Clin. Nutr.* **2013**, *97*, 155–166. [CrossRef]
- 87. Liu, H.; Zhou, Y.; Tang, L. Caffeine induces sustained apoptosis of human gastric cancer cells by activating the caspase9/caspase3 signalling pathway. *Mol. Med. Rep.* **2017**, *16*, 2445–2454. [CrossRef]
- 88. Prediger, R.D. Effects of caffeine in Parkinson's disease: From neuroprotection to the management of motor and non-motor symptoms. *J. Alzheimers Dis.* **2010**, *20* (Suppl. 1), S205–S220. [CrossRef]
- Tiwari, K.K.; Chu, C.; Couroucli, X.; Moorthy, B.; Lingappan, K. Differential concentration-specific effects of caffeine on cell viability, oxidative stress, and cell cycle in pulmonary oxygen toxicity in vitro. *Biochem. Biophys. Res. Commun.* 2014, 450, 1345–1350. [CrossRef]
- Akbar, A.; Medina, A.; Magan, N. Efficacy of different caffeine concentrations on growth and ochratoxin A production by *Aspergillus* species. *Lett. Appl. Microbiol.* 2016, 63, 25–29. [CrossRef]
- 91. Buchanan, R.L.; Tice, G.; Marino, D. Caffeine inhibition of ochratoxin A production. *J. Food Sci.* **1982**, 47, 319–321. [CrossRef]
- Tsubouchi, H.; Terada, H.; Yamamoto, K.; Hisada, K.; Sakabe, Y. Caffeine degradation and increased ochratoxin A production by toxigenic strains of Aspergillus ochraceus isolated from green coffee beans. *Mycopathologia* 1985, 90, 181–186. [CrossRef]
- 93. Cheng, B.; Furtado, A.; Henry, R.J. The coffee bean transcriptome explains the accumulation of the major bean components through ripening. *Sci. Rep.* **2018**, *8*, 11414. [CrossRef]
- 94. Di Stefano, V.; Pitonzo, R.; Cicero, N.; D'Oca, M.C. Mycotoxin contamination of animal feedingstuff: Detoxification by gamma-irradiation and reduction of aflatoxins and ochratoxin A concentrations. *Food Addit. Contam. Part. A Chem. Anal. Control Expo. Risk Assess.* 2014, *31*, 2034–2039. [CrossRef]
- 95. Kabak, B.; Dobson, A.D.; Var, I. Strategies to prevent mycotoxin contamination of food and animal feed: A review. *Crit Rev. Food Sci. Nutr.* **2006**, *46*, 593–619. [CrossRef]
- 96. Nascimento, L.C.; Lima, L.C.O.; Picolli, R.H.; Fiorini, J.E.; Duarte, S.M.S.; Silva, J.M.S.F.; Oliveira, N.M.S.; Veiga, S.M.O.M. Ozone and ultrasound: Alternative processes in the treatment of fermented coffee. *Food Sci. Technol.* 2008, 28, 282–294. [CrossRef]



© 2019 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).