

Article

A Hydrogel as a Bespoke Delivery Platform for Stromal Cell-Derived Factor-1

Yi Wang^{1,2}, Vanessa Penna³, Richard J. Williams⁴, Clare L. Parish^{3,†} and David R. Nisbet^{1,5,6,7,*}

¹ The Graeme Clark Institute, The University of Melbourne, Melbourne, Australia; yi.wang24@unimelb.edu.au

² Department of Biomedical Engineering, Faculty of Engineering and Information Technology, The University of Melbourne, Melbourne, Australia

³ The Florey Institute of Neuroscience and Mental Health, The University of Melbourne, Parkville, VIC 3052, Australia; vanessa.penna@florey.edu.au (V.P.); clare.parish@florey.edu.au (C.L.P.)

⁴ Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Deakin University, Warun Ponds, VIC 3216, Australia; richard.williams@deakin.edu.au

⁵ Laboratory of Advanced Biomaterials, Research School of Chemistry and the John Curtin School of Medical Research, The Australian National University, Canberra 2601, Australia

⁶ Melbourne Medical School, Faculty of Medicine, Dentistry and Health Science, The University of Melbourne, Melbourne, Australia

⁷ Department of Biomedical Engineering, Faculty of Engineering and Information Technology, The University of Melbourne, Melbourne, Australia

* Correspondence: david.nisbet@unimelb.edu.au

† These authors contributed equally to this work.

Supporting information




Figure S1. Representative photomicrographs staining with GFP+, β III-tubulin+, DAPI+ and merged images in Cells group, Cells + sSDF group, and Cells + SAP-shSDF group, demonstrating immature neurons in all groups.