

Supplementary Materials: Zero-Field Splitting in Cyclic Molecular Magnet $\{\text{Cr}_8\text{Y}_8\}$: a High-Frequency ESR Study

Zhendong Fu ¹, Zhong-Wen Ouyang ^{2,*}, Qian-Cheng Luo ³, Yan-Zhen Zheng ^{3,*}, Wei Tong ⁴, Huanpeng Bu ¹, Hanjie Guo ¹ and Jin-Kui Zhao ^{1,5}

¹ Songshan Lake Materials Laboratory, Dongguan 523808, China

² Wuhan National High Magnetic Field Center, Huazhong University of Science and Technology, Wuhan 430074, China

³ Frontier Institute of Science and Technology (FIST), State Key Laboratory for Mechanical Behavior of Materials, School of Chemistry and School of Physics, Xi'an Jiaotong University, Xi'an 710054, China

⁴ Anhui Province Key Laboratory of Condensed Matter Physics at Extreme Conditions, High Magnetic Field Laboratory, Chinese Academy of Sciences, Hefei 230031, China

⁵ Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

* Correspondence: zhouyang@mail.hust.edu.cn (Z.-W.O.); zheng.yanzhen@xjtu.edu.cn (Y.-Z.Z.)




Figure S1. A Schematic representation of the energy levels for $S = 11$ and $S = 12$ as a function of magnetic field (H //easy axis). The allowed transitions for 154 GHz are marked with red arrows. The $S = 11$ and $S = 12$ levels are separated for the sake of clarity.