

Supplementary Table S1: Proportion of initial CK-MM concentration for LC, MD, and NB samples at time points 10, 20, 30, 40, and 50 in each storage condition. Results are calculated mean proportions of initial CK-MM concentrations with standard error (SE).

<i>Time Since Initial (Days)</i>	<i>LC Samples</i>				<i>MD Samples</i>				<i>NB Samples</i>	
	<i>Low Humidity</i>		<i>Ambient Humidity</i>		<i>Low Humidity</i>		<i>Ambient Humidity</i>		<i>Low Humidity</i>	
	<i>Proportion of Initial</i>	<i>SE</i>	<i>Proportion of Initial</i>	<i>SE</i>	<i>Proportion of Initial</i>	<i>SE</i>	<i>Proportion of Initial</i>	<i>SE</i>	<i>Proportion of Initial</i>	<i>SE</i>
10	1.0683	0.0394	1.0247	0.0455	0.9331	0.0149	0.9039	0.0201	0.9184	0.0079
20	0.8675	0.0330	0.8630	0.0453	0.9520	0.0189	0.8508	0.0233	0.9065	0.0094
30	0.9655	0.0375	0.8757	0.0361	0.9151	0.0255	0.8291	0.0293	0.8614	0.0072
40	0.8947	0.0519	0.8365	0.0427	0.9791	0.0330	0.8891	0.0278	0.8714	0.0068
50	1.0513	0.0542	0.7884	0.0600	0.8875	0.0275	0.7666	0.0259	0.8159	0.0081

Supplementary Table S2: Proportion of initial CK-MM concentration for LC, MD, and NB samples at time points 2, 4, 6, and 8 days in high temperature and high humidity storage. Results are calculated mean proportions of initial CK-MM concentrations with standard error (SE).

LC Samples			MD Samples		NB Samples	
Time Since Initial (Days)	Proportion of Initial	SE	Proportion of Initial	SE	Proportion of Initial	SE
2	0.9483	0.0380	0.8735	0.0246	1.0167	0.0240
4	0.8182	0.0269	0.8143	0.0259	0.9782	0.0227
6	0.8624	0.0337	0.8213	0.0314	0.8826	0.0199
8	0.8467	0.0343	0.7643	0.0257	0.7944	0.0182