

## Review

# Herbicides versus Nitrogen Cycle: Assessing the Trade-Offs for Soil Integrity and Crop Yield—An In-Depth Systematic Review

Maura Gabriela da Silva Brochado , Laryssa Barbosa Xavier da Silva, Alessandro da Costa Lima ,  
Yure Marin Guidi  and Kassio Ferreira Mendes \* 

Department of Agronomy, Federal University of Viçosa, Viçosa 36570-900, MG, Brazil;  
maura.brochado@ufv.br (M.G.d.S.B.); laryssa.b.silva@ufv.br (L.B.X.d.S.); alessandro.lima@ufv.br (A.d.C.L.);  
yure.guidi@ufv.br (Y.M.G.)

\* Correspondence: kfmendes@ufv.br

**Abstract:** The interaction of herbicides in the nitrogen cycle and their consequences on soil health and agricultural production are essential topics in agronomic research. In this systematic review article, we have synthesized recent studies on this subject. The results revealed that the indiscriminate use of herbicides can have negative effects on vital processes in the nitrogen cycle, such as reduced enzymatic activity and microbial respiration. Moreover, herbicides alter the soil microbial composition, affecting nitrogen cycling-related activities. Symbiotic nitrogen fixation is also impaired, resulting in a reduction in the population of nitrogen-fixing bacteria and a decrease in the availability of this nutrient in the soil. These effects compromise soil fertility and the release of nitrogen to plants. Therefore, sustainable agricultural practices must be adopted, considering nitrogen cycling efficiency and the preservation of soil and natural resources. This understanding is crucial for guiding appropriate management strategies aimed at minimizing the negative effects of herbicides on the nitrogen cycle and ensuring soil health and agricultural productivity.

**Keywords:** soil health; sustainable agricultural practices; microbial respiration; enzymatic activity; symbiotic nitrogen fixation



**Citation:** Brochado, M.G.d.S.; Silva, L.B.X.d.; Lima, A.d.C.; Guidi, Y.M.; Mendes, K.F. Herbicides versus Nitrogen Cycle: Assessing the Trade-Offs for Soil Integrity and Crop Yield—An In-Depth Systematic Review. *Nitrogen* **2023**, *4*, 296–310. <https://doi.org/10.3390/nitrogen4030022>

Academic Editor: Angel Llamas

Received: 20 July 2023

Revised: 10 August 2023

Accepted: 29 August 2023

Published: 1 September 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

In order to increase productivity, the use of pesticides becomes necessary in agricultural practices. Pesticides are chemical compounds used to control pests, fungi, and weeds in crops [1,2]. Among these pesticides, herbicides are of great importance. They are crucial for effective weed control, as weeds compete with agricultural crops for essential resources, resulting in reduced productivity [3]. By eliminating or selectively suppressing weeds, herbicides keep crops weed-free and contribute to increased food production [1].

However, it is important to note that excessive use of these products can have negative impacts on the environment and vital cycles for agriculture, as well as consequences for human health. This has prompted the development of more sustainable and ecological agricultural practices [3,4].

Among the vital cycles in agriculture, nitrogen cycling is a fundamental process in crop development but can be negatively affected by the excessive use of herbicides, leading to direct toxicity [4–7]. Crucial microorganisms involved in the nitrogen cycle, such as nitrifying and nitrogen-fixing bacteria, are impacted by the toxic effects of certain herbicides, resulting in damage, mortality, and impairment of vital processes. In a study conducted by Crouzet et al. [6], it was found that high doses of the herbicide mesotrione were associated with a decrease in enzymatic activity and microbial respiration rates, indicating a reduction in the activity of key microorganisms in the nitrogen cycle. Additionally, the use of herbicides can modify the soil microbial composition, promoting the selective growth of certain microbial groups at the expense of others, which affects microbial activity related to

nitrogen cycling, including processes such as nitrification and denitrification [8]. In this regard, Du et al. [4] observed changes in the composition of soil microbial communities after mesotrione application, resulting in alterations in the structure and diversity of microbial communities important for nitrogen cycling.

The symbiotic nitrogen fixation is an indispensable process in the nitrogen cycle, where bacteria of the genus *Rhizobium* establish symbiotic associations with leguminous plants, such as the common bean (*Phaseolus vulgaris*) [9], soybean (*Glycine max*) [10] and pea (*Pisum sativum*) [11]. These bacteria have the ability to convert atmospheric nitrogen into a form assimilable by plants [12]. Studies have shown that diclosulam and glyphosate were able to reduce the population of rhizobia in soil by 89% and 93%, respectively [5]. This nitrogen fixation is an important source of nitrogen availability in the soil, and its reduction can adversely affect soil fertility [7,12,13]. Another implication is the alteration of organic residue decomposition, in which herbicides can exert an unfavorable influence [8]. These changes can affect the release of nitrogen from residues and its subsequent availability to plants [8,14]. Consequently, changes in decomposition processes can lead to reduced nitrogen cycling in the soil [14].

Therefore, understanding the impacts of herbicides on the nitrogen cycle and their consequences for soil health, agricultural efficiency, and environmental preservation is of utmost importance for the scientific community, academics, producers, and researchers. This systematic review article plays a crucial role in synthesizing and analyzing the most recent studies, highlighting the need for integrated approaches that consider efficient nitrogen cycling, soil quality preservation and protection of natural resources.


## 2. Interaction between Herbicides and the Nitrogen Cycle: Negative Impacts on Soil Health and Crop Yield

The interactions between herbicides and the nitrogen cycle involve complex processes of chemical, biological, and physical nature that can negatively affect the availability and cycling of this essential element in the soil [15]. From a molecular standpoint, herbicides can interact with nitrogen compounds, leading to their degradation or volatilization, thereby reducing the accessibility of these nutrients for plants [16]. Biologically, herbicides can interfere with the activity of nitrogen-fixing bacteria, diminishing their ability to convert atmospheric nitrogen into usable forms for plants [17]. This inhibition of biological nitrogen fixation directly affects the viability of this nutrient in the agricultural ecosystem [7]. Additionally, from a physical perspective, excessive herbicide application can cause modifications in soil structure, compaction and reduced porosity, leading to detrimental impacts on soil health [18] and the activity of nitrogen-fixing bacteria [19].

In order to provide a basis for decision-making in agricultural management, it is of utmost importance to understand the context of the interaction between the nitrogen cycle and herbicides.

### 2.1. Mechanism of Interaction

Interaction mechanisms play a pivotal role in crop development, exerting a critical influence on the intricate processes that orchestrate the nitrogen cycle. This intricate web of interactions encompasses Biological Nitrogen Fixation (BNF), carried out by both symbiotic bacteria [20] and those operating in a non-symbiotic manner [21], catalyzing the incorporation of nitrogen into the soil (Figure 1A,B). In parallel, nitrification plays a crucial role in transforming organic nitrogen compounds into readily accessible inorganic forms for plants (Figure 1E), while denitrification emerges as a key process, releasing atmospheric nitrogen from nitrogenous compounds (Figure 1H,I) [22]. Subsequent nitrogen assimilation by plants emerges as an essential process, playing a central role in the robust growth and development of these plant organisms [23] (Figure 1F).



**Figure 1.** Nitrogen cycle in the environment. (A) Nitrogen fixation, (B) Nitrogen fixing bacteria, (C) Nitrifying bacteria, (D) Decomposition, (E) Nitrification, (F) Absorption, (H) Denitrifying bacteria, and (I) Denitrification. Source: Adapted from Geisler et al. [23].

All these mechanisms are essential for the proper and balanced functioning of the nitrogen cycle [23]. However, excessive and improper use of herbicides can potentially damage the efficiency of nitrogen cycling [15]. In a study conducted by Du et al. [24], it was observed that the presence of mesosulfuron-methyl had an effect on soil biodegradability, negatively influencing the capacity for decomposition and degradation of organic compounds in the soil. This interference is directly related to nitrogen transformation in the soil, encompassing crucial mechanisms such as nitrification, denitrification, and biological nitrogen fixation (Figure 1A–C,H,I).


On the other hand, in a study by Hungria et al. [25], it was found that glyphosate had adverse effects on soybean plant nodulation. Nodulation is a crucial symbiotic mechanism for legumes, where an association occurs between plants and nitrogen-fixing bacteria, which are important for nitrogen fixation in the soil. Although the herbicide interfered with nodulation, the authors did not observe any effects of glyphosate on grain production.

These studies reveal the intrinsic complexity of these mechanisms for accurately assessing the effects of herbicides on nitrogen cycling dynamics. Such effects can have negative impacts on crop development, resulting in reduced nitrogen availability in the soil.

#### 2.1.1. Nitrogen Fixation Inhibition

Nitrogen fixation in the soil is a biological process in which diazotrophic bacteria associated with leguminous plants, such as *Rhizobium* spp., have the ability to reduce atmospheric nitrogen ( $N_2$ ) into ammonia ( $NH_3$ ) through the enzyme nitrogenase (Figure 2) [26]. Ammonia can then be converted into ammonium ions ( $NH_4^+$ ) available for plants. For

instance, the symbiosis between beans and the bacterium *Rhizobium leguminosarum* results in the formation of nodules on the roots, where nitrogen fixation occurs (Figure 2) [27]. This process is crucial for providing nitrogen to plants, enhancing soil fertility, and contributing to agricultural productivity. Additionally, nitrogen fixation reduces the dependence on synthetic nitrogen fertilizers, which has direct implications for the environmental sustainability of agriculture.



**Figure 2.** Nitrogen fixation and root nodule reduction in the presence of herbicide misuse. (A–C) Rhizobial nitrogen-fixing bacteria invade the roots, (D) Undergo multiplication, (E) Form root nodules, establishing a symbiotic relationship. Source: Adapted from Threatt et al. [26], Barberie et al. [27], Zablotowicz and Reddy [28], and Delong [29].

However, nitrogen fixation in non-leguminous crops differs from that in leguminous crops, as non-leguminous plants lack the intrinsic ability to form root nodules housing nitrogen-fixing bacteria, as seen in legumes (Figure 2). Instead, nitrogen fixation in non-leguminous crops involves symbiotic interactions with diazotrophic bacteria colonizing the rhizosphere (the soil region surrounding roots) to provide nitrogen to the host plants [30,31].

These bacteria facilitate nitrogen fixation through the enzyme nitrogenase, which catalyzes the conversion of  $N_2$  into usable nitrogenous compounds [31]. In many cases, optimizing nitrogen fixation in non-leguminous crops can be achieved through sustainable agricultural practices, such as crop rotation and cover cropping. Introducing specific cover crops that host diazotrophic bacteria can enhance nitrogen availability in the soil for primary crops [30]. Moreover, due to the absence of symbiotic association in non-leguminous crops, the regular application of nitrogen fertilizers is commonly employed as a conventional approach to meet nutritional demands.

Some herbicides can inhibit the activity of nitrogen-fixing bacteria present in plant roots, resulting in reduced nitrogen availability in the soil (Figure 2) [15]. A study conducted by Chen et al. [32] investigated the effects of butachlor at doses of 0.15 and 1.5 kg ha<sup>-1</sup> on the diversity of diazotrophic bacteria in the soil. The results revealed changes in the diversity of these bacteria in response to exposure to butachlor. Initially, the authors observed a suppression in nitrogen fixation activity, as evidenced by a reduction in acetylene production. However, in a subsequent stage, an increase in the suppression of this activity was observed. These findings suggest that butachlor interferes with the soil diazotrophic bacterial community, negatively affecting its nitrogen fixation capacity. Similar results were found in the study by Angelini et al. [17], where S-metolachlor, diclosulam, glyphosate, imazethapyr, and imazapic were applied to peanuts, resulting in a reduction in the diversity and abundance of nitrogen-fixing bacteria in the soil. Additionally, a pronounced decrease in nitrogen fixation activity by these bacteria was observed (Figure 2). These results indicated that exposure to the mentioned herbicides has negative effects on the community of nitrogen-fixing bacteria, reducing nitrogen availability for peanut plants.

Despite the importance of the nitrogen fixation process in providing plants with a usable form of nitrogen, it is also crucial to recognize the significance of denitrification in maintaining the balance of this essential element in the soil–plant–environment relationship.

#### 2.1.2. Inhibition of Denitrifying Bacteria

Denitrification is an essential step in the nitrogen cycle, where the bacteria transform nitrate (NO<sub>3</sub><sup>-</sup>) into gaseous nitrogen (N<sub>2</sub>), releasing it into the atmosphere [33–36]. Some bacteria, such as *Pseudomonas denitrificans* [37,38] and *Paracoccus denitrificans* [39,40], are mainly responsible for this process. These bacteria are able to utilize nitrate as an alternative energy source under anaerobic conditions [23,33,34].

The inhibition of denitrifying bacteria can occur due to various factors, such as the presence of toxic chemicals in the soil, including herbicides [36]. These substances can interfere with the metabolism of denitrifying bacteria, compromising their ability to efficiently carry out denitrification [40].

The inhibition of these bacteria can have significant implications on the nitrogen cycle and nutrient balance in the soil. This can lead to the accumulation of nitrate in the soil and a reduction in nitrogen availability for plants. The excessive accumulation of nitrate in the soil can disrupt the absorption of other nutrients by plants, destabilizing the nutritional equilibrium and consequently diminishing the effective availability of essential nitrogen for healthy growth. This phenomenon arises from ion antagonism, which affects nutrient transport processes in plant roots. As observed by Yu et al. [14], the application of acetochlor at a concentration of 10 mg kg<sup>-1</sup> resulted in the inhibition of soil denitrification potentials. This inhibition could be attributed to a decrease in both the abundance and activity of denitrifying bacteria after the application of acetochlor. Similar results were found by Crouze et al. [41], where mesotrione increased the ammonium content in the soil and caused a decrease in nitrate content observed in treated soils. These functional impacts were mainly correlated with changes in the abundance of oxidizing bacteria or denitrifying bacteria.

Since herbicides have the potential to inhibit vital processes, one of the main impacts is the reduction of nutrient absorption by plants. This decrease in absorption can result in various issues, such as nutrient deficiency in the plant and, consequently, a decrease in productivity.

#### 2.1.3. Reduced Nitrogen Uptake by Plants

The reduction in nitrogen uptake by plants can be attributed to multiple factors, including decreased soil microorganism activity, direct interference of herbicides with plant membrane transporters, and modulation of enzymes involved in nitrogen metabolism [42]. Nitrogen uptake by plants is a complex phase and vital for their growth and development. It involves the uptake of nitrate (NO<sub>3</sub><sup>-</sup>) and ammonium (NH<sub>4</sub><sup>+</sup>) from the soil by the



roots, followed by transport to other parts of the plant [43]. This process primarily takes place in the roots, where membrane transporters play a pivotal role by facilitating the selective transport of nitrogen ions [44]. The reliance on these transporters is pivotal for the effectiveness of nitrogen uptake, exerting a substantial influence on nutritional equilibrium and plant productivity [44].

Nitrogen assumes a vital role for plants, fulfilling a dual function in synthesizing pivotal compounds, including amino acids, proteins, nucleotides, and other nitrogenous substances within plant cells [44]. These essential compounds significantly contribute to plant structure and functionality, encompassing critical processes such as growth, development, reproduction, and the plant's adaptive response to environmental stresses. Furthermore, the potential compromise in plant growth due to the reduction of assimilable nitrogen forms may consequently impact the entirety of these fundamental functions. [23,44].

Furthermore, herbicides can interfere with membrane transporters, which are responsible for the active or passive transport of nitrogen ions within plant cells. This interference can compromise the efficiency of nitrogen uptake, affecting the availability of this essential nutrient for plants [42].

Another noteworthy aspect is the potential influence of herbicides on nitrogen metabolism enzymes in plants. Inhibition of these enzymes could impact the assimilation of nitrate and ammonium, impairing the conversion of these compounds into usable forms for amino acid and protein synthesis [45]. This would result in an imbalance in nitrogen metabolism, directly impacting plant growth and development [45].

Therefore, the reduction in nitrogen uptake may be attributed to a combination of factors, including decreased microbial activity in the soil, direct interference with plant membrane transporters, and disruptions in enzymes involved in nitrogen metabolism.

Due to the impact of herbicides on processes essential for nitrogen availability [15], significant interference with plant development occurs. This interference includes the reduction of the bacterial population responsible for converting the element into a form assimilable by plants and the formation of root nodules [28]. According to Zablotowicz and Reddy [28], glyphosate interfered with the symbiosis between soybean and nitrogen-fixing bacteria, compromising the uptake of this nutrient by the crop. In addition, Singh and Wright [46], when evaluating terbutylazine, simazine, prometryn, and bentazone in pea crops, observed a decrease in plant growth due to the impairment of root nodule formation, which made it impossible to absorb nitrogen from the soil.

#### 2.1.4. Changing the Rate of Organic Matter Decomposition

Soil organic matter (SOM) is made up of a variety of organic compounds, both polymers and monomers, which exhibit different sizes and levels of decomposition. These compounds interact with their surrounding environment, and it is these physicochemical interactions that play a key role in microbial dynamics in the rhizosphere [47]. These interactions are responsible, for example, for the decomposition of organic molecules and, consequently, for their persistence in the soil [48,49]. Interactions between herbicides and soil organic compounds can occur through several physical forces, such as hydrogen bonds, van der Waals forces, electrostatic forces, covalent bonds and hydrophobic interactions. These interactions act simultaneously on the sorption of herbicide molecules in soil colloids [50,51].

The use of organic compounds in the soil presents different responses in the retention of herbicides. For example, Mendes et al. [52] evaluated the addition of cow bone char on sorption–desorption and mobility of hexazinone, metribuzin and quinclorac applied to sandy loam soil under laboratory conditions. The results obtained proved that bone char was an excellent sorbent for reducing the mobility of the three herbicides due to the high sorption in the soil. On the other hand, Prata et al. [53] evaluated the effect of stillage addition on the behavior of diuron and ametryn in sandy and clayey soils. The authors reported that the addition of stillage did not affect the sorbed amount of these molecules

and did not result in changes in the organic carbon content of the soil after four days of application.

The addition of nitrogen sources can alter the decomposition rate of SOM. Nitrogen availability directly affects microbial activity in the soil, being a key factor in the decomposition of SOM [47]. Li et al. [54] reported that urea fertilization decreased the decomposition of SOM and maize straw, evidenced by the higher nitrogen and carbon content compared to soil without urea.

In addition, the application of mixtures with herbicides and nitrogen fertilizers can influence the development characteristics of plant species. Dupont et al. [55] evaluated the isolated and combined effect of glyphosate and nitrogen doses on *Tanacetum vulgare* plants. Based on the results obtained, the authors showed that the application of glyphosate nitrogen and their interactions affected the reproductive characteristics of the plants evaluated, mainly the density and flowering phenology. Flowering was severely delayed by glyphosate application (10.5 days delay per 100 g a.e. ha<sup>-1</sup> year<sup>-1</sup>). Although nitrogen partially attenuated the reduction in floral abundance, the delay in flowering was amplified when nitrogen was added.

The use of nitrogen fertilizers can also interfere with the selectivity of herbicides. Langaro et al. [56] evaluated the selectivity of herbicides according to the time of nitrogen application in irrigated rice. The results obtained proved that the application of nitrogen, in general, resulted in an increase in the height and dry matter of rice plants, and the application of nitrogen before waterlogging and split applications (50% before and 50% after waterlogging) were favorable to the crop, consequently, the selectivity of the crop to herbicides. Bispyribac-sodium resulted in the greatest damage and reduction in rice plant height, followed by bentazone and carfentrazone-ethyl, while the lowest damage and highest grain yield were obtained with the quinclorac application.

Therefore, the addition of nitrogen sources to soils is still a point to be elucidated by the scientific literature in view of the variety of responses listed so far. The decomposition of SOM by providing an additional supply of nitrogen to soil microorganisms. This may result in a more rapid release of nutrients contained in the decomposing SOM, making them available to plants in a more readily usable form, thus influencing the behavior of herbicides in the soil.

#### 2.1.5. Change in the Rate of Nitrogen Fixation by Symbiotic Bacteria

The use of symbiotic bacteria applied as inoculants can meet the nitrogen needs of the plant. Bacteria of the genus *Bradyrhizobium* are the most used in agriculture due to the high world production of soybeans, enabling the reduction in the use of industrialized nitrogen fertilizers [57].

However, the increased use of pesticides, especially herbicides, can interfere with the process of biological nitrogen fixation by plants, both with positive and negative effects. The positive effect is observed when herbicides stimulate the growth and development of the fixation nodules. On the other hand, the negative effect is manifested in the reduction of nitrogen fixation efficiency, resulting in lower plant growth and yield [58,59].

After application, herbicides may come into contact with rhizobia immediately or, in the case of herbicides with residual effect, throughout the development of the main crop. These herbicides have the ability to impact the rhizobia, the host plant and also the establishment and development of the symbiosis [60]. The use of herbicides with a residual effect on the soil is a technique that enhances weed management, as it allows control in the pre-emergence of weeds [61]. However, the problem involved in the use of these herbicides is the possibility of carryover, that is, the presence of bioavailable residues of the herbicide in the soil, which can influence the microbial activity of the soil [62].

The behavior of herbicides in the soil is complex since the reduction of the population of a certain species of microorganisms can occur in contrast to the development of another population, which is not so affected by the toxic effects of the product. The species or individuals that survive start to use nutrients, such as carbon, sulfur and nitrogen, as

well as the chemical energy of the herbicide molecules, which are released during the degradation process. Barroso et al. [63] reported that *Bradyrhizobium* sp. strain BR 3901 was able to produce nitrogen to catalyze degradation reactions of diuron, sulfentrazone, oxyfluorfen, and 2,4-D.

In studies carried out by Vercellino and Gómez [64], the authors evaluated the growth parameters of 81 strains of different genera of rhizobia (*Rhizobium*, *Mesorhizobium*, *Ensifer* and *Bradyrhizobium*) when exposed to the application of glyphosate, 2,4-D and atrazine, in addition to the ability of these strains to degrade herbicides. The genera studied showed different responses to the herbicides evaluated, with the genus *Bradyrhizobium* showing a greater ability to transform herbicides into compounds of lower phytotoxicity. In addition, among the compounds evaluated, atrazine was the most used as an energy source for bacteria. The results obtained confirmed that *Bradyrhizobium* strains were able to both denitrify and use atrazine as an energy substrate.

The relationship between herbicide application and symbiotic nitrogen fixation may have different effects depending on the crop variety, as well as the symbiotic strains used. Bossolani et al. [65] evaluated the effect of glyphosate doses on *Bradyrhizobium* strains (*Bradyrhizobium elkanii*—SEMIA 5019—and *Bradyrhizobium japonicum*—SEMIA 5079) and biological nitrogen fixation in soybean plants (BMX Potência RR). The authors reported that the inoculated RR soybean did not present alteration in the leaf chlorophyll index by the application of glyphosate, and regardless of the Inoculation, the soybean plants were able to recover from the application of glyphosate, not impairing their development.

The influence of herbicides on the nodulation of nitrogen-fixing bacteria is not a specific point of the soybean crop. Paniagua-López et al. [2] reported that the application of pendimethalin and clethodim reduced the ability of *Phaseolus vulgaris* and *Medicago sativa* to fix nitrogen by inhibiting root growth and modifying the composition of root exudates, as well as rhizospheric bacterial fitness. The authors cited 30% reductions in nodulation after clethodim application, while pendimethalin totally inhibited nodulation, causing reduced growth and motility of nitrogen-fixing nodules.

In another study, Khan et al. [66] evaluated bentazone, isoproturon, fluchloralin and 2,4-D, applied to soil, on growth characteristics, chlorophyll contents, nitrogen and protein contents, nodulation and seed yield in chickpea inoculated with *Mesorhizobium ciceri*. Fluchloralin and 2,4-D caused negative effects on chickpea seed yield.

In the study carried out by Santos et al. [67], the growth of *Rhizobium tropici* strains BR 322 and BR 520, used as inoculants in bean cultivation in Brazil, was analyzed in culture medium based on mannitol and yeast extract added with bentazone, metolachlor, imazamox, fluazifop-p-butyl, fomesafen, and paraquat. The authors found that paraquat was the herbicide with the highest growth inhibition of the strains evaluated, followed by the commercial mixture of fomesafen and fluazifop-p-butyl. As for the other herbicides, there was no reduction in growth. Overall, strain BR 520 showed greater tolerance to the herbicides tested, except for paraquat.

Thus, an efficient alternative for the remediation of contaminated soils may be the symbiosis between phytoremediating plants and microorganisms that possess enzymatic activity capable of metabolizing herbicides. Several groups of microorganisms degrade and utilize herbicide residues as a source of carbon and nitrogen. Bioremediation agents can absorb and degrade herbicides in secondary metabolic pathways or stimulate the soil microbiota to promote the transformation of these contaminants into compounds with lower or no toxicity [68–70].

Inoculation of symbiotic bacteria can contribute to the phytoremediation of herbicides in soil. Some *Bradyrhizobium* strains have the potential to degrade herbicides in soils, which is important for environmental decontamination purposes. Barroso et al. [63] evaluated the tolerance and in vitro growth of *Bradyrhizobium* sp BR 3901 in media without carbon and nitrogen sources, exposed to 2,4-D, oxyfluorfen, clomazone, glufosinate-ammonium, atrazine, ametryn, glyphosate, sulfentrazone, and diuron). The results obtained showed a reduction in nodulation.




In the study by Mielke et al. [71], the potential for positive interactions between symbiotic microorganisms (*Bradyrhizobium* sp. BR 2003—SEMIA 6156) and *Canavalia ensiformis* was evaluated, aiming at the phytoremediation of the herbicide sulfentrazone in the soil. The authors highlighted that microbiological indicators showed satisfactory results, mainly for the dose of 400 g ha<sup>-1</sup>. Thus, the symbiosis between herbicide-metabolizing microorganisms and phytoremediating plants can be an efficient alternative to remediate soils contaminated with herbicides.

### 3. Perspective of Contemporary Literature Regarding the Effects of Herbicides on Nitrogen Cycle Dynamics and Agricultural Systems

#### 3.1. Distribution of Articles by Year and Cumulative Total

Figure 3 represents the number of papers published over time on the effect of herbicides on soil nitrogen. There has been a steady increase in the number of publications from 1960 to the present day. However, in the first decade analyzed (until 1970), there were few publications, only nine, and some years with no new publications. From the 1970s onwards, the number of articles increased but was still relatively low, totaling 50 until 1980. The 1990s saw a significant increase in the total number of publications, reaching 272, with a highlight being the 43 articles in 1989, which saw a ~5-fold increase in the total number of articles published each decade until 1990.




**Figure 3.** Articles on the effect of herbicides on soil nitrogen published by year (gray bars) and total accumulated (red line) between 1960 and 2023. Source: CAPES [72]; Clarivate [73]; Elsevier [74]; PMC [75].

Until 2000, there was a less expressive but significant increase in the total number of publications, totaling 890 articles (Figure 3). The year 1989 stood out with 77 new articles. In 2010, the total number of publications almost doubled, reaching 1716 publications. The decade from 2011 to 2020 was the period of greatest growth in the total number of publications, with a total of 2712 articles published, with moments with more than 100 articles published in a single year. The years 2021 and 2022 had the highest number of publications, with 143 and 173 articles published, respectively. By March 2023, 43 articles had already been published, suggesting that this year may have the highest number of publications in the historical series.

According to the data presented, it is evident that the topic of the effect of herbicides on soil nitrogen is widely addressed in the scientific literature and is at its research peak.

### 3.2. Distribution of Articles by Index Bases

Figure 4 illustrates the distribution of articles on the impact of herbicides on soil nitrogen published between 1960 and 2023 in various collections. The Science Citation Index Expanded, a subset within the Web of Science, has the highest number of published articles at 2114, suggesting that the scientific community has a significant interest in this area of research. This can be attributed to the fact that Web of Science covers a wide range of research areas and publishes in over 9500 journals, including those related to agriculture and environmental sciences, which are likely to feature studies on herbicides and their effects on soil nitrogen.



**Figure 4.** Articles on the effect of herbicides on soil nitrogen have been published in different collections. The range of the red area represents the number of articles. Source: CAPES [72]; Clarivate [73]; Elsevier [74]; PMC [75].


ScienceDirect is the second largest collection with 839 published articles, indicating its importance in this field (Figure 4), being known to be a vital source of research in science, technology and health and offering access to 4756 journals. PubMed, with 721 articles, is also widely used in the areas of health and biology, with 3998 journals, which may explain the relatively high number of publications on the effect of herbicides on soil nitrogen. Gale Academic OneFile, with 576 articles, and DOAJ Directory of Open Access Journals, with 569 articles, also have a substantial number of publications.

Other research platforms with a smaller number of articles on the effect of herbicides on soil nitrogen include Wiley Online Library Journals (491 articles), Springer Nature Journals (479 articles), Journals Ovid Complete (459 articles), EBSCO Academic Search Premier (307 articles), BioOne (242 articles), Taylor & Francis Journals Complete (210 articles), Highwire Press Free (156 articles), American Chemical Society Journals (ACS) (118 articles), Canadian Science Publishing (104 articles) and SciELO (69 articles) (Figure 4).

It is worth mentioning the presence of open-access platforms, such as the DOAJ Directory of Open Access Journals and SciELO, which can facilitate the democratization of scientific knowledge and broaden access to research on the effect of herbicides on soil nitrogen. Overall, these data suggest that research on the impact of herbicides on soil nitrogen is extensive, and there is considerable interest in this topic among the scientific community. In addition, several research platforms offer a wide range of sources to obtain information on the subject.

### 3.3. Distribution of Published Articles by Thematic Areas

Figure 5 indicates the distribution of published articles on the topic of herbicides and their effect on soil nitrogen, categorized by subject area. The highest number of published articles was in Science and Technology, Life Sciences, and Biomedicine, which suggests that this topic is of great interest to researchers in these areas.



**Figure 5.** Articles on the effect of herbicides on soil nitrogen indexed in different subjects. The range of the red area represents the number of articles published in each subject area. Source: CAPES [72]; Clarivate [73]; Elsevier [74]; PMC [75].

Agriculture also had a high number of articles published, which is not surprising given the importance of herbicides in weed control (Figure 5). It is noteworthy that the area of Herbicides had the fourth highest number of articles published, indicating that there is a considerable amount of research being conducted specifically on this topic.

Other areas with a significant number of published articles include Environmental Sciences and Ecology, Agronomy and Soil Science (Figure 5). This suggests that researchers in these areas are also interested in studying the effects of herbicides on soil nitrogen.

The area of Phytopathology, which deals with plant diseases and pests, also had a relatively high number of published articles (Figure 5). On the other hand, the fields of Physical Sciences, Chemistry and Pesticides had relatively low numbers of published articles (Figure 5). This may be because these fields may not have as much focus on soil and agriculture as other fields or that the effects of herbicides on soil nitrogen are not seen as a priority within these fields.

Overall, these data highlight the interest and importance of studying the effects of herbicides on soil nitrogen in a variety of different thematic areas. This suggests that there is still much to be learned about this topic and that researchers from different disciplines can contribute to a better understanding of the effects of herbicides on soil health.

#### 4. Concluding Remarks

Overall, this systematic review highlights the need to deepen our understanding of the interaction of herbicides in the nitrogen cycle and the resulting implications for soil health and crop yield. The findings obtained from reviewed studies suggest that herbicides can have diverse effects on the dynamics of the nitrogen cycle, influencing key processes, such as symbiotic fixation, denitrification, and nitrogen uptake by plants. These effects can compromise nutrient availability in the soil, leading to negative consequences for crop growth, development, and yield.

It is crucial to consider these interactions in agricultural practice by adopting good practices that minimize the negative impacts of herbicides on the nitrogen cycle. This may include the appropriate use of herbicides, selecting products with lower potential for interfering with nitrogen fixation and uptake, as well as embracing integrated weed management practices.

Furthermore, more studies are needed to deepen our knowledge of herbicide effects on the nitrogen cycle and to develop more efficient and sustainable management strategies. These research efforts are essential to ensure soil health, agricultural productivity, and food security, given the importance of nitrogen as an essential nutrient for plants.

**Author Contributions:** Literature revision, M.G.d.S.B., L.B.X.d.S., A.d.C.L. and K.F.M.; writing—original draft preparation, M.G.d.S.B., L.B.X.d.S., A.d.C.L., Y.M.G. and K.F.M.; writing—review and editing M.G.d.S.B., Y.M.G. and K.F.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Data Availability Statement:** No new data were created.

**Acknowledgments:** The authors would like to thank the *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPq) for PhD scholarship.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

- Sharma, A.; Kumar, V.; Shahzad, B.; Tanveer, M.; Sidhu, G.P.S.; Handa, N.; Kohli, S.K.; Yadav, P.; Bali, A.S.; Parihar, R.D.; et al. Worldwide Pesticide Usage and Its Impacts on Ecosystem. *SN Appl. Sci.* **2019**, *1*, 1446. [\[CrossRef\]](#)
- Paniagua-López, M.; Jiménez-Pelayo, C.; Gómez-Fernández, G.O.; Herrera-Cervera, J.A.; López-Gómez, M. Reduction in the Use of Some Herbicides Favors Nitrogen Fixation Efficiency in *Phaseolus vulgaris* and *Medicago Sativa*. *Plants* **2023**, *12*, 1608. [\[CrossRef\]](#) [\[PubMed\]](#)
- Mandal, A.; Sarkar, B.; Mandal, S.; Vithanage, M.; Patra, A.K.; Manna, M.C. Impact of Agrochemicals on Soil Health. In *Agrochemicals Detection, Treatment and Remediation*; Butterworth-Heinemann: Oxford, UK, 2020; pp. 161–187. [\[CrossRef\]](#)
- Du, P.; Wu, X.; Xu, J.; Dong, F.; Liu, X.; Zheng, Y. Effects of Trifluralin on the Soil Microbial Community and Functional Groups Involved in Nitrogen Cycling. *J. Hazard. Mater.* **2018**, *353*, 204–213. [\[CrossRef\]](#) [\[PubMed\]](#)
- Ampofo, J.A.; Tetteh, W.; Bello, M. Impact of Commonly Used Agrochemicals on Bacterial Diversity in Cultivated Soils. *Indian J. Microbiol.* **2009**, *49*, 223–229. [\[CrossRef\]](#) [\[PubMed\]](#)
- Crouzet, O.; Batisson, I.; Besse-Hoggan, P.; Bonnemoy, F.; Bardot, C.; Poly, F.; Bohatier, J.; Mallet, C. Response of Soil Microbial Communities to the Herbicide Mesotrione: A Dose-Effect Microcosm Approach. *Soil Biol. Biochem.* **2010**, *42*, 193–202. [\[CrossRef\]](#)
- Du, Z.; Zhu, Y.; Zhu, L.; Zhang, J.; Li, B.; Wang, J.; Wang, J.; Zhang, C.; Cheng, C. Effects of the Herbicide Mesotrione on Soil Enzyme Activity and Microbial Communities. *Ecotoxicol. Environ. Saf.* **2018**, *164*, 571–578. [\[CrossRef\]](#)
- Cao, J.; Zhang, Y.; Dai, G.; Cui, K.; Wu, X.; Qin, F.; Xu, J.; Dong, F.; Pan, X.; Zheng, Y. The Long-Acting Herbicide Mesosulfuron-Methyl Inhibits Soil Microbial Community Assembly Mediating Nitrogen Cycling. *J. Hazard. Mater.* **2023**, *443*, 130293. [\[CrossRef\]](#)
- Brito, M.d.M.P.; Muraoka, T.; Silva, E.C.d. Contribuição Da Fixação Biológica de Nitrogênio, Fertilizante Nitrogenado e Nitrogênio Do Solo No Desenvolvimento de Feijão e Caupi. *Bragantia* **2011**, *70*, 206–215. [\[CrossRef\]](#)
- Cunha, L.d.S.; Duarte Júnior, J.B.; Lana, M.d.C.; Ribeiro, L.L.O.; Shimada, B.S.; Richart, A.; Costa, A.C.T.d.; Rosa, W.B. Inoculation, Co-Inoculation and Nitrogen Fertilization in Soybean Culture. *Concillium* **2023**, *23*, 454–472. [\[CrossRef\]](#)

11. Kamran, A.; Mushtaq, M.; Arif, M.; Rashid, S. Role of Biostimulants (Ascorbic Acid and Fulvic Acid) to Synergize Rhizobium Activity in Pea (*Pisum sativum* L. Var. Meteor). *Plant Physiol. Biochem.* **2023**, *196*, 668–682. [CrossRef]
12. Dixon, R.; Kahn, D. Genetic Regulation of Biological Nitrogen Fixation. *Nat. Rev. Microbiol.* **2004**, *2*, 621–631. [CrossRef] [PubMed]
13. Xiang, L.; Harindintwali, J.D.; Wang, F.; Bian, Y.; Zhao, Z.; Wang, Z.; Wang, Y.; Mei, Z.; Jiang, X.; Schäffer, A.; et al. Manure- and Straw-Derived Biochars Reduce the Ecological Risk of PBDE and Promote Nitrogen Cycling by Shaping Microbiomes in PBDE-Contaminated Soil. *Chemosphere* **2023**, *312*, 137262. [CrossRef] [PubMed]
14. Yu, J.; Zhang, J.; Zheng, X.; Zhang, Y.; Chen, D.; Ding, H. Divergent Modulation of Land Use-Driven Changes in Soil Properties and Herbicide Acetochlor Application on Soil Nitrogen Cycling. *Soil Tillage Res.* **2022**, *215*, 105231. [CrossRef]
15. Rose, M.T.; Cavnagaro, T.R.; Scanlan, C.A.; Rose, T.J.; Vancov, T.; Kimber, S.; Kennedy, I.R.; Kookana, R.S.; Van Zwieten, L. Impact of Herbicides on Soil Biology and Function. *Adv. Agron.* **2016**, *136*, 133–220. [CrossRef]
16. Carmo, J.B.d.; Andrade, C.A.d.; Cerri, C.C.; Piccolo, M.d.C. Disponibilidade de Nitrogênio e Fluxos de N<sub>2</sub>O a Partir de Solo Sob Pastagem Após Aplicação de Herbicida. *Rev. Bras. Ciência Solo* **2005**, *29*, 735–746. [CrossRef]
17. Angelini, J.; Silvina, G.; Taurian, T.; Ibáñez, F.; Tonelli, M.L.; Valetti, L.; Anzuay, M.S.; Ludueña, L.; Muñoz, V.; Fabra, A. The Effects of Pesticides on Bacterial Nitrogen Fixers in Peanut-Growing Area. *Arch. Microbiol.* **2013**, *195*, 683–692. [CrossRef]
18. Mishra, R.K.; Mohammad, N.; Roychoudhury, N. Soil Pollution: Causes, Effects and Control. *Trop. For. Res. Inst.* **2015**, *3*, 20–30.
19. Cycoń, M.; Wójcik, M.; Borymski, S.; Piotrowska-Seget, Z. Short-Term Effects of the Herbicide Napropamide on the Activity and Structure of the Soil Microbial Community Assessed by the Multi-Approach Analysis. *Appl. Soil Ecol.* **2013**, *66*, 8–18. [CrossRef]
20. Gonzalez, A.; Gonzalez-Murua, C.; Royuela, M. Influence of Imazethapyr on Rhizobium Growth and Its Symbiosis with Pea (*Pisum sativum*). *Weed Sci.* **1996**, *44*, 31–37. [CrossRef]
21. Kennedy, I. Non-Symbiotic Bacterial Diazotrophs in Crop-Farming Systems: Can Their Potential for Plant Growth Promotion Be Better Exploited? *Soil Biol. Biochem.* **2004**, *36*, 1229–1244. [CrossRef]
22. Bloom, A.J. The Increasing Importance of Distinguishing among Plant Nitrogen Sources. *Curr. Opin. Plant Biol.* **2015**, *25*, 10–16. [CrossRef] [PubMed]
23. Geisler, G.; Lea, P.J.; Morot-Gaudry, J.-F. (Eds.) 2001: Plant Nitrogen. *J. Agron. Crop Sci.* **2003**, *189*, 197–198. [CrossRef]
24. Du, P.; He, H.; Wu, X.; Xu, J.; Dong, F.; Liu, X.; Zheng, Y. Mesosulfuron-Methyl Influenced Biodegradability Potential and N Transformation of Soil. *J. Hazard. Mater.* **2021**, *416*, 125770. [CrossRef] [PubMed]
25. Hungria, M.; Mendes, I.C.; Nakatani, A.S.; dos Reis-Junior, F.B.; Morais, J.Z.; de Oliveira, M.C.N.; Fernandes, M.F. Effects of the Glyphosate-Resistance Gene and Herbicides on Soybean: Field Trials Monitoring Biological Nitrogen Fixation and Yield. *Field Crops Res.* **2014**, *158*, 43–54. [CrossRef]
26. Threatt, S.D.; Rees, D.C. Biological Nitrogen Fixation in Theory, Practice, and Reality: A Perspective on the Molybdenum Nitrogenase System. *FEBS Lett.* **2022**, *597*, 45–58. [CrossRef]
27. Barbieri, P.; Starck, T.; Voisin, A.-S.; Nesme, T. Biological Nitrogen Fixation of Legumes Crops under Organic Farming as Driven by Cropping Management: A Review. *Agric. Syst.* **2023**, *205*, 103579. [CrossRef]
28. Zablotowicz, R.M.; Reddy, K.N. Impact of Glyphosate on the Symbiosis with Glyphosate-Resistant Transgenic Soybean. *J. Environ. Qual.* **2004**, *33*, 825. [CrossRef]
29. Delong, G. Soybean Root Structure with Nodule. Available online: <https://pixels.com/featured/soybean-root-structure-with-nodule-garry-delong.html> (accessed on 18 May 2023).
30. Chalk, P.M. The Contribution of Associative and Symbiotic Nitrogen Fixation to the Nitrogen Nutrition of Non-Legumes. *Plant Soil* **1991**, *132*, 29–39. [CrossRef]
31. Kennedy, I.R.; Tchan, Y.-T. Biological Nitrogen Fixation in Non-Leguminous Field Crops: Recent Advances. *Plant Soil* **1992**, *141*, 93–118. [CrossRef]
32. Chen, W.C.; Yen, J.H.; Chang, C.S.; Wang, Y.S. Effects of Herbicide Butachlor on Soil Microorganisms and on Nitrogen-Fixing Abilities in Paddy Soil. *Ecotoxicol. Environ. Saf.* **2009**, *72*, 120–127. [CrossRef]
33. Yeomans, J.C.; Bremner, J.M. Denitrification in Soil: Effects of Herbicides. *Soil Biol. Biochem.* **1985**, *17*, 447–452. [CrossRef]
34. Bollag, J.-M.; Henninger, N.M. Influence of Pesticides on Denitrification in Soil and with an Isolated Bacterium. *J. Environ. Qual.* **1976**, *5*, 15–18. [CrossRef]
35. Pell, M.; Stenberg, B.; Stenström, J.; Torstensson, L. Potential Denitrification Activity Assay in Soil—With or without Chloramphenicol? *Soil Biol. Biochem.* **1996**, *28*, 393–398. [CrossRef]
36. Gulhane, P.A.; Ashok, V.G.; Kajal, M.S. Influence of pesticides on nitrogen fixing bacteria. *Int. J. Tech. Res. Appl.* **2015**, *3*, 157–160.
37. Koike, I.; Hattori, A. Growth Yield of a Denitrifying Bacterium, *Pseudomonas Denitrificans*, under Aerobic and Denitrifying Conditions. *J. Gen. Microbiol.* **1975**, *88*, 1–10. [CrossRef]
38. Grenier, V.; Laur, J.; Gonzalez, E.; Pitre, F.E. Glyphosate Has a Negligible Impact on Bacterial Diversity and Dynamics during Composting. *Environ. Microbiol.* **2023**, early view. [CrossRef]
39. Wan, X.; Wan, G.; Snozzi, M. Microbiological Denitrification and Denitrifying Activity of *Paracoccus Denitrificans*. *Chin. J. Geochem.* **2000**, *19*, 186–192. [CrossRef]
40. Su, G.; Chen, B.; Wu, X.; Xu, J.; Yang, K.; Lin, D. nZVI Decreases N<sub>2</sub>O Emission from Pesticide-Contaminated Paddy Soil. *Sci. Total Environ.* **2023**, *892*, 164613. [CrossRef]
41. Crouzet, O.; Wiszniowski, J.; Donnadiou, F.; Bonnemoy, F.; Bohatier, J.; Mallet, C. Dose-Dependent Effects of the Herbicide Mesotrione on Soil Cyanobacterial Communities. *Arch. Environ. Contam. Toxicol.* **2012**, *64*, 23–31. [CrossRef]



42. Forde, B.G.; Clarkson, D.T. Nitrate and Ammonium Nutrition of Plants: Physiological and Molecular Perspectives. *Adv. Bot. Res.* **1999**, *30*, 1–90. [\[CrossRef\]](#)
43. Abd-Alla, M.H.; Omar, S.A.; Karanxha, S. The Impact of Pesticides on Arbuscular Mycorrhizal and Nitrogen-Fixing Symbioses in Legumes. *Appl. Soil Ecol.* **2000**, *14*, 191–200. [\[CrossRef\]](#)
44. Taiz, L.; Zeiger, E.; Santarém, E.R. *Fisiologia Vegetal*, 4th ed.; Artmed: Porto Alegre, Brazil, 2009; Volume 1.
45. Patnaik, G.K.; Bose, L.K.; Mehta, A.M.; Rajaramamohan Rao, V. Rhizosphere Nitrogenase and *Azospirillum* sp. Association with Wild, Trisomic and Cultivated Rice. *Microbiol. Res.* **1994**, *149*, 42–46. [\[CrossRef\]](#)
46. Singh, G.; Wright, D. Effects of Herbicides on Nodulation, Symbiotic Nitrogen Fixation, Growth and Yield of Pea (*Pisum sativum*). *J. Agric. Sci.* **1999**, *133*, 21–30. [\[CrossRef\]](#)
47. Angst, G.; Mueller, K.E.; Nierop, K.G.J.; Simpson, M.J. Plant- or Microbial-Derived? A Review on the Molecular Composition of Stabilized Soil Organic Matter. *Soil Biol. Biochem.* **2021**, *156*, 108189. [\[CrossRef\]](#)
48. Schmidt, M.W.I.; Torn, M.S.; Abiven, S.; Dittmar, T.; Guggenberger, G.; Janssens, I.A.; Kleber, M.; Kögel-Knabner, I.; Lehmann, J.; Manning, D.A.C.; et al. Persistence of Soil Organic Matter as an Ecosystem Property. *Nature* **2011**, *478*, 49–56. [\[CrossRef\]](#)
49. Totsche, K.U.; Amelung, W.; Gerzabek, M.H.; Guggenberger, G.; Klumpp, E.; Knief, C.; Lehdorff, E.; Mikutta, R.; Peth, S.; Prechtel, A.; et al. Microaggregates in Soils. *J. Plant Nutr. Soil Sci.* **2017**, *181*, 104–136. [\[CrossRef\]](#)
50. Prata, F.; Lavoretti, A.; Regitano, J.B.; Tornisielo, V.L. Influência Da Matéria Orgânica Na Sorção e Dessorção Do Glifosato Em Solos Com Diferentes Atributos Mineralógicos. *Rev. Bras. Ci. Solo* **2000**, *24*, 947–951. [\[CrossRef\]](#)
51. Takeshita, V.; Mendes, K.F.; Alonso, F.G.; Tornisielo, V.L. Effect of Organic Matter on the Behavior and Control Effectiveness of Herbicides in Soil. *Planta Daninha* **2019**, *37*, 019214401. [\[CrossRef\]](#)
52. Mendes, K.F.; de Sousa, R.N.; Takeshita, V.; Alonso, F.G.; Régo, A.P.J.; Tornisielo, V.L. Cow Bone Char as a Sorbent to Increase Sorption and Decrease Mobility of Hexazinone, Metribuzin, and Quinclorac in Soil. *Geoderma* **2019**, *343*, 40–49. [\[CrossRef\]](#)
53. Prata, F.; Lavoretti, A.; Regitano, J.B.; Tornisielo, V.L. Degradação e Sorção de Ametrina Em Dois Solos Com Aplicação de Vinhaça. *Pesqui. Agropecuária Bras.* **2001**, *36*, 975–981. [\[CrossRef\]](#)
54. Li, X.G.; Jia, B.; Lv, J.; Ma, Q.; Kuzyakov, Y.; Li, F. Nitrogen Fertilization Decreases the Decomposition of Soil Organic Matter and Plant Residues in Planted Soils. *Soil Biol. Biochem.* **2017**, *112*, 47–55. [\[CrossRef\]](#)
55. Dupont, Y.L.; Strandberg, B.; Damgaard, C. Effects of Herbicide and Nitrogen Fertilizer on Non-Target Plant Reproduction and Indirect Effects on Pollination in *Tanacetum vulgare* (Asteraceae). *Agric. Ecosyst. Environ.* **2018**, *262*, 76–82. [\[CrossRef\]](#)
56. Langaro, A.C.; Agostinetto, D.; Oliveira, C.; Franco, J.J.; Zandoná, R.R.; Vargas, L. Influence of Nitrogen Fertilization on Herbicide Selectivity in Rice. *Planta Daninha* **2018**, *36*, 018180161. [\[CrossRef\]](#)
57. Hungria, M.; Mendes, I.C. Nitrogen Fixation with Soybean: The Perfect Symbiosis? In *Biological Nitrogen Fixation*; de Bruijn, F.J., Ed.; Wiley: Hoboken, NJ, USA, 2015; pp. 1009–1024. [\[CrossRef\]](#)
58. Rodrigues, T.F.; Bender, F.R.; Sanzovo, A.W.S.; Ferreira, E.; Nogueira, M.A.; Hungria, M. Impact of Pesticides in Properties of Bradyrhizobium Spp. and in the Symbiotic Performance with Soybean. *World J. Microbiol. Biotechnol.* **2020**, *36*, 172. [\[CrossRef\]](#)
59. Burul, F.; Barić, K.; Lakić, J.; Milanović-Litre, A. Herbicides Effects on Symbiotic Nitrogen-Fixing Bacteria. *J. Cent. Eur. Agric.* **2022**, *23*, 89–102. [\[CrossRef\]](#)
60. Aliverdi, A.; Ahmadvand, G. Herbicide Toxicity to Soybean–Rhizobium Symbiosis as Affected by Soil pH. *Bull. Environ. Contam. Toxicol.* **2018**, *101*, 434–438. [\[CrossRef\]](#)
61. Nunes, A.L.; Lorenset, J.; Gubiani, J.E.; Santos, F.M. A Multy-Year Study Reveals the Importance of Residual Herbicides on Weed Control in Glyphosate-Resistant Soybean. *Planta Daninha* **2018**, *36*, 018176135. [\[CrossRef\]](#)
62. Mehdizadeh, M.; Mushtaq, W.; Anusha Siddiqui, S.; Ayadi, S.; Kaur, P.; Yeboah, S.; Mazraedoost, S.; AL-Taey, D.K.A.; Tampubolon, K. Herbicide Residues in Agroecosystems: Fate, Detection, and Effect on Non-Target Plants. *Rev. Agric. Sci.* **2021**, *9*, 157–167. [\[CrossRef\]](#)
63. Madureira Barroso, G.; dos Santos, J.B.; de Oliveira, I.T.; Rocha Nunes, T.K.M.; Alves Ferreira, E.; Marinho Pereira, I.; Valadão Silva, D.; de Freitas Souza, M. Tolerance of *Bradyrhizobium* sp. BR 3901 to Herbicides and Their Ability to Use These Pesticides as a Nutritional Source. *Ecol. Indic.* **2020**, *119*, 106783. [\[CrossRef\]](#)
64. Vercellino, M.; Gómez, M.A. Denitrifying Capacity of Rhizobial Strains of Argentine Soils and Herbicide Sensitivity. *Ann. Microbiol.* **2013**, *63*, 1563–1570. [\[CrossRef\]](#)
65. Bossolani, J.W.; Poloni, N.M.; Lazarini, E.; Bettiol, J.V.T.; Fischer Filho, J.A.; Negrisoli, M.M. Development of RR Soybean in Function of Glyphosate Doses and Bradyrhizobium Inoculation. *Rev. Bras. Eng. Agrícola Ambient.* **2018**, *22*, 854–858. [\[CrossRef\]](#)
66. Khan, M.d.S.; Zaidi, A.; Aamil, M. Influence of Herbicides on Chickpea-Mesorhizobium Symbiosis. *Agronomie* **2004**, *24*, 123–127. [\[CrossRef\]](#)
67. Santos, J.B.; Silva, A.A.; Costa, M.D.; Jakelaitis, A.; Vivian, R.; Santos, E.A. Ação de Herbicidas Sobre o Crescimento de Estirpes de Rhizobium Tropici. *Planta Daninha* **2006**, *24*, 457–465. [\[CrossRef\]](#)
68. Melo, C.A.D.; Massenssini, A.M.; Passos, A.B.R.J.; Carvalho, F.P.; Ferreira, L.R.; Silva, A.A.; Costa, M.D. Isolation and Characteristics of Sulfentrazone-Degrading Bacteria. *J. Environ. Sci. Health Part B* **2016**, *52*, 115–121. [\[CrossRef\]](#)
69. Cao, B.; Zhang, Y.; Wang, Z.; Li, M.; Yang, F.; Jiang, D.; Jiang, Z. Insight into the Variation of Bacterial Structure in Atrazine-Contaminated Soil Regulating by Potential Phytoremediator: *Pennisetum americanum* (L.) K. Schum. *Front. Microbiol.* **2018**, *9*, 864. [\[CrossRef\]](#)

70. Mielke, K.C.; Bertuani, R.R.; Laube, A.F.S.; Brochado, M.G.d.S.; Ribeiro, A.C.M.; Paula, D.F.d.; Pires, F.R.; D'Angeri, R. Fitorremediação de Solos Contaminados Com Sulfentrazone Em Função Da Inoculação de Bactérias Simbióticas Em *Crotalaria juncea*. *Rev. Ibero-Am. Ciências Ambient.* **2021**, *12*, 140–151. [CrossRef]
71. Mielke, K.C.; Bertuani, R.R.; Pires, F.R.; Bueno Cotta, A.J.; Egreja Filho, F.B.; Madalão, J.C. Does *Canavalia ensiformis* Inoculation with *Bradyrhizobium* sp. Enhance Phytoremediation of Sulfentrazone-Contaminated Soil? *Chemosphere* **2020**, *255*, 127033. [CrossRef]
72. CAPES—Coordenação de Aperfeiçoamento de Pessoal de Nível Superior. Portal de Periódicos da Capes. Available online: <https://www-periodicos-capes-gov-br.ez1.periodicos.capes.gov.br/index.php> (accessed on 31 March 2023).
73. CLARIVATE Science Citation Index-Expanded. Available online: <https://clarivate.com/products/scientific-and-academic-research/research-discovery-and-workflow-solutions/web-of-science/web-of-science-core-collection/science-citation-index-expanded/#resources> (accessed on 23 March 2023).
74. ELSEVIER Science Direct. Available online: <https://www.sciencedirect.com/browse/journals-and-books> (accessed on 23 March 2023).
75. PMC National Library of Medicine. Available online: <https://www.ncbi.nlm.nih.gov/pmc/journals/#csvfile> (accessed on 23 March 2023).

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.