

Supplementary Material

Table S1. Demographic and clinical properties of patients with bilateral facial palsy/diplegia in complete Heerfordt's syndrome.

No	Age	Gend er	Heerfordt's syndrome manifestations/H.B. Scale	Other Systemic Manifestation	Diagnostic finding	Treatment	Prognosis	References
1	32 Y	Male	SBFP Fever Bilateral parotid swelling Bilateral anterior uveitis	Lung LNP+ (mediastinal,hilar)	-Serum ACE test ↑ -Parotid gland biopsy +	First-line therapy	CR	Srirangaramasamy et al. - 2016 [28]
2	42Y	Femal e	SBFD/- Fever Bilateral parotid swelling Unilateral anterior uveitis	Lung/Eye/Skin LNP+ (mediastinal,hilar)	- Skin biopsy +	First-line therapy	PR	King et al - 2007 [29]
3	26Y	Male	SBFD/- Fever Bilateral parotid swelling Unilateral anterior uveitis	EMG: proximal demyelination facial nerves.	-	Fist-line therapy Second-line therapy	CR	Glocker et al. - 1999 [30]
4	52Y	Femal e	SBFD/IV-IV/ Bilateral parotid swelling Unilateral anterior uveitis	Lung LNP+ (mediastinal,hilar)	-Serum ACE test ↑ - Parotid gland biopsy+	Fist-line therapy	PR (II-I)	Chappity et al - 2015 [8]
5	NA	NA	SBFD/- Fever Bilateral parotid swelling Anterior uveitis	-	Parotid gland FNA cytology+	-	-	
6	NA	NA	SBFD/- Fever Bilateral parotid swelling Anterior uveitis	-	Parotid gland FNA cytology+	-	-	
7	NA	NA	SBFD/- Fever Bilateral parotid swelling Anterior uveitis	-	Parotid gland FNA cytology+ -Bilateral Gasser's ganglia MRI impairment			Chappity et al - 2015 [8]
8	NA	NA	SBFD/- Bilateral parotid swelling Anterior uveitis	-	Parotid gland FNA cytology+	-	-	

CR: complete recovery; /-: unknown; HB Scale: House Brackmann Scale; FNA: fine-needle aspiration LNP+: lymphadenopathy; PPR: partial recovery; S: skin; SBFD: simultaneous bilateral facial diplegia; SBFP/RBFP: simultaneous/recurrent bilateral facial palsy