Effects of Geraniol on Survival, Reproduction, Endophytes, and Transcriptome of Tea Green Leafhoppers (*Empoasca onukii*)

Junjie Tian [1,2], Hainuo Hong [2], Shiliang Mo [3], Shiqin Mo [3], Hongliang Shi [1], Juan Shi [1,*],Ying Zhou [2,*]

[1] Beijing Key Laboratory for Forest Pest Control, Beijing Forestry University, Beijing100083, China; JunjieTian@bjfu.edu.cn (T.J.), shihl@bjfu.edu.cn (H.S.)

[2] Hainan Institute, Zhejiang University, Sanya 572025, China; 12216111@zju.edu.cn (H.H.)

[3] Jianfengling Branch Bureau of Hainan Tropical Rainforest National Park Service, Ledong572542, China; ShiliangMo1990@gmail.com (S.M.), ShiqinMo1989@gmail.com (S.M.)

[*] Correspondence: yzhyzb@zju.edu.cn (Y.Z.); bjshijuan@bjfu.edu.cn (J.S.)




Figure S1: Functional enrichment of tea green leafhopper genes compared with KEGG database. The numbers represent the number of genes annotated to this pathway, and the percentage values represent the proportion of the number of genes annotated to this pathway to the total number of annotated genes.