

Building Triazolated Macrocycles from Bis-propargylated Calix[4]arenes and Bis-azidomethylated Azobenzene or Stilbene

Ivan Lentin, Ilia Korniltsev, Alexander Gorbunov, Dmitry Cheshkov, Stanislav Bezzubov,
Vladimir Kovalev and Ivan Vatsouro

Supplementary Materials




Figure S1. ¹H NMR spectrum of compound 4 (400 MHz, CDCl₃).




Figure S2. ^{13}C NMR spectrum (APT) of compound 4 (100 MHz, CDCl_3).




Figure S3. ¹H NMR spectrum of compound 5 (400 MHz, CDCl₃+CD₃OD, 10:1).




Figure S4. ^{13}C NMR spectrum (APT) of compound 5 (100 MHz, $\text{CDCl}_3+\text{CD}_3\text{OD}$, 10:1).




Figure S5. ¹H NMR spectrum of compound c5 (400 MHz, CDCl₃).




Figure S6. ^{13}C NMR spectrum (APT) of compound **c5** (100 MHz, CDCl_3).




Figure S7. ¹H NMR spectrum of compound 6 (400 MHz, CDCl₃+CD₃OD, 10:1).




Figure S8. ^{13}C NMR spectrum (APT) of compound **6** (100 MHz, $\text{CDCl}_3 + \text{CD}_3\text{OD}$, 10:1).




Figure S9. ¹H NMR spectrum of compound **c6** (400 MHz, CDCl₃).




Figure S10. ^{13}C NMR spectrum (APT) of compound **c6** (100 MHz, CDCl_3).




Figure S11. ¹H NMR spectrum of compound 8 (400 MHz, CDCl₃).




Figure S12. ^{13}C NMR spectrum (APT) of compound 8 (100 MHz, CDCl_3).