

Correction

# Correction: Otón, G., et al. Global Detection of Long-Term (1982–2017) Burned Area with AVHRR-LTDR Data. *Remote Sensing* 2019, 11, 2079

Gonzalo Otón , Rubén Ramo , Joshua Lizundia-Loiola  and Emilio Chuvieco 

Department of Geology, Geography and the Environment, University of Alcalá, 28801 Alcalá de Henares, Spain; ruben.ramo@uah.es (R.R.); joshua.lizundia@uah.es (J.L.-L.); emilio.chuvieco@uah.es (E.C.)

\* Correspondence: gonzalo.oton@uah.es

Received: 10 July 2020; Accepted: 15 July 2020; Published: 20 July 2020



The authors wish to make the following corrections to this paper [1], which includes an error in the following formula (Section 2.5 of the paper) relate.

It currently indicates:

$$LBI = z(T5) - z(T5\_diff) - z(Red) - z(Red\_diff) - z(NIR) - z(NIR\_diff) - z(GEMI) - z(BAI) - z(BAI_{t+1})$$

The correct formula of the LBI is:

$$LBI = z(T5) - z(T5\_diff) - z(Red) + z(Red\_diff) - z(NIR) + z(NIR\_diff) + z(GEMI) + z(BAI) + z(BAI_{t+1})$$

Please, note that signs are not negative for some of the input variables.

The authors would like to apologize for any inconvenience caused to the readers by the original typo.

## References

1. Otón, G.; Ramo, R.; Lizundia-Loiola, J.; Chuvieco, E. Global Detection of Long-Term (1982–2017) Burned Area with AVHRR-LTDR Data. *Remote Sens.* **2019**, *11*, 2079. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).