

Supporting Information for

Emerging Signal of Englacial Debris on One Clean Surface Glacier Based on High Spatial Resolution Remote Sensing Data in Northeastern Tibetan Plateau

Yuwei Wu ¹, Wei Zhang ^{2,*}, Shiqiang Zhang ¹, Zhonglei Mao ¹, Anan Chen ¹, Zhen Li ²,
Quan Zhang ¹, Zhongming Guo ¹, Xi Jiang ³ and Yongqing Long ¹

¹ Shaanxi Key Laboratory of Earth Surface System and Environmental Carrying Capacity,

College of Urban and Environmental Science, Northwest University, Xi'an 710127, China;

yuwei.wu@nwu.edu.cn (Y.W.); zhangsq@nwu.edu.cn (S.Z.);

maozl@stumail.nwu.edu.cn (Z.M.); aachen@nwu.edu.cn (A.C.);

zhangquanzq@nwu.edu.cn (Q.Z.); gzm@nwu.edu.cn (Z.G.); sjzxlyq@nwu.edu.cn (Y.L.)

² State Key Laboratory of Cryospheric Sciences, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China; lizhen@lzb.ac.cn

³ Center for Data Assimilation Research and Application, School of Atmospheric Sciences, Nanjing University of Information Science & Technology, Nanjing 210044, China; jiangxi@nuist.edu.cn

* Correspondence: zhangw06@lzb.ac.cn

Contents of this file

Figures S1 to S2



Figure S1. Exposed englacial debris in five zones A-E (a-e, respectively; taken on summer 2022).




Figure S2. Ice disintegration in the terminus of the Qiyi Glacier that was first observed on summer 2013. a: ice crevasse; b: ice disintegration from the ice crevasse.