

Electronic Supplementary Information for

Adipic acid route: oxidation of cyclohexene *vs.* cyclohexane

Ana P. C. Ribeiro ¹, Elisa Spada, Roberta Bertani ² and Luísa M. D. R. S. Martins ^{1,*}

¹ Centro de Química Estrutural and Departamento de Engenharia Química, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal; e-mail: apribeiro@tecnico.ulisboa.pt

² Department of Industrial Engineering, University of Padova, Padova, Italy; elisaspada91@hotmail.it; roberta.bertani@unipd.it

* Correspondence: luisammartins@tecnico.ulisboa.pt Tel.: +351-218419389

CONTENTS

	Page
¹ H NMR spectrum of isolated adipic acid (Figure S1).	2
Far infrared spectrum of [FeCl ₂ {κ ³ -HC(pz) ₃ }] after the oxidation reaction (Figure S2).	2




Figure S1 - ^1H NMR spectrum of isolated adipic acid in DMSO.




Figure S2 - Far infrared spectrum of $[\text{FeCl}_2\{\kappa^3\text{-HC}(\text{pz})_3\}]$ after acting as catalyst for the MW-assisted oxidation reaction of cyclohexene with H_2O_2 .