

Article

Long-Term Monitoring of a Surface Flow Constructed Wetland Treating Agricultural Drainage Water in Northern Italy

Stevo Lavrnić ¹, Ilaria Braschi ¹, Stefano Anconelli ², Sonia Blasioli ¹, Domenico Solimando ², Paolo Mannini ² and Attilio Toscano ^{1,*}

¹ Department of Agricultural and Food Sciences, Alma Mater Studiorum -University of Bologna, Viale Giuseppe Fanin 40-50, 40127 Bologna, Italy; stevo.lavrnice@unibo.it (S.L.); ilaria.braschi@unibo.it (I.B.); sonia.blasioli@unibo.it (S.B.)

² Consorzio di Bonifica Canale Emiliano Romagnolo, Via Ernesto Masi 8, 40137 Bologna, Italy; anconelli@consorzioicer.it (S.A.); solimando@consorzioicer.it (D.S.); mannini@consorzioicer.it (P.M.)

* Correspondence: attilio.toscano@unibo.it; Tel.: +39-051-20-9-6179

Supporting information

Table SI.1. Farm TN input and SFCW nitrogen balance (all the values are in kg).

	Farm		SFCW		
	Fertilisation	Rain	Inflow	Rain	Outflow
2003–2004	1518	60	311	2	125
2004–2005	1515	226	210	6	115
2005–2006	1750	162	215	4	39
2006–2007	1834	178	1	5	0
2007–2008	1319	330	3	9	0
2008–2009	2531	366	170	10	11