

Article

A Facile Synthesis of α -*N*-Ribosyl-Asparagine and α -*N*-Ribosyl-Glutamine Building Blocks

Gaetano Speciale, Anna Bernardi and Filippo Nisic *

Universita' degli Studi di Milano, Dipartimento di Chimica, via Golgi 19, 20133 Milano, Italy;
E-Mails: g.speciale@student.unimelb.edu.au (G.S.); anna.bernardi@unimi.it (A.B.)

* Author to whom correspondence should be addressed; E-Mail: filippo.nisic@unimi.it;
Tel.: +39-02-5031-4092, Fax: +39-02-5031-4072.

Received: 2 July 2013; in revised form: 16 July 2013 / Accepted: 19 July 2013 /

Published: 24 July 2013

Abstract: Adenosine diphosphate ribosylation (ADP-ribosylation) is a widely occurring post-translational modification of proteins at nucleophilic side chain of amino acid residues. Elucidation of ADP-ribosylation events would benefit greatly from the availability of well-defined ADP-ribosylated peptides and analogues thereof. In this paper we present a novel approach to the chemical synthesis of ribosylated amino acid building blocks using traceless Staudinger ligation. We describe an efficient and stereoselective synthesis of α -*N*-ribosyl-asparagine (α -*N*-ribosyl-Asn) and α -*N*-ribosyl-glutamine (α -*N*-ribosyl-Gln) building blocks starting from 5-*tert*-butyldiphenylsilyl- β -D-ribofuranosyl azide. The *N*-glycosyl aminoacids are produced in good yields as pure α -anomers, suitably protected for peptide synthesis.


Keywords: ADP-ribosylation; glycoconjugates; ribofuranosyl aminoacids; Staudinger ligation; stereoselective synthesis

1. Introduction

Glycosylated peptides play a fundamental role in biological systems: in fact, more than half of all proteins carry carbohydrate moieties, generating different glycoforms whose exact composition often controls protein function and distribution in biological systems [1–4]. The details of glycan regulation of protein activity and stability are still under intense scrutiny and the synthesis of well-defined glycopeptides is therefore an important target [5,6], which still presents many challenges to organic


chemistry. One of the processes that is currently being elucidated is adenosine diphosphate ribosylation (ADP-ribosylation), a wide-occurring post-translational modification effected by enzymes that transfer ADP-ribose from NAD^+ to Asn, Glu, Asp, Arg or Cys residues of proteins, so altering their function [7]. An important contribution for the clarification of the role of ADP-ribosylation events came in 2010, when Filippov and coworkers [8] reported the synthesis of ribosylated oligopeptides using α -*N*-ribosyl-asparagine (α -*N*-ribosyl-Asn) and α -*N*-ribosyl-glutamine (α -*N*-ribosyl-Gln) building blocks. Despite the relevance of this work, the approach suffers from a poorly selective synthesis of the required ribosylated building blocks **4** α and **5** α (Scheme 1), which were prepared by PtO_2 reduction of azide **3**, followed by EDC-mediated coupling of the resulting mixture of epimeric amines with *Z*-Asp-OBn or *Z*-Glu-OBn, respectively. The *N*-ribosyl-aminoacids were obtained as a 3:1 α : β anomeric mixtures, from which the desired α -anomers were to be chromatographically isolated. Acetyl transfer to the anomeric nitrogen from the 2-*O*-acetate also occurs during the reaction, further reducing the process yields, particularly for the asparagine derivative **4** α .

Scheme 1. Filippov's synthesis of ribosylated amino acids (from ref. [8]).



We have recently developed a protocol for the stereoselective synthesis of either α - or β -*N*-glycofuranosyl amides using the Staudinger traceless ligation [9,10]. In this reaction, a starting anomeric glycosyl azide is partially reduced by an appropriate phosphine and then intramolecularly acylated in order to afford the amide (Scheme 2) [11]. We have shown that for furanosyl azides the outcome of the process is controlled by the configuration and the protection state of the contiguous hydroxyl group, so that 1,2-*cis* amides (α -, in the *ribo* series) are obtained when this group is unprotected and 1,2-*trans* amides (β -, in the *ribo* series) when it is acetylated. For unprotected furanoses, the anomeric configuration of the product amides appears to be dictated by O-P coordination, as supported by ^{31}P -NMR studies [10].

Scheme 2. Traceless Staudinger ligation of furanosyl azides with functionalized phosphines (from ref [9]).




As an application of this methodology, we here report the facile stereoselective synthesis of **4a** and **5a** that could be obtained in good yields and with full stereoselectivity with a single synthetic operation starting from β -5-*O*-*tert*-butyldiphenylsilyl-ribosylazide **2**.

2. Results and Discussion


Previous research in our laboratories had shown that transfer of aminoacid chains in Staudinger ligations of glycosyl azides was particularly effective using fluorinated phosphines **6** [12]. Thus we envisaged that phosphines **6a** and **6b** (Scheme 3), functionalized with *Z*-Asp-OBn and *Z*-Glu-OBn acyl chains, respectively, would be the reagent of choice to prepare the target α -*N*-ribosyl-Asn and α -*N*-ribosyl-Gln building blocks. Phosphines **6a** and **6b** were synthesized in excellent yields by EDC-mediated acylation of 2-diphenylphosphanyl-5-fluorophenol (**7**) [13] and purified by filtration on short pads of silica.

Scheme 3. Synthesis of functionalized phosphines **6a** and **6b**.



An early trial of ligation was conducted using the unprotected β -D-ribofuranosyl azide **1** [14] with phosphines **6a** and **6b** for 20 h at 70 °C in a 98:2 DMA:DMPU [15] solution, followed by water quenching (Scheme 4). In both ligation reactions 45% of a 3:2 mixture of α -ribofuranosyl amide (**8a** or **9a**) and β -ribofuranosyl amide (**10** or **11**) was obtained. The isomer distribution was determined by $^1\text{H-NMR}$ analysis of the crude reaction mixtures: the α -ribofuranosyl amide (**8a** or **9a**) was characterized by the presence of the H1 proton signal at 5.65 ppm and by the high chemical shift of C4 (84.3 ppm), which is diagnostic for the furanose form. The structures of **10** and **11** were assigned on the basis of the vicinal coupling constant of the anomeric protons ($J_{1,2} = 8$ Hz, H1 = 5.12 ppm) and of the low chemical shift value of C4 (68.8 ppm), that confirms the pyranose structure. The β -glycopyranosyl isomers **10** and **11** obtained in these reactions must derive from ring-expansion occurring after a ring-opening process.


Scheme 4. Ligation of **1** with phosphines **6a** and **6b**.



Reagents and Conditions: DMA/DMPU 98:2, 70 °C, 20 h; (b) H_2O , 2 h, 70 °C.

To avoid ring expansion, the 5-hydroxy group of **1** was protected as a *tert*-butyldiphenylsilyl ether and the 5-*tert*-butyldiphenylsilyl- β -D-ribofuranosyl azide **2** was synthesized in good yields as described by Filippov [8]. Ligation of **2** with **6a** or **6b** was carried out as above (20 h at 70 °C in a 98:2 DMA:DMPU mixture) (Scheme 5). In both cases, 400 MHz ^1H -NMR analysis of the crude showed only one signal in the furanose anomeric region, implying that no pyranose was formed, as expected, and that the α/β ratio of the resulting ribofuranosyl amides **12a** and **13a** was $\geq 99:1$. The two compounds were isolated by flash chromatography, which afforded the desired building blocks in 60% and 76% yields, respectively, as single α isomers. After acetylation (Ac_2O , cat. DMAP, CH_2Cl_2 , quant.), **4a** and **5a** were obtained. Their spectroscopic data were fully consistent with the already published characterization [8].

Scheme 5. Synthesis of ribofuranosyl amides **4a** and **5a**.



Reagents and Conditions: DMA/DMPU 98:2, 70 °C, 20 h; (b) H_2O , 2 h, 70 °C; (c) Ac_2O , cat. DMAP, CH_2Cl_2 .

3. Experimental

Solvents were dried by standard procedures: dichloromethane and *N,N*-diisopropylethylamine were dried over calcium hydride; *N,N*-dimethylacetamide (DMA), 1,3-dimethyltetrahydro-2(1*H*)-pyrimidinone (DMPU), chloroform and pyridine were dried over activated molecular sieves. Reactions requiring anhydrous conditions were performed under nitrogen. ^1H -, ^{13}C - and ^{31}P -NMR spectra were recorded at 400 MHz on a Bruker AVANCE-400 instrument. Chemical shifts (δ) for ^1H and ^{13}C spectra are expressed in ppm relative to internal Me_4Si as standard. Chemical shifts (δ) for ^{31}P are expressed in ppm relative to internal H_3PO_4 as standard. Signals were abbreviated as s, singlet; bs, broad singlet; d, doublet; t, triplet; q, quartet; m, multiplet. Mass spectra were obtained with a Bruker ion-trap Esquire 3000 apparatus (ESI ionization) or FT-ICR Mass Spectrometer APEX II & Xmass software (Bruker Daltonics)—4.7 Magnet and Autospec Fission Spectrometer (FAB ionization). Thin layer chromatography (TLC) was carried out with pre-coated Merck F_{254} silica gel plates. Flash chromatography (FC) was carried out with Macherey-Nagel silica gel 60 (230–400 mesh). The typical scale used for the ligation reactions in this paper was 20 mg (0.1 mmol) of unprotected glycosyl azide.

General Procedure for the Synthesis of 2-(diphenylphosphanyl)-4-fluorophenyl Esters 6

A solution of the *o*-diphenylphosphinophenol (**7**, 1 equiv.), the commercially available *N*-carbobenzyloxy-*L*-aminoacid-1-benzyl ester (*Z*-Asp-OBn or *Z*-Glu-OBn, 1.2 equiv.) and *N,N*-dimethylaminopyridine (0.1 equiv.) in dry CH_2Cl_2 (0.1 M) were added, at room temperature and under nitrogen, to a suspension of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDC, 1.4 equiv.) and dry *N,N*-diisopropylethylamine (1.4 equiv.) in dry CH_2Cl_2 . The mixture was

stirred at RT for 2 h, monitoring by TLC (60:40 hexane/AcOEt). The reaction mixture was diluted with CH_2Cl_2 and washed with 5% aqueous HCl and water: the organic layer was dried over Na_2SO_4 and concentrated. The crude product obtained was purified by flash chromatography, as indicated in each case.

1-Benzyl 5-[2-(Diphenylphosphanyl)-4-fluorophenyl] N-(Benzyloxycarbonyl)-L-aspartate (6a). The crude product was purified by flash chromatography (hexane/AcOEt 60:40). yield = 85%. $^1\text{H-NMR}$ (CDCl_3 , 25 °C): δ = 7.37–7.23 (m, 20H, Ph), 7.02–6.84 (m, 2H, H-2, H-3), 6.49 (m, 1H, H-1), 5.69 (d, $J_{\text{NH-CH}}$ = 8.6 Hz, 1H, NH), 5.17 (s, 2H, CH_2O), 5.13 (s, 2H, CH_2O), 4.62 (m, $J_{\text{CH-CH}_2}$ = 4.4 Hz, 1H, CH), 3.00 (dt, J = 4.9 Hz, J = 17.4 Hz, 1H, H_a , CH_2), 2.67 (dd, J = 4.3 Hz, J = 17.4 Hz, 1H, H_b , CH_2). $^{13}\text{C-NMR}$ (CDCl_3 , 25 °C): δ = 170.4, 169.2, 161.9, 159.4 ($\text{CO}_{(\text{Cbz})}$), 134.6, 134.0, 132.0, 131.7, 129.0, 128.9, 128.7, 128.6, 128.4, 128.3, 128.2 (C_{Ar}), 124.0, 116.9, 116.6, 67.8 (CH_2O), 67.3 (CH_2O), 50.4 (CH), 36.5 (CH_2). $^{31}\text{P-NMR}$ (161 MHz, CDCl_3 , 25 °C): δ = -14.3 ppm. FAB-MS: m/z 636 (M+1).

1-Benzyl 5-[2-(Diphenylphosphanyl)-4-fluorophenyl] N-(Benzyloxycarbonyl)-L-glutamate (6b). The crude product was purified by flash chromatography (hexane/AcOEt 70:30). yield = 85%. $^1\text{H-NMR}$ (CDCl_3 , 25 °C): δ = 7.37–7.24 (m, 20 H, Ph), 7.05 (m, 1H, H-2), 6.99 (m, 1H, H-3), 6.45 (m, 1H, H-1), 5.37 (d, $J_{\text{NH-CH}}$ = 8.1 Hz, 1H, NH), 5.16 (s, 2H, $\text{CH}_2\text{-O}$), 5.07 (s, 2H, $\text{CH}_2\text{-O}$), 4.38 (m, 1H, CH), 2.26 (m, 2H, $\text{CH}_2\text{-CO}$), 2.05 (m, 1H, H_a , $\text{CH}_2\text{-CH}$), 1.80 (m, 1H, H_b , $\text{CH}_2\text{-CH}$). $^{13}\text{C-NMR}$ (CDCl_3 , 25 °C): δ = 170.7, 161.7, 159.3, 156.1 ($\text{CO}_{(\text{Cbz})}$), 134.4, 134.3, 134.2, 134.0, 129.1, 129.0, 128.8, 128.5, 128.4 (C_{Ar}), 124.2, 124.1, 120.0, 119.8, 116.8, 116.5, 67.6 ($\text{CH}_2\text{-O}$), 67.4 ($\text{CH}_2\text{-O}$), 53.5 (CH), 30.1 ($\text{CH}_2\text{-CO}$), 27.4 (CH_2). $^{31}\text{P-NMR}$ (161 MHz, CDCl_3 , 25 °C): δ = -14.3 ppm. FAB-MS: m/z 650 (M+1).

General Procedure for the Staudinger Traceless Ligation of Ribofuranosyl Azides

Phosphine **6a** or **6b** (2 equiv.) was added, at room temperature, to a 0.1 M solution of ribofuranosyl azide **2** (1 equiv.) in 98: 2 *N,N*-dimethylacetamide and DMPU. The solution was stirred for 20 h at 70 °C, then 10% water was added and the mixture was stirred for an additional 2 h at the same temperature. The solvent was evaporated under reduced pressure, and the residue was purified by flash chromatography as indicated in each case.

General Procedure for the Acetylation of the Ribofuranosyl Amides

Ac_2O (6 equiv.), pyridine (6 equiv.) and a catalytic amount of *N,N*-dimethylaminopyridine were added, at room temperature, to a solution of substrate (1 equiv.) in dry CH_2Cl_2 (0.1 M). The solution was stirred for 24 h and then was concentrated in vacuo. The residue was dissolved in AcOEt and washed with aqueous 5% HCl, aqueous 5% NaHCO_3 and water. The organic layer was dried over Na_2SO_4 and concentrated. The crude was then purified by flash chromatography (hexane:AcOEt, 80:20 to 50:50). Purification afforded products with spectroscopic data coherent with the fully characterized compounds published by Filippov and coworkers [8].

N^δ-(5-tert-Butyldiphenylsilyl- α -D-ribose)-N ^{α} -benzyloxycarbonyl asparagine benzyl ester (12a). The crude product was purified by flash chromatography ($\text{CHCl}_3/\text{MeOH}$ 95:5). yield = 60%. $^1\text{H-NMR}$ (CD_3OD , 25 °C): δ = 7.71–7.64, 7.42–7.24 (3 \times m, 20H, Arom.), 5.73 (d, $J_{1,2}$ = 4.8 Hz, 1H, H-1), 5.15, 5.07 (2 \times s, 2H, CH_2Bn , CH_2Cbz), 4.65 (m, 1H, CH α -Asn), 4.25 (t, J = 5 Hz, 1H, H-3), 4.15 (t, J = 4.8 Hz,

1H, H-2), 3.97 (m, 1H, H-4), 3.72 (dd, $J_{4,5} = 3.2$ Hz, $J_{gem} = 11.3$ Hz, 2H, H-5 H-5'), 2.84 (m, 2H, CH₂ β-Asn), 1.04 (s, 9H, t-Bu). ¹³C-NMR (CD₃OD, 25 °C): 172.9, 172.4 (CO α-Asn, CO γ-Asn), 158.6 (CO Cbz), 138.2 (Cq Arom.), 137.3 (Cq Arom.), 136.9, 136.8 (Arom.), 134.6, 134.4 (Cq Arom.), 131.1, 131.0, 129.7, 129.6, 129.5, 129.5, 129.3, 129.3, 129.0, 128.8, 128.4, 128.1 (Arom.), 84.7 (C-4), 81.9 (C-1), 72.8 (C-3), 72.1 (C-2), 68.4, 67.9 (CH₂ Bn, CH₂ Cbz), 65.3 (C-5), 52.4 (CH, α-Asn), 38.9 (CH₂, β-Asn), 27.4 (CH₃, t-Bu), 20.2 (Cq, t-Bu). FT-ICR (ESI) calcd. for C₄₀H₄₆N₂O₉Si [M+Na]⁺ 749.29726; found 749.29721.

N^δ-(5-*tert*-Butyldiphenylsilyl-α-*D*-ribose)-*N*^α-benzyloxycarbonyl glutamine benzyl ester (**13α**). The crude product was purified by flash chromatography (CHCl₃/MeOH 95:5). yield = 76%. ¹H-NMR (CD₃OD, 25 °C): δ = 7.71–7.64, 7.43–7.23 (3 x m, 20H, Arom.), 5.72 (d, $J_{1,2} = 4.6$ Hz, 1H, H-1), 5.15, 5.06 (2 × s, 2H, CH₂Bn, CH₂Cbz), 4.25 (m, 1H, CH α-Gln), 4.23 (t, $J = 4.9$ Hz, 1H, H-3), 4.15 (t, $J = 4.8$ Hz, 1H, H-2), 3.95 (m, 1H, H-4), 3.70 (dd, $J_{4,5} = 3.2$ Hz, $J_{gem} = 10.2$ Hz, 2H, H-5 H-5'), 2.35 (m, 2H, CH₂ γ-Asn), 2.13 (m, 1H, CH₂ β-Asn), 1.95 (m, 1H, CH₂ β-Asn), 1.02 (s, 9H, t-Bu). ¹³C-NMR (CD₃OD, 25 °C): 136.7, 136.6 (Arom.), 134.6, 134.4 (Cq Arom.), 130.8, 130.8, 129.7, 129.5, 129.4, 129.1, 128.9, 128.8, 128.7 (Arom.), 84.5 (C-4), 81.6 (C-1), 72.7 (C-3), 71.9 (C-2), 67.9, 67.6 (CH₂ Bn, CH₂ Cbz), 65.0 (C-5), 55.1 (CH α-Gln), 33.3 (CH₂ β-Gln), 28.0 (CH₂ γ-Gln), 27.3 (t-Bu), 22.9 (Cq t-Bu). FT-ICR (ESI) calcd. for C₄₁H₄₈N₂O₉Si [M+Na]⁺ 763.31291; found 763.31298.

4. Conclusions

In conclusion, we have described an improved synthesis of the building blocks **4α** and **5α**: application of the Staudinger Traceless Ligation protocol allowed to obtain the ribosylated amino acids from **2** in good yields and with excellent selectivity for the α anomer. This represents a major advantage over pre-existing synthesis of these glycosyl aminoacids [8,16,17] and should stimulate further research in the field of ADP-ribosylation.

Acknowledgments

This work was supported by the University of Milan (PUR) and Ministero dell'Università e della Ricerca (PRIN). Exact mass were obtained from CIGA.

Conflict of Interest

The authors declare no conflict of interest.

References and Notes

1. Varki, A. Biological roles of oligosaccharides: all of the theories are correct. *Glycobiology* **1993**, *3*, 97–130.
2. Dwek, R.A. Glycobiology: Toward understanding the function of sugars. *Chem. Rev.* **1996**, *96*, 683–720.
3. Reuter, G.; Gabius, H.J. Eukaryotic glycosylation: whim of nature or multipurpose tool? *Cell. Mol. Life Sci.* **1999**, *55*, 368–422.

4. Gamblin, D.P.; Scanlan, E.M.; David, B.G. Glycoprotein synthesis: an update. *Chem. Rev.* **2009**, *109*, 131–163.
5. Marcaurelle, L.A.; Bertozzi, C.R. Recent advances in the chemical synthesis of mucin-like glycoproteins. *Glycobiology* **2002**, *12*, 69R–77R.
6. Warren, J.D.; Miller, J.S.; Keding, S.J.; Danishefsky, S.J. Toward fully synthetic glycoproteins by ultimately convergent routes: A solution to a long-standing problem. *J. Am. Chem. Soc.* **2004**, *126*, 6576–6578.
7. Corda, D.; Di Girolamo, M. Functional aspects of protein mono-ADP-ribosylation. *EMBO J.* **2003**, *22*, 1953–1958.
8. van der Heden van Noort, G.J.; van der Horst, M.G.; Overkleeft, H.S.; van der Marel, G.A.; Filippov, D.V. Synthesis of Mono-ADP-Ribosylated oligopeptides using ribosylated amino acid building blocks. *J. Am. Chem. Soc.* **2010**, *132*, 5236–5240.
9. Nisic, F.; Bernardi, A. Stereoselective synthesis of *N*-galactofuranosyl amides *Carbohydr. Res.* **2011**, *346*, 465–471.
10. Nisic, F.; Speciale, G.; Bernardi, A. Stereoselective synthesis of α - and β -Glycofuranosyl amides by traceless ligation of glycofuranosyl azides. *Chem. Eur. J.* **2012**, *18*, 6895–6906.
11. Bianchi, A.; Bernardi, A. Traceless Staudinger ligation of glycosyl azides with triaryl phosphines: Stereoselective synthesis of glycosyl amides. *J. Org. Chem.* **2006**, *71*, 4565–4577.
12. Nisic, F.; Andreini, M.; Bernardi, A. Stereoselective synthesis of *N*-Glycosyl amino acids by traceless Staudinger ligation of unprotected glycosyl azides. *Eur. J. Org. Chem.* **2009**, *33*, 5744–5751.
13. Heinicke, J.; Köhler, M.; Peulecke, N.; He, M.; Kindermann, M.K.; Keim, W.; Fink, G. 2-phosphanilphenolate nickel catalysts for the polymerization of ethylene. *Chem. Eur. J.* **2003**, *9*, 6093–6107.
14. Stimac, A.; Kobe, J. An improved preparation of 2,3,5-tri-O-acyl- β -D-ribofuranosyl azides by the Lewis acid-catalysed reaction of β -D-ribofuranosyl acetates and trimethylsilyl azide: an example of concomitant formation of the α anomer by trimethylsilyl triflate catalysis. *Carbohydr. Res.* **1992**, *232*, 359–365.
15. This solvent mixture was found to perform optimally in Staudinger ligations of unprotected glycosyl azides, see: Nisic, F.; Bernardi, A. Stereoselective synthesis of glycosyl amides by traceless Staudinger ligation of unprotected glycosyl azides. *Carb. Res.* **2008**, *343*, 1636–1643.
16. Bonache, M.A.; Nuti, F.; Le Chevalier Isaad A.; Real-Fernandez F.; Chelli M.; Rovero, P.; Papini A.M. Synthesis of new ribosylated Asn building blocks as useful tools for glycopeptide and glycoprotein synthesis. *Tetrahedron Lett.* **2009**, *50*, 4151–4153.
17. After the ECSOC conference, while this paper was under review, a novel rybosylation of Asn and Gln has appeared in the literature: Kistemaker, H.A.V.; van der Heden van Noort, G.J.; Overkleeft, H.S.; van der Marel, G.A.; Filippov, D.V. Stereoselective Ribosylation of Amino Acids. *Org. Lett.* **2013**, *15*, 2306–2308.

Sample Availability: Samples of the compounds **1**, **6** and **7** are available from the authors.