Supplementary Information

Figures S1–S6 The ¹H-, ¹³C-NMR, HSQC, HMBC, COSY and ROESY of botryosphaerin G in CHCl₃.

Figures S7–S12 The ¹H-, ¹³C-NMR, HSQC, HMBC, COSY and ROESY of botryosphaerin H in C₅D₅N.




Figure S1. The ¹H-NMR of botryosphaerin G in CHCl₃.




Figure S2. The ¹³C-NMR of botryosphaerin G in CHCl₃.




Figure S3. The HSQC NMR of botryosphaerin G in CHCl₃.




Figure S4. The HMBC NMR of botryosphaerin G in CHCl₃.




Figure S5. The COSY NMR of botryosphaerin G in CHCl₃.




Figure S6. The ROESY NMR of botryosphaerin G in CHCl₃.




Figure S7. The ¹H-NMR of botryosphaerin H in C₅D₅N.




Figure S8. The ¹³C-NMR of botryosphaerin H in C₅D₅N.




Figure S9. The HSQC NMR of botryosphaerin H in C₅D₅N.




Figure S10. The HMBC NMR of botryosphaerin H in C₅D₅N.




Figure S11. The COSY NMR of botryosphaerin H in C₅D₅N.




Figure S12. The ROESY NMR of botryosphaerin H in C₅D₅N.