## Supplementary Materials: Synthesis, Characterization and Molecular Docking of Novel Bioactive Thiazolyl-Thiazole Derivatives as promising Cytotoxic Antitumor Drug

oc decoupled spectra MOSTAFA A3 in DMSO BŘ ÉR 40.51 40.39 40.25 39.97 39.83 39.69 .69 A 20130510 19.36 spect INSTRUM PROBHD PULPROG CHa PU TD SO NS DS LVENT SWH FIDRES AQ RG DW DE TE VH-D1 D11 TD0 14 a SFO1 NUC1 MHZ P1 PLW1 8.80 uses 78.13500214 W ANNEL f2 ======= 600.1324005 MHz 1H waltz65 70.00 use NUC2 CPDPF PCPD2 PLW2 PLW12 PLW13 0.63804001 W 0.31264001 W F2 - Pro SI SF WDW -SSB LB GB GB -PC ssing parameters 32768 150.9028090 MHz EM 1.00 Hz 1.40 0 200 180 160 140 120 100 80 60 40 20 ppm

Sobhi M. Gomha, Taher A. Salah, Huwaida M. E. Hassaneen and Mohammed A. Khedr

Figure S1. <sup>13</sup>C-NMR of compound 14a.




Figure S2. <sup>1</sup>H-NMR of compound 14a.




Figure S3. Mass of compound 14a.




Figure S5.<sup>13</sup>C-NMR of compound 14b.




Figure S6. <sup>1</sup>H-NMR of compound 14b.




Figure S7. Mass of compound 14b.




Figure S9. <sup>13</sup>C-NMR of compound 14c.




Figure S10. <sup>1</sup>H-NMR of compound 14c.




Figure S11. Mass of compound 14c.




Figure S12. <sup>13</sup>C-NMR of compound 14d.




Figure S13. Mass of compound 14d.




Figure S14. <sup>13</sup>C-NMR of compound 14e.




Figure S15. <sup>1</sup>H-NMR of compound 14e.




Figure S16. Mass of compound 14e.




Figure S17. <sup>13</sup>C-NMR of compound 17.




Figure S18. <sup>1</sup>H-NMR of compound 14e.




Figure S19. Mass of compound 17.




Figure S20. Mass of compound 3a.




Figure S21. Mass of compound 6b




Figure S22. Mass of compound 6c.