

Supplementary Materials: New Sesquiterpenoids and Anti-Platelet Aggregation Constituents from the Rhizomes of *Curcuma zedoaria*

Jih-Jung Chen, Tung-Han Tsai, Hsiang-Ruei Liao, Li-Chai Chen, Yueh-Hsiung Kuo, Ping-Jyun Sung, Chun-Lin Chen and Chun-Sheng Wei




Figure S1. ESI-MS spectrum of 1.






Figure S2. HR-ESI-MS spectrum of 1.

Figure S3. ¹H-NMR spectrum of 1 (CDCl₃, 500 MHz).Figure S4. ¹³C-NMR spectrum of 1 (CDCl₃, 125 MHz).




Figure S5. ^1H - ^1H COSY spectrum of 1.






Figure S6. NOESY spectrum of 1.

Figure S7. HMBC spectrum of **1**.Figure S8. HSQC spectrum of **1**.




Figure S9. ESI-MS spectrum of 2.




Figure S10. HR-ESI-MS spectrum of 2.




Figure S11. ¹H-NMR spectrum of 2 (CDCl₃, 500 MHz).




Figure S12. ¹³C-NMR spectrum of 2 (CDCl₃, 125 MHz).




Figure S13. ^1H - ^1H COSY spectrum of 2.




Figure S14. NOESY spectrum of 2.




Figure S15. HMBC spectrum of 2.




Figure S16. HSQC spectrum of 2.