

Article

Bioassay-Guided Isolation of Cytotoxic Isocryptoporic Acids from *Cryptoporus volvatus*

Ling-Yun Zhou ^{1,2}, Xiao-Hong Yu ¹, Bin Lu ¹ and Yan Hua ^{1,*}

¹ Key Laboratory for Forest Resources Conservation and Use in the Southwest Mountains of China, Southwest Forestry University, Kunming 650224, China; zly4321@sina.com (L.-Y.Z.); 13187815521@163.com (X.-H.Y.); nongxue022@163.com (B.L.)

² School of Pharmacy, Wannan Medical College, Wuhu 241002, China

* Correspondence: huayan1216@163.com; Tel.: +86-871-6386-3145

Academic Editor: Thomas J. Schmidt

Received: 19 October 2016; Accepted: 6 December 2016; Published: 8 December 2016

Abstract: The present work constitutes a contribution to the phytochemical investigation of *Cryptoporus volvatus* aiming to search for effective cytotoxic constituents against tumor cell lines in vivo. Bioassay-guided separation of the ethylacetate extract of *C. volvatus* afforded four new isocryptoporic acid (ICA) derivatives, ICA-B trimethyl ester (**1**), ICA-E (**2**), ICA-E pentamethyl ester (**3**), and ICA-G (**4**), together with nine known cryptoporic acids. These isocryptoporic acids are isomers of the cryptoporic acids with drimenol instead of albicanol as the terpenoid fragment; their structures were elucidated on the basis of spectroscopic evidences (UV, IR, HRMS, and NMR) and comparison with literature values. All isolates show certain cytotoxic activities against five tumor cell lines. Among them, compound **4** showed a comparable activity to that of the positive control *cis*-platin, while other compounds exhibited weak cytotoxic activities.

Keywords: *Cryptoporus volvatus*; isocryptoporic acids; cytotoxic activities against cancer cell lines

1. Introduction

Cryptoporus volvatus (Pk.) Hubbard, belonging to the Polyporaceae, grows on living trees or dead wood. The fruiting bodies of this fungus have long been used for the treatment of asthma and bronchitis in Chinese traditional medicine [1]. In previous reports, several cryptoporic acid (CA) derivatives have been isolated from this plant [2–5]. Pharmacological investigations have indicated that this type of compound has strong superoxide-release inhibition [3] and antitumor-promotion properties [6]. In order to find other constituents with higher cytotoxic activities, we examined the fruiting bodies of *C. volvatus* by the bioassay-guided separation, which led to the isolation of four new chemical components as well as nine known components (CA-A, CA-A trimethyl ester, CA-B, CA-B trimethyl ester, CA-C, CA-C pentamethyl ester, CA-D, CA-D pentamethyl ester, and CA-E pentamethyl ester). In this paper, the isolation, structural elucidation, and cytotoxic properties towards five human cancer cell lines of all fractions and isolates were reported.

2. Results and Discussion

2.1. Identification of New Compounds

Compound **1** was obtained as a yellow oil and the molecular formula was established by MALDI-TOF/TOF-MS as C₂₄H₃₈O₈, with a quasi-molecular ion peak of [M + Na]⁺ at *m/z* 477.2464 (calcd. for C₂₄H₃₈O₈Na, 477.2459). The ¹H-NMR spectrum (Table 1) showed two quaternary methyls at δ_H 0.83 and 0.84, a broad singlet at 5.43 ppm characteristic of a double bond, a methyl group at δ_H 1.74 attached to a double bond, and six signals between 2.5 and 4.1 ppm, indicative of the presence

of heteroatoms in the molecular structure. The ^{13}C -NMR spectrum (Table 2) revealed the presence of a double bond with carbon chemical shifts at δ_{C} 122.5 and 133.4, two carbons attached to oxygen, at δ_{C} 72.0 (CH_2) and 78.7 (CH), and three carbonyl carbons at δ_{C} 170.8, 171.1, and 172.2. The NMR data for **1** were similar to that of isocryptoporic acid H (ICA-H) [7], except for the presence of a hydroxymethyl group in place of one tertiary methyl group in ICA-H, indicating that one of the three tertiary methyl groups of ICA-H was replaced by a hydroxymethyl group. The additional hydroxyl group was found to be linked to C-15, which was determined by the presence of the NOE between H-5 and H-15 as well as HMBC correlations of δ_{H} 1.33 and 1.45 (2H, m, H-3), 0.84 (3H, s, H-14), and δ_{C} 72.0 (C-15). The planar structure of **1** was confirmed by analyses of ^1H - ^1H COSY, HSQC, and HMBC data, where a drimenol instead of an albicanol moiety of cryptoporic acid B trimethyl ester [3] is present in the molecule structure. The relative configuration of **1** was established by NOESY analyses, where the cross-peak network observed between 11- CH_3 /13- CH_3 /14- CH_3 /1'- CH confirmed that they were located on the same ring face. The absolute configurations of the isocitric acid moiety were proposed to be either (1'R, 2'S) or (1'S, 2'R) by comparing ^1H -NMR spectral analysis (4.09 (H-1') and 3.45 (H-2')) to the four synthesized diastereoisomeric cryptoporic acid [8]. Comparison of the optical rotation with that of cryptoporic acid B trimethyl ester [3], the absolute configuration of **1** were finally confirmed as 1'R, 2'S (Figure 1). Compound **1** was then named isocryptoporic acid trimethyl ester (ICA-B trimethyl ester).




Figure 1. The structure of compounds **1**–**4**.

Compound **2** was obtained as a yellow oil. The molecular formula was established as $\text{C}_{45}\text{H}_{68}\text{O}_{15}$ on the basis of MALDI-TOF/TOF-MS, which showed a quasimolecular-ion peak ($[\text{M} + \text{Na}]^+$) at m/z 871.4446. The ^1H -NMR and ^{13}C -NMR spectra (Tables 1 and 2) showed that the molecular contained four quaternary methyls, one methyl group attached to a double bond, three methoxys, one exocyclic methylene, three methylenes bearing an oxygen, four ester carbonyls, and two carboxylic groups (Tables 1 and 2). The ^1H - ^1H and ^{13}C - ^1H 2D COSY spectra of **2** displayed the presence of signals corresponding to ICA-B and CA-B, indicating that compound **2**, similar to CA-E [3], is the dimer of two subunits, esterified between the hydroxyl group at C-15 of CA-B and one of the carboxylic acid groups in the isocitric moiety of ICA-B. The position of two carboxylic acids at C-3' and C-3''' and of dimerization at C-5' with the primary alcohol of ICA-B were further established by the 2D NMR spectra of **2**. The relative and absolute configurations of C-1', C-1''', C-2', and C-2''' in the isocitrate moieties were determined to be (1'R, 1'''R, 2'S, 2'''S), in the same way described for compound **1**. On the basis of the above data, the structure of **2** was determined and named isocryptoporic acid E (ICA-E).

Compound **3** was isolated as a yellow oil. Its molecular formula $C_{47}H_{72}O_{15}$ was inferred from the MALDI-TOF/TOF-MS ($[M + Na]^+$) at m/z 899.4776, 28 mass more than that of **2**, which suggests that **3** should be an ester derivant of **2**. In addition, the 1H - and ^{13}C -NMR spectra of **3** further confirmed the deduction, which were quite similar to those of **2**, except for the additional signal of two methoxyl groups. Furthermore, two carboxylic groups at C-3' and C-3''' esterified into two methyl esters were indicated by ^{13}C - 1H 2D COSY and HMBC spectra. The absolute configuration of **3** was established as that of **2**, which was identical to that of **2**. Accordingly, the structure of **2** was assigned as shown and was named ICA-E pentamethyl ester.

Table 1. The 1H -NMR spectral data for compounds **1–4** (500 MHz, $CDCl_3$).

H	1	2	3	4
1	1.03 (m), 1.93 (m)	1.03 (m), 1.88 (m)	1.03 (m), 1.84 (m)	1.03 (m), 1.88 (m)
2	1.51 (m), 1.62 (m)	1.50 (m), 1.63 (m)	1.48 (m), 1.63 (m)	1.50 (m), 1.63 (m)
3	1.33 (m), 1.45 (m)	1.34 (m), 1.42 (m)	1.33 (m), 1.42 (m)	1.32 (m), 1.42 (m)
5	1.48 (m)	1.52 (m)	1.52 (m)	1.53 (m)
6	1.82 (m)	1.88 (m)	1.86 (m)	1.86 (m)
7	5.43 (brs)	5.42 (brs)	5.41 (brs)	5.44 (brs)
9	1.95 (m)	1.94 (m)	1.93 (m)	1.94 (m)
11	3.43 (dd, 9.8, 3.2)	3.46 (m)	3.43 (m)	3.48 (m)
	3.80 (dd, 9.8, 6.0)	3.84 (m)	3.86 (m)	3.82 (m)
12	1.87 (brs)	1.73 (brs)	1.70 (brs)	1.75 (brs)
13	0.83 (s)	0.84 (s)	0.83 (s)	0.84 (s)
14	0.84 (s)	0.84 (s)	0.84 (s)	0.85 (s)
15	3.11 (d, 10.8)	3.12 (d, 11.2)	3.11 (d, 11.2)	3.12 (d, 11.2)
	3.35 (d, 10.8)	3.35 (d, 11.2)	3.35 (d, 11.2)	3.37 (d, 11.2)
1'	4.09 (d, 4.2)	4.10 (d, 4.8)	4.08 (d, 4.8)	4.11 (d, 4.8)
2'	3.45 (m)	3.41 (m)	3.43 (m)	3.43 (m)
3'	2.53 (dd, 16.8, 4.4)	2.64 (m)	2.55 (m)	2.64 (m)
	2.80 (dd, 16.8, 9.2)	2.82 (m)	2.78 (m)	2.80 (m)
1''		1.13 (m), 1.66 (m)	1.15 (m), 1.68 (m)	1.12 (m), 1.65 (m)
2''		1.52 (m), 1.63 (m)	1.50 (m), 1.63 (m)	1.50 (m), 1.66 (m)
3''		1.21 (m), 1.58 (m)	1.23 (m), 1.58 (m)	1.20 (m), 1.58 (m)
5''		1.34 (m)	1.31 (m)	1.35 (m)
6''		1.35 (m), 1.57 (m)	1.33 (m), 1.55 (m)	1.34 (m), 1.55 (m)
7''		2.00 (m), 2.38 (m)	2.01 (m), 2.36 (m)	2.01 (m), 2.38 (m)
9''		1.99 (m)	1.96 (m)	2.03 (m)
11''		3.53 (m)	3.51 (m)	3.55 (m)
		3.92 (m)	3.89 (m)	3.94 (m)
12''		4.78 (brs), 4.84 (brs)	4.75 (brs), 4.84 (brs)	4.78 (brs), 4.86 (brs)
13''		0.74 (s)	0.75 (s)	0.75 (s)
14''		0.75 (s)	0.74 (s)	0.76 (s)
15''		3.62 (d, 11.2)	3.65 (d, 11.2)	3.45 (d, 11.2)
		4.06 (d, 11.2)	3.86 (d, 11.2)	3.96 (d, 11.2)
1'''		4.14 (d, 4.8)	4.10 (d, 4.8)	4.18 (d, 4.8)
2'''		3.40 (m)	3.42 (m)	3.38 (m)
3'''		2.72 (m)	2.53 (m)	2.60 (m)
		2.75 (m)	2.78 (m)	2.68 (m)
OMe	3.65 (s)	3.68 (s)	3.67 (s)	3.75 (s)
	3.65 (s)	3.76 (s)	3.67 (s)	3.76 (s)
	3.68 (s)	3.76 (s)	3.68 (s)	
			3.74 (s)	
			3.75 (s)	

Table 2. The ^{13}C -NMR spectral data for compounds **1–4** (125 MHz, CDCl_3).

C	1	2	3	4
1	39.1	38.8	39.0	38.7
2	18.1	18.5	18.3	18.5
3	35.5	35.4	35.4	35.3
4	35.5	37.4	37.4	37.8
5	44.5	44.6	44.6	44.7
6	23.4	23.4	23.4	23.4
7	122.5	122.5	122.5	123.0
8	133.4	133.2	133.4	132.9
9	54.8	54.9	54.7	55.7
10	37.5	38.5	38.5	38.5
11	69.5	68.9	69.7	68.4
12	21.7	21.8	21.8	21.8
13	15.2	15.7	15.7	15.7
14	17.7	17.6	17.5	17.6
15	72.0	71.8	71.9	71.5
1'	78.7	78.4	78.5	77.7
2'	43.3	43.2	43.2	43.4
3'	31.9	32.0	32.1	32.3
4'	171.1	171.1	171.2	172.6
5'	170.8	170.7	170.8	170.8
6'	172.2	175.0	172.2	174.9
1''		38.5	38.5	38.5
2''		18.1	18.0	18.4
3''		35.3	35.6	35.3
4''		37.6	37.1	37.8
5''		48.9	48.9	48.3
6''		23.6	23.6	23.6
7''		36.8	36.7	36.8
8''		146.3	146.3	146.5
9''		55.1	54.8	55.3
10''		38.9	39.0	38.7
11''		68.2	68.2	68.4
12''		108.2	108.2	108.2
13''		15.6	15.6	15.6
14''		17.7	17.6	17.7
15''		73.6	73.8	73.2
1'''		78.8	78.9	78.8
2'''		44.6	44.4	44.7
3'''		31.7	31.7	32.0
4'''		170.9	170.8	171.1
5'''		171.1	170.7	175.0
6'''		175.2	172.1	176.0
OMe	52.0	52.1	51.9	52.2
	52.0	52.2	51.9	52.2
	52.2	52.2	52.0	
			52.0	
			52.0	

Compound **4** was assigned the molecular formula $\text{C}_{44}\text{H}_{66}\text{O}_{15}$ by analysis of its ^{13}C -NMR and MALDI-TOF/TOF-MS (m/z 833.4343, $[\text{M} - \text{H}]^+$). Expect for the absence of one methoxyl group, the ^1H - and ^{13}C -NMR spectra (Tables 1 and 2) were very close to those of **2**, indicating that **3** is a monode-methyl derivative of **2**. The position of being demethylated was confirmed for the methyl ester at C-2'' by the HMBC cross-peaks $\text{MeO-C}(4')/\text{C-4'}$, $\text{MeO-C}(4'')/\text{C-4''}$, $\text{H-C}(2')/\text{C-4'}$, C-5', and C-6'. In fact, the structure of **4** is the isomer of the known compound CA-G. According to the previous judgment method, the absolute configuration of **4** presented the same conclusion as that of the above compounds. Thus, the structure was finally characterized to be **4** and named isocryptoporic acid G (ICA-G).

2.2. Cytotoxic Activities

The cytotoxic activities of different crude extracts (see Table 3) were assessed in vitro by the MTS cell proliferation assay against a panel of five human cancer cell lines. The activity of the EtOAc extract was similar to that of the MeOH extract, but the extraction rate was higher. In addition, both these extracts displayed better cytotoxic activity than the 60% EtOH extract. Therefore, we finally used ethyl acetate as the extraction solvent. The EtOAc extract was chromatographed on silica gel CC to yield eight fractions. All fractions were further screened via MTS assay. Fraction 1 and Fraction 2, containing a large number of fatty acids and cholesterol, showed weak cytotoxic activities, while Fractions 3–6 with certain cytotoxic activities were finally proved to be consisted mostly of sesquiterpene acids, one of which had been reported to have certain cytotoxic activities. With the eluent polarity increased, more pigment-like impurities were eluted. As a result, the cytotoxic activities of these elutions were decreased. Further bioassay-guided separation of Fractions 3–6 yielded 13 sesquiterpene acids. All isolates were tested for their cytotoxic activity against five human cancer cell lines. *Cis-platin* was a broad spectrum first-line anti-tumor drug and was used as a positive control. Among these compounds, compound 4 showed comparable activity to that of the positive control *cis-platin* (see Table 4), while other compounds exhibited weak cytotoxic activities with IC_{50} values of 50–100 μ M (data not shown). It is concluded that the ratio of carboxyl groups and ester groups may play an important role in attaining better and more selective activity. Further analysis of the structure–activity relationship requires more compounds of this type.

Table 3. Cytotoxic activities of column chromatography fractions.

Samples	IC_{50} (μ g/mL)				
	HL-60	A-549	SMMC-7721	MCF-7	SW480
The EtOAc extract	>200	>300	>300	>300	>300
The MeOH extract	>200	>300	>300	>300	>300
The 60% EtOH-H ₂ O extract	>300	>500	>400	>400	>400
Fraction 1	>300	>400	>400	>400	>400
Fraction 2	>200	>300	>200	>200	>200
Fraction 3	80.81 \pm 1.90	>100	95.19 \pm 2.80	>100	87.82 \pm 0.50
Fraction 4	70.22 \pm 1.70	>100	88.97 \pm 1.54	92.14 \pm 1.90	80.13 \pm 3.30
Fraction 5	67.35 \pm 1.30	94.28 \pm 1.35	85.53 \pm 0.98	90.62 \pm 1.50	85.67 \pm 1.30
Fraction 6	80.87 \pm 2.90	>100	>100	>100	97.14 \pm 1.80
Fraction 7	98.26 \pm 1.90	>100	>100	>100	>100
Fraction 8	>100	>300	>200	>200	>200

Table 4. Antitumor activities of compound 4 and the positive control *cis-platin*.

Compounds	IC_{50} (μ M)				
	HL-60	A-549	SMMC-7721	MCF-7	SW480
4	20.92 \pm 1.20	26.91 \pm 1.80	21.19 \pm 2.10	26.62 \pm 1.60	17.82 \pm 0.70
<i>Cis-platin</i>	1.68 \pm 0.22	25.22 \pm 1.15	14.85 \pm 0.82	11.77 \pm 2.28	4.28 \pm 1.51

3. Experimental Section

3.1. General Procedures

The optical rotations of the compounds were measured with a JASCO P-1020 DIP digital polarimeter (Jasco, Tokyo, Japan). The IR spectra were recorded on a Bruker Tensor-27 spectrometer with KBr pellets (Bruker, Bremen, Germany). The UV spectra were recorded on a Shimadzu UV-2401 PC spectrometer (Shimadzu, Suzhou, China). The 1D- and 2D-NMR spectra were obtained using an Avance III-500 instruments (Bruker) with TMS as the internal standard. The ESI-MS experiment

was performed on Bruker HTC/Esquire mass spectrometer. MADLI-TOF/TOF-MS was carried out with an Bruker Daltonics Flex Analysis spectrometer (Bruker). An Agilent 1100 series machine (NO. 2008052910000035, Agilent, Beijing, China) equipped with a semi-preparative Agilent Zorbax SB-C18 column (9.4 mm × 250 mm, 5 mm, Beijing, China) was used for sample preparation. Column chromatography was performed using silica gel (200–300 mesh, Qingdao Haiyang Chemical Co., Ltd., Qingdao, China), Sephadex LH-20 (GE Healthcare Bio-Sciences AB, Fairfield, CT, USA), and reversed-phase C18 silica gel (50 mm, Merck, Darmstadt, Germany). Analytical and preparative TLC were performed using precoated GF₂₅₄ plates (0.25 mm thickness, Qingdao Haiyang Chemical Co., Ltd., Qingdao, China). The detection was performed by spraying the plates with 5% sulfuric acid followed by heating.

3.2. Plant Material

The fruiting bodies of *Cryptoporus volvatus* (Basidiospores dimensions: 10~11 μm × 5~5.5 μm) was collected in Lijiang, Yunnan Province, China, in April 2015 and identified by Doctor Ying Zhang of Southwest Forestry University, a specialist in the field of fungal identification. A voucher specimen (No. YK01) is deposited at College of Forestry, Southwest Forestry University, Kunming, China.

3.3. Extraction and Isolation

The fruiting bodies of *C. volvatus* (10 kg) were dried by the sun and then cut with a knife. Small pieces of the material were extracted three times with EtOAc (15 L each time) under 60 °C for 2 h, and the combined solution was concentrated to produce a crude extract (2.1 kg). The extract was chromatographed on silica gel (5 kg) using a petroleum ether–ethylacetate gradient (1:0 to 0:1) to yield 8 fractions (Fractions 1~8). Fraction 1 (300 g) was composed of fatty acids. Fraction 2 (250 g) consisted of ergosterol derivatives. Fraction 3 (200 g) was further chromatographed on silica gel (1 kg) using petroleum ether–ethylacetate (8:1 to 4:1) to furnish compound 1 (15 mg). Fraction 4 (100 g) was subjected to further silica gel CC, eluted with petroleum ether–acetone (6:1 to 1:1) to afford four fractions (Fractions 4-A~D). Fraction 4-C (10 g) was rechromatographed on Sephadex LH-20 CC (CHCl₃–EtOH = 7:3) to afford the mixtures (4 g) of drimane-type sesquiterpenoids, which were further purified by semi-HPLC (CH₃CN:H₂O = 65:35, λ = 210 nm) to produce compound 2 (12.5 mg, Retention time = 8.5 min) and compound 3 (10 mg, Retention time = 10.3 min). Fraction 5 (150 g) was fractionated by MCI gel CC, eluted with MeOH–H₂O (6:4 to 1:0) to yield 3 fractions (Fractions 5-A~D). Fraction 5-B (10 g) was purified using silica gel CC and eluted with petroleum ether–acetone (2:1 to 1:2) to yield compound 4 (14 mg).

Isocryptoporic acid B trimethyl ester (1). Yellow oil; UV (MeOH) ν_{\max} (log ϵ): 204 (0.27) nm; $[\alpha]_{\text{D}}^{20} +36$ (c 2.10, MeOH); IR: 3431, 2950, 2929, 1742, 1663, 1044 cm^{−1}; ¹H-NMR (500 MHz, CHCl₃) and ¹³C-NMR (125 MHz, CHCl₃) data, see Table 1; MALDI-TOF/TOF-MS (m/z): 477.2464 [M + Na]⁺ (calcd. for C₂₄H₃₈O₈Na, 477.2459).

Isocryptoporic acid E (2). Yellow oil; UV (MeOH) ν_{\max} (log ϵ): 203 (0.26) nm; $[\alpha]_{\text{D}}^{20} +23$ (c 2.20, MeOH); IR: 3431, 2946, 2929, 1739, 1640, 1040 cm^{−1}; ¹H-NMR (500 MHz, CHCl₃) and ¹³C-NMR (125 MHz, CHCl₃) data, see Table 1; MALDI-TOF/TOF-MS (m/z): 871.4446 [M + Na]⁺ (calcd. for C₄₅H₆₈O₁₅Na, 871.4439).

Isocryptoporic acid E pentamethyl ester (3). Yellow oil; UV (MeOH) ν_{\max} (log ϵ): 203 (0.22) nm; $[\alpha]_{\text{D}}^{20} +24$ (c 2.20, MeOH); IR: 3433, 2945, 2930, 1740, 1644, 1044 cm^{−1}; ¹H-NMR (500 MHz, CHCl₃) and ¹³C-NMR (125 MHz, CHCl₃) data, see Table 1; MALDI-TOF/TOF-MS (m/z): 899.4776 [M + Na]⁺ (calcd. for C₄₇H₇₂O₁₅Na, 899.4763).

Isocryptoporic acid G (4). Yellow oil; UV (MeOH) ν_{\max} (log ϵ): 203 (0.35) nm; $[\alpha]_{\text{D}}^{20} +21$ (c 1.70, MeOH); IR: 3426, 2946, 2929, 1739, 1638, 1034 cm^{−1}; ¹H-NMR (500 MHz, CHCl₃) and ¹³C-NMR (125 MHz, CHCl₃) data, see Table 1; MALDI-TOF/TOF-MS (m/z): 833.4343 [M − H]⁺ (calcd. for C₄₄H₆₆O₁₅, 833.4329).

3.4. Assays of Cytotoxic Activity

Five human cancer cell lines, i.e., lung cancer (A-549), human myeloid leukemia (HL-60), breast cancer (MCF-7), hepatocellular carcinoma (SMMC-7721), and colon cancer (SW-480) were purchased from American Type Culture Collection (ATCC) (Manassas, VA, USA) and cultured in RPMI 1640 medium (HL-60 and SMMC-7721) or DMEM medium (A-549, MCF-7 and SW-480). The cytotoxic activities of tumor cell lines were performed in vitro by the MTS cell proliferation assay. Briefly, the cells were seeded at a concentration of 5×10^3 cell/mL in a volume of 0.1 mL in 96-well plates and incubated for 24 h at 37 °C in a CO₂ incubator. Test sample portions of 100 µL at varied concentrations, prepared in culture medium, were added to each well. After 48 h of incubation at 37 °C in the CO₂ incubator, the cells in each well were incubated in culture medium with 20 µL of a 5 M solution of MTS, which was purchased from Promega Biotechnology Corporation (Beijing, China) and dissolved in PBS (pH = 6.0), for 1 h at 37 °C. After another hour of incubation, absorption was measured at 490 nm via MULTISKAN FC. All experiments were carried out in triplicate. The cytotoxicity of each sample was expressed as IC₅₀. The inhibition ratio of cells proliferation was calculated as follows [9]:

$$\text{Inhibition ratio (\%)} = (1 - (\text{OD} - \text{OD}_{\text{blank}})/(\text{OD}_0 - \text{OD}_{\text{blank}})) \times 100\% \quad (1)$$

where OD, OD₀, and OD_{blank} are the absorbance of treated cells, untreated cells, and the blank control, respectively.

4. Conclusions

In this study, the EtOAc extract of *Cryptoporus volvatus* (Pk.) Hubbard was chromatographed to get different fractions. On the basis of the evaluation in vitro cytotoxic activity using the MTS assay to all fractions, the bioassay-guided separation led to the isolation of four new isocryptoporic acid derivatives, together with nine known cryptoporic acids. This kind of isocryptoporic acid was reported in this genus for the first time. In addition, the activity data indicated that all compounds had certain inhibitory activities against five cancer cells. Compound 4 especially had moderate inhibitory activities, exhibiting close inhibitions with the positive control *cis*-platin.

Acknowledgments: The authors are grateful to the members of the State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, for measuring the spectroscopic data and cytotoxic activities.

Author Contributions: L.-Y.Z. and Y.H. designed research; L.-Y.Z., X.-H.Y., and B.L. performed the isolation and purification of plants, and analyzed the data; L.-Y.Z. wrote the paper. All authors read and approved the final manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Zhao, J.D. *Flora Fungorum Sinicorum*; Science Press: Beijing, China, 1998; pp. 88–89.
2. Hirotsu, M.; Furuya, T.; Shiro, M. Cryptoporic acid-H and acid-I, drimane-sesquiterpenes from *ganoderma-neo-japonicum* and *Cryptoporus volvatus*. *Phytochemistry* **1991**, *30*, 1555–1559. [[CrossRef](#)]
3. Asakawa, Y.; Hashimoto, T.; Mizuno, Y.; Toria, M.; Fukazawa, Y. Cryptoporic acids AG, drimane-type sesquiterpenoid ethers of isocitric acid from the fungus *Cryptoporus volvatus*. *Phytochemistry* **1992**, *31*, 579–592. [[CrossRef](#)]
4. Wang, J.C.; Li, G.Z.; Gao, L.; Cao, L.; Lv, N.; Shen, L.G.; Si, J.Y. Two new cryptoporic acid derivatives from the fruiting bodies of *Cryptoporus volvatus*. *Phytochem. Lett.* **2015**, *14*, 64–66. [[CrossRef](#)]
5. Wang, J.C.; Li, G.Z.; Lv, N.; Gao, L.; Cao, L.; Shen, L.G.; Si, J.Y. Chemical constituents from the fruiting bodies of *Cryptoporus volvatus*. *Arch. Pharm. Res.* **2016**, *39*, 747–754. [[CrossRef](#)] [[PubMed](#)]
6. Wu, W.; Zhao, F.; Ding, R.; Bao, L.; Gao, H.; Lu, J.C.; Yao, X.S.; Zhang, X.Q.; Liu, H.W. Four New Cryptoporic Acid Derivatives from the Fruiting Bodies of *Cryptoporus sinensis*, and Their Inhibitory Effects on Nitric Oxide Production. *Chem. Biodivers.* **2011**, *8*, 1529–1538. [[CrossRef](#)]

7. Xu, X.B.; Gu, Z.X.; Liu, S.; Gao, N.; He, X.Z.; Xin, X. Purification and characterization of a glucan from *Bacillus Calmette Guerin* and the antitumor activity of its sulfated derivative. *Carbohydr. Polym.* **2015**, *128*, 138–146. [[CrossRef](#)] [[PubMed](#)]
8. Tori, M.; Hamada, N.; Sono, M.; Sono, Y.; Ishikawa, M.; Nakashima, K.; Hashimoto, T.; Asakawa, Y. Synthesis of cryptoporic acid A methyl ester. *Tetrahedron Lett.* **2000**, *41*, 3099–3102. [[CrossRef](#)]
9. Huang, K.Y.; Kao, S.H.; Wang, W.L.; Chen, C.Y.; Hsiao, T.H.; Salunke, S.B.; Chen, J.W.; Su, C.K.; Yang, S.C.; Hong, T.M.; et al. Small Molecule T315 Promotes Casitas B-Lineage Lymphoma-Dependent Degradation of Epidermal Growth Factor Receptor via Y1045 Autophosphorylation. *Am. J. Respir. Crit. Care Med.* **2016**, *193*, 753–766. [[CrossRef](#)] [[PubMed](#)]

Sample Availability: Samples of the compounds are not available from the authors because of the limited amount.



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).