

Investigation of ‘Head-to-Tail’-Connected Oligoaryl N,O-Ligands as Recognition Motifs for Cancer-Relevant G-Quadruplexes

Natalia Rizeq¹ and Savvas N. Georgiades^{1,*}



¹ Department of Chemistry, University of Cyprus, 1 Panepistimiou Avenue,
Aglandjia, 2109, Nicosia, Cyprus



* Correspondence: georgiades.savvas@ucy.ac.cy; Tel.: +357-22-892779



Supplementary Materials

Table of Contents



• ^1H and ^{13}C NMR spectra	S-2
• Fluorescence titration curves	S-13
• FRET melting temperature curves	S-16
• ΔT_m for Fds26T	S-22













S-4











S-6


















22AG (telomeric)/K⁺/5-mer (**10**):




22AG (telomeric)/K⁺/6-mer (**13**):




22AG (telomeric)/K⁺/7-mer (**14**):




22AG (telomeric)/Na⁺/5-mer (**10**):




22AG (telomeric)/Na⁺/6-mer (**13**):




22AG (telomeric)/Na⁺/7-mer (**14**):




Myc2345-Pu22/K⁺/5-mer (**10**):




Myc2345-Pu22/K⁺/6-mer (**13**):




Myc2345-Pu22/K⁺/7-mer (**14**):




F21T/K⁺/7mer (**14**)/ds26:




F21T/K⁺/6mer (**13**)/ds26:




F21T/K⁺/5mer (**10**)/ds26:




F21T/Na⁺/7mer (**14**)/ds26:




F21T/ Na^+ /6mer (**13**)/ds26:




F21T/ Na^+ /5mer (**10**)/ds26:




FmycT/K⁺/7mer (**14**)/ds26:




FmycT/K⁺/6mer (**13**)/ds26:




FmycT/K⁺/5mer (**10**)/ds26:



Fds26T/K⁺/7mer (**14**):



Fds26T/K⁺/6mer (**13**):



Fds26T/K⁺/5mer (**10**):

