

Supplementary Data

Studies on the inclusion complexes of daidzein with β -cyclodextrin and derivatives

Shujing Li^{1*}, Li Yuan¹, Yong Chen², Wei Zhou¹ and Xinrui Wang¹

¹ Beijing Advanced Innovation Center for Food Nutrition and Human Health/Department of Chemistry,
School of Science, Beijing Technology and Business University, Beijing 100048, PR China;

² Key Laboratory of Photochemical Conversion and Optoelectronic Materials, Technical Institute of
Physics and Chemistry, Chinese Academy of Sciences, Beijing 100190, PR China

* Correspondence: lishujing@mail.ipc.ac.cn; Tel.: +86-10-6898-5573




Figure S1 Job's continuous variation plot of the daidzein- β -CD complex




Figure S2 Job's continuous variation plot of the daidzein-Me- β -CD complex




Figure S3 Job's continuous variation plot of the daidzein-HP- β -CD complex




Figure S4 TG curve of daidzein, β -CD, daidzein/ β -CD physical mixture, and daidzein- β -CD inclusion complex.




Figure S5 TG curve of daidzein, Me- β -CD, daidzein/Me- β -CD physical mixture, and daidzein-Me- β -CD inclusion complex.




Figure S6 TG curve of daidzein, HP- β -CD, daidzein/HP- β -CD physical mixture, daidzein-HP- β -CD inclusion complex.




Figure S7 Scanning electron microphotographs: (A) daidzein, (B) β -CD, (C) daidzein/ β -CD physical mixture, (D) daidzein- β -CD inclusion complex.




Figure S8 Scanning electron microphotographs: (A) daidzein, (B) Me- β -CD, (C) daidzein/Me- β -CD physical mixture, (D) daidzein-Me- β -CD inclusion complex.




Figure S9 ^1H NMR spectra of β -CD in D_2O at 25°C




Figure S10 ^1H NMR spectra of daidzein- β -CD inclusion complex in D_2O at 25°C




Figure S11 ^1H NMR spectra of Me- β -CD in D_2O at 25°C




Figure S12 ^1H NMR spectra of daidzein-Me- β -CD inclusion complex in D_2O at 25°C




Figure S13 ^1H NMR spectra of HP- β -CD in D_2O at 25°C




Figure S14 ^1H NMR spectra of daidzein-HP- β -CD inclusion complex in D_2O at 25 $^\circ\text{C}$




Figure S15 ROESY spectrum of daidzein- β -CD inclusion complex in D_2O at 25 $^\circ\text{C}$




Figure S16 ROESY spectrum of daidzein-Me- β -CD inclusion complex in D_2O at 25 $^\circ\text{C}$